

Classification overview: proteins



All 1,900 proteins, one row each. Rows are ordered by average−linkage clustering (1 − Pearson r) of limpa's matrix, for display only. Median change colours stop at the 99th percentile of |change|, 1. Grey: not tested. Table: c_data/01_proteins.xlsx, auc.

Training, Hi (T1 to T2)


229 of 1900 proteins at p < 0.05, 0 at BH < 0.05. Page 1 of 2.



Wilcoxon signed−rank test on up to 7 pairs; smallest attainable p 0.016. Header: median paired change, p, BH q within task, and the feature's complete pairs. Table: c_data/01_proteins.xlsx, auc.

Training, Hi (T1 to T2)


229 of 1900 proteins at p < 0.05, 0 at BH < 0.05. Page 2 of 2.



Wilcoxon signed−rank test on up to 7 pairs; smallest attainable p 0.016. Header: median paired change, p, BH q within task, and the feature's complete pairs. Table: c_data/01_proteins.xlsx, auc.

Training, Lo (T1 to T2)


75 of 1900 proteins at p < 0.05, 0 at BH < 0.05. Page 1 of 2.



Wilcoxon signed−rank test on up to 6 pairs; smallest attainable p 0.031. Header: median paired change, p, BH q within task, and the feature's complete pairs. Table: c_data/01_proteins.xlsx, auc.

Training, Lo (T1 to T2)


75 of 1900 proteins at p < 0.05, 0 at BH < 0.05. Page 2 of 2.



Wilcoxon signed−rank test on up to 6 pairs; smallest attainable p 0.031. Header: median paired change, p, BH q within task, and the feature's complete pairs. Table: c_data/01_proteins.xlsx, auc.

Acute bout, Hi (T2 to T3)


200 of 1900 proteins at p < 0.05, 0 at BH < 0.05. Page 1 of 2.



Wilcoxon signed−rank test on up to 7 pairs; smallest attainable p 0.016. Header: median paired change, p, BH q within task, and the feature's complete pairs. Table: c_data/01_proteins.xlsx, auc.

Acute bout, Hi (T2 to T3)


200 of 1900 proteins at p < 0.05, 0 at BH < 0.05. Page 2 of 2.



Wilcoxon signed−rank test on up to 7 pairs; smallest attainable p 0.016. Header: median paired change, p, BH q within task, and the feature's complete pairs. Table: c_data/01_proteins.xlsx, auc.

Acute bout, Lo (T2 to T3)


106 of 1900 proteins at p < 0.05, 0 at BH < 0.05. Page 1 of 2.



Wilcoxon signed−rank test on up to 7 pairs; smallest attainable p 0.016. Header: median paired change, p, BH q within task, and the feature's complete pairs. Table: c_data/01_proteins.xlsx, auc.

Acute bout, Lo (T2 to T3)


106 of 1900 proteins at p < 0.05, 0 at BH < 0.05. Page 2 of 2.



Wilcoxon signed−rank test on up to 7 pairs; smallest attainable p 0.016. Header: median paired change, p, BH q within task, and the feature's complete pairs. Table: c_data/01_proteins.xlsx, auc.

Hi vs Lo at T1


121 of 1900 proteins at p < 0.05, 0 at BH < 0.05. Page 1 of 2.



Hi (up to 8) against Lo (up to 6); AUC above 0.5 means higher in Hi. Wilcoxon rank−sum test; smallest attainable p 0.00067. Header: AUC with its DeLong 95% interval, p, none at AUC 0 or 1, where DeLong's variance is ze

Hi vs Lo at T1


121 of 1900 proteins at p < 0.05, 0 at BH < 0.05. Page 2 of 2.



Hi (up to 8) against Lo (up to 6); AUC above 0.5 means higher in Hi. Wilcoxon rank−sum test; smallest attainable p 0.00067. Header: AUC with its DeLong 95% interval, p, none at AUC 0 or 1, where DeLong's variance is ze

Hi vs Lo at T2


76 of 1900 proteins at p < 0.05, 0 at BH < 0.05. Page 1 of 2.



Hi (up to 7) against Lo (up to 7); AUC above 0.5 means higher in Hi. Wilcoxon rank−sum test; smallest attainable p 0.00058. Header: AUC with its DeLong 95% interval, p, none at AUC 0 or 1, where DeLong's variance is ze

Hi vs Lo at T2


76 of 1900 proteins at p < 0.05, 0 at BH < 0.05. Page 2 of 2.



Hi (up to 7) against Lo (up to 7); AUC above 0.5 means higher in Hi. Wilcoxon rank−sum test; smallest attainable p 0.00058. Header: AUC with its DeLong 95% interval, p, none at AUC 0 or 1, where DeLong's variance is ze

Hi vs Lo, training change


123 of 1900 proteins at p < 0.05, 0 at BH < 0.05. Page 1 of 2.



Hi (up to 7) against Lo (up to 6); AUC above 0.5 means higher in Hi. Wilcoxon rank−sum test; smallest attainable p 0.0012. Header: AUC with its DeLong 95% interval, p, none at AUC 0 or 1, where DeLong's variance is zero

Hi vs Lo, training change


123 of 1900 proteins at p < 0.05, 0 at BH < 0.05. Page 2 of 2.



Hi (up to 7) against Lo (up to 6); AUC above 0.5 means higher in Hi. Wilcoxon rank−sum test; smallest attainable p 0.0012. Header: AUC with its DeLong 95% interval, p, none at AUC 0 or 1, where DeLong's variance is zero

Hi vs Lo, acute change


26 of 1900 proteins at p < 0.05, 0 at BH < 0.05. Page 1 of 2.



Hi (up to 7) against Lo (up to 7); AUC above 0.5 means higher in Hi. Wilcoxon rank−sum test; smallest attainable p 0.00058. Header: AUC with its DeLong 95% interval, p, none at AUC 0 or 1, where DeLong's variance is ze

Hi vs Lo, acute change

26 of 1900 proteins at p < 0.05, 0 at BH < 0.05. Page 2 of 2.



Hi (up to 7) against Lo (up to 7); AUC above 0.5 means higher in Hi. Wilcoxon rank−sum test; smallest attainable p 0.00058. Header: AUC with its DeLong 95% interval, p, none at AUC 0 or 1, where DeLong's variance is ze