

Discrimination overview: proteins



All 1,900 proteins, one row each. Rows are ordered by average−linkage clustering (1 − Pearson r) of limpa's matrix, for display only. Grey: not tested. Table: c_data/01_proteins.xlsx, auc.

HR vs LR at T1


44 of 1900 proteins at p < 0.05, 0 at BH < 0.05. Page 1 of 2.



HR (up to 7) against LR (up to 8); AUC above 0.5 means higher in HR. Header: AUC with its stratified bootstrap 95% interval (2000 resamples), Wilcoxon rank−sum p (smallest attainable 0.00031), BH q within task, and the

HR vs LR at T1


44 of 1900 proteins at p < 0.05, 0 at BH < 0.05. Page 2 of 2.



HR (up to 7) against LR (up to 8); AUC above 0.5 means higher in HR. Header: AUC with its stratified bootstrap 95% interval (2000 resamples), Wilcoxon rank−sum p (smallest attainable 0.00031), BH q within task, and the

HR vs LR at T2


57 of 1900 proteins at p < 0.05, 0 at BH < 0.05. Page 1 of 2.



HR (up to 7) against LR (up to 8); AUC above 0.5 means higher in HR. Header: AUC with its stratified bootstrap 95% interval (2000 resamples), Wilcoxon rank−sum p (smallest attainable 0.00031), BH q within task, and the

HR vs LR at T2


57 of 1900 proteins at p < 0.05, 0 at BH < 0.05. Page 2 of 2.



HR (up to 7) against LR (up to 8); AUC above 0.5 means higher in HR. Header: AUC with its stratified bootstrap 95% interval (2000 resamples), Wilcoxon rank−sum p (smallest attainable 0.00031), BH q within task, and the

HR vs LR at T3


100 of 1900 proteins at p < 0.05, 0 at BH < 0.05. Page 1 of 2.



HR (up to 7) against LR (up to 8); AUC above 0.5 means higher in HR. Header: AUC with its stratified bootstrap 95% interval (2000 resamples), Wilcoxon rank−sum p (smallest attainable 0.00031), BH q within task, and the

HR vs LR at T3


100 of 1900 proteins at p < 0.05, 0 at BH < 0.05. Page 2 of 2.



HR (up to 7) against LR (up to 8); AUC above 0.5 means higher in HR. Header: AUC with its stratified bootstrap 95% interval (2000 resamples), Wilcoxon rank−sum p (smallest attainable 0.00031), BH q within task, and the

HR vs LR, training change


50 of 1900 proteins at p < 0.05, 0 at BH < 0.05. Page 1 of 2.



HR (up to 6) against LR (up to 8); AUC above 0.5 means higher in HR. Header: AUC with its stratified bootstrap 95% interval (2000 resamples), Wilcoxon rank−sum p (smallest attainable 0.00067), BH q within task, and the

HR vs LR, training change


50 of 1900 proteins at p < 0.05, 0 at BH < 0.05. Page 2 of 2.



HR (up to 6) against LR (up to 8); AUC above 0.5 means higher in HR. Header: AUC with its stratified bootstrap 95% interval (2000 resamples), Wilcoxon rank−sum p (smallest attainable 0.00067), BH q within task, and the

HR vs LR, acute change


67 of 1900 proteins at p < 0.05, 0 at BH < 0.05. Page 1 of 2.



HR (up to 7) against LR (up to 8); AUC above 0.5 means higher in HR. Header: AUC with its stratified bootstrap 95% interval (2000 resamples), Wilcoxon rank−sum p (smallest attainable 0.00031), BH q within task, and the

HR vs LR, acute change

67 of 1900 proteins at p < 0.05, 0 at BH < 0.05. Page 2 of 2.



HR (up to 7) against LR (up to 8); AUC above 0.5 means higher in HR. Header: AUC with its stratified bootstrap 95% interval (2000 resamples), Wilcoxon rank−sum p (smallest attainable 0.00031), BH q within task, and the