





All 1,900 proteins by 42 biopsies, grouped by arm and time point; column labels are subjects. Fill: log2 abundance minus the protein's mean over biopsies. Colours stop at the 99th percentile of |value|, 1.9. Rows are ordered by average−linkage clustering (1 − Pearson r) of limpa's matrix, for display only. Grey: missing.

# Contrast overview: proteins





All 1,900 proteins by the nine contrasts. Fill: log2 fold change from the limma fit; status: p and BH q within contrast. Colours stop at the 99th percentile of |value|, 1.1. Rows are ordered by average−linkage clustering (1 − Pearson r) of limpa's matrix, for display only. Grey: not tested. Table: c_data/02_contrasts.xlsx, DEP_matrix.





Dashed line: a flat null (tested / 20). A spike at zero on a flat background is signal; a slope up toward 1 is a conservative test. Table: c_data/02_contrasts.xlsx, DEP_matrix.

# Volcano per contrast

34 to 175 proteins at p < 0.05 per contrast; 0 at BH < 0.05 across all nine.




limma on limpa's quantification, subject blocked, blood as a covariate. Dashed line: p = 0.05. BH within contrast. Table: c_data/02_contrasts.xlsx, DEP_matrix.


Samples on the complete, blood−corrected matrix
PERMANOVA, time within subject: R2 0.062, p = 0.0044
PC2 (10.0%)
25
0
−25
−50
−25
0
25
PC1 (12.9%)
arm
Hi
Lo
timepoint
T1
T2
T3
Proteins scaled to unit variance. Table: c_data/02_contrasts.xlsx, pca, permanova.