


Nominal associations against chance




Chance is 5% of proteins tested per cell. Table: c_data/04_associate.xlsx, chance_expectation.

log2 abundance against fc_{sa}_mixed, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2



log2 abundance against fc_{sa}_mixed, between-person term


One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects

HR LR T1 T2



log2 abundance against fc_{sa}_mixed, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects




67 of 1640 proteins reach p < 0.05 on this term; chance predicts 82. limma on 30 biopsies, subject blocked; q from BH within trait and term. Table: c_data/04_associate.xlsx, model.

log2 abundance against fc_{sa_mixed}, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2



log2 abundance against fcsa_mixed, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2



log2 abundance against fcsc_mixed, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2



log2 abundance against fcsc_mixed, within-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2



log2 abundance against fcsa_mixed, within-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2



log2 abundance against fcsc_mixed, within-person term


One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects

HR LR T1 T2



log2 abundance against fcsc_mixed, within-person term


One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects

61 of 1640 proteins reach $p < 0.05$ on this term; chance predicts 82. limma on 30 biopsies, subject blocked; q from BH within trait and term. Table: c_data/04_associate.xlsx, model.

log2 abundance against fcsa_mixed, within-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects

HR LR T1 T2




log2 abundance against fc_{sa}_mixed, within-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects

HR LR T1 T2


PARVA
within -0.000353 p 0.05 q 0.99



log2 abundance against fcsc_type1, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2



log2 abundance against fcsa_type1, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2



log2 abundance against fc_{sa}_type1, between-person term


One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects

HR LR T1 T2



log2 abundance against fc_{sa}_type1, between-person term


One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects

64 of 1640 proteins reach $p < 0.05$ on this term; chance predicts 82. limma on 30 biopsies, subject blocked; q from BH within trait and term. Table: c_data/04_associate.xlsx, model.

log2 abundance against fc_{sa}_type1, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2



log2 abundance against fcsc_type1, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2



log2 abundance against fcsc_type1, within-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2



log2 abundance against fc_{sa}_type1, within-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects

HR LR ○ T1 ● T2




59 of 1640 proteins reach p < 0.05 on this term; chance predicts 82. limma on 30 biopsies, subject blocked; q from BH within trait and term. Table: c_data/04_associate.xlsx, model.

log2 abundance against fc_{sa_type1}, within-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2



log2 abundance against fcsc_type1, within-person term


One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects

HR LR T1 T2




log2 abundance against fc_{sa_type1}, within-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects




log2 abundance against fcsa_type2, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects

64 of 1640 proteins reach $p < 0.05$ on this term; chance predicts 82. limma on 30 biopsies, subject blocked; q from BH within trait and term. Table: c_data/04_associate.xlsx, model.


log2 abundance against fcsa_type2, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects



log2 abundance against fcsc_type2, between-person term


One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects



64 of 1640 proteins reach p < 0.05 on this term; chance predicts 82. limma on 30 biopsies, subject blocked; q from BH within trait and term. Table: c_data/04_associate.xlsx, model.


log2 abundance against fc_{sa}_type2, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects

64 of 1640 proteins reach $p < 0.05$ on this term; chance predicts 82. limma on 30 biopsies, subject blocked; q from BH within trait and term. Table: c_data/04_associate.xlsx, model.

log2 abundance against fcsa_type2, between-person term


One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects

64 of 1640 proteins reach $p < 0.05$ on this term; chance predicts 82. limma on 30 biopsies, subject blocked; q from BH within trait and term. Table: c_data/04_associate.xlsx, model.

log2 abundance against fc_{sa}_type2, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2



log2 abundance against fc_{sa_type2}, within-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2

74 of 1640 proteins reach $p < 0.05$ on this term; chance predicts 82. limma on 30 biopsies, subject blocked; q from BH within trait and term. Table: c_data/04_associate.xlsx, model.

log2 abundance against fcsc_type2, within-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects

HR LR T1 T2




74 of 1640 proteins reach p < 0.05 on this term; chance predicts 82. limma on 30 biopsies, subject blocked; q from BH within trait and term. Table: c_data/04_associate.xlsx, model.

log2 abundance against fcsc_type2, within-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects

HR LR T1 T2




74 of 1640 proteins reach p < 0.05 on this term; chance predicts 82. limma on 30 biopsies, subject blocked; q from BH within trait and term. Table: c_data/04_associate.xlsx, model.

log2 abundance against fc_{sa}_type2, within-person term


One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects

HR LR T1 T2




log2 abundance against fcsa_type2, within-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects

74 of 1640 proteins reach $p < 0.05$ on this term; chance predicts 82. limma on 30 biopsies, subject blocked; q from BH within trait and term. Table: c_data/04_associate.xlsx, model.

log2 abundance against fc_{sa_type2}, within-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects




74 of 1640 proteins reach p < 0.05 on this term; chance predicts 82. limma on 30 biopsies, subject blocked; q from BH within trait and term. Table: c_data/04_associate.xlsx, model.

log2 abundance against fcsa_type2, within-person term


One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects

HR LR T1 T2



log2 abundance against fibres_mixed, between-person term


One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects

83 of 1640 proteins reach $p < 0.05$ on this term; chance predicts 82. limma on 30 biopsies, subject blocked; q from BH within trait and term. Table: c_data/04_associate.xlsx, model.

log2 abundance against fibres_mixed, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects

HR LR T1 T2




83 of 1640 proteins reach p < 0.05 on this term; chance predicts 82. limma on 30 biopsies, subject blocked; q from BH within trait and term. Table: c_data/04_associate.xlsx, model.

log2 abundance against fibres_mixed, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects

HR LR T1 T2




83 of 1640 proteins reach p < 0.05 on this term; chance predicts 82. limma on 30 biopsies, subject blocked; q from BH within trait and term. Table: c_data/04_associate.xlsx, model.

log2 abundance against fibres_mixed, between-person term


One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects

HR LR T1 T2




log2 abundance against fibres_mixed, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects



log2 abundance against fibres_mixed, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects




83 of 1640 proteins reach p < 0.05 on this term; chance predicts 82. limma on 30 biopsies, subject blocked; q from BH within trait and term. Table: c_data/04_associate.xlsx, model.

log2 abundance against fibres_mixed, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2



log2 abundance against fibres_mixed, within-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2



log2 abundance against fibres_mixed, within-person term


One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects

HR LR T1 T2



log2 abundance against fibres_mixed, within-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects




46 of 1640 proteins reach p < 0.05 on this term; chance predicts 82. limma on 30 biopsies, subject blocked; q from BH within trait and term. Table: c_data/04_associate.xlsx, model.

log2 abundance against fibres_mixed, within-person term


One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects

HR LR T1 T2



log2 abundance against fibres_type1, between-person term


One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects

63 of 1640 proteins reach $p < 0.05$ on this term; chance predicts 82. limma on 30 biopsies, subject blocked; q from BH within trait and term. Table: c_data/04_associate.xlsx, model.

log2 abundance against fibres_type1, between-person term


One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects

HR LR T1 T2



log2 abundance against fibres_type1, between-person term


One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects

63 of 1640 proteins reach $p < 0.05$ on this term; chance predicts 82. limma on 30 biopsies, subject blocked; q from BH within trait and term. Table: c_data/04_associate.xlsx, model.

log2 abundance against fibres_type1, between-person term


One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects

HR LR T1 T2



log2 abundance against fibres_type1, between-person term


One line per subject from T1 to T2: within is the slope of the lines, between is the spread of subjects

63 of 1640 proteins reach $p < 0.05$ on this term; chance predicts 82. limma on 30 biopsies, subject blocked; q from BH within trait and term. Table: c_data/04_associate.xlsx, model.

log2 abundance against fibres_type1, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2



log2 abundance against fibres_type1, within-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2



log2 abundance against fibres_type1, within-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2



log2 abundance against fibres_type1, within-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2



52 of 1640 proteins reach $p < 0.05$ on this term; chance predicts 82. limma on 30 biopsies, subject blocked; q from BH within trait and term. Table: c_data/04_associate.xlsx, model.

log2 abundance against fibres_type1, within-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects




52 of 1640 proteins reach $p < 0.05$ on this term; chance predicts 82. limma on 30 biopsies, subject blocked; q from BH within trait and term. Table: c_data/04_associate.xlsx, model.

log2 abundance against fibres_type1, within-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2



log2 abundance against leg_ext_1rm, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2



log2 abundance against leg_ext_1rm, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects

HR LR T1 T2




130 of 1615 proteins reach $p < 0.05$ on this term; chance predicts 81. limma on 29 biopsies, subject blocked; q from BH within trait and term. Table: c_data/04_associate.xlsx, model.

log2 abundance against leg_ext_1rm, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2



log2 abundance against leg_ext_1rm, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects

HR LR T1 T2




130 of 1615 proteins reach $p < 0.05$ on this term; chance predicts 81. limma on 29 biopsies, subject blocked; q from BH within trait and term. Table: c_data/04_associate.xlsx, model.

log2 abundance against leg_ext_1rm, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects

HR LR T1 T2




130 of 1615 proteins reach $p < 0.05$ on this term; chance predicts 81. limma on 29 biopsies, subject blocked; q from BH within trait and term. Table: c_data/04_associate.xlsx, model.

log2 abundance against leg_ext_1rm, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects

HR LR T1 T2




130 of 1615 proteins reach $p < 0.05$ on this term; chance predicts 81. limma on 29 biopsies, subject blocked; q from BH within trait and term. Table: c_data/04_associate.xlsx, model.

log2 abundance against leg_ext_1rm, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2



log2 abundance against leg_ext_1rm, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2



log2 abundance against leg_ext_1rm, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2



log2 abundance against leg_ext_1rm, between-person term


One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects

HR LR T1 T2

130 of 1615 proteins reach $p < 0.05$ on this term; chance predicts 81. limma on 29 biopsies, subject blocked; q from BH within trait and term. Table: c_data/04_associate.xlsx, model.

log2 abundance against leg_ext_1rm, between-person term


One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


130 of 1615 proteins reach $p < 0.05$ on this term; chance predicts 81. limma on 29 biopsies, subject blocked; q from BH within trait and term. Table: c_data/04_associate.xlsx, model.

log2 abundance against leg_ext_1rm, within-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects

HR LR T1 T2




122 of 1615 proteins reach $p < 0.05$ on this term; chance predicts 81. limma on 29 biopsies, subject blocked; q from BH within trait and term. Table: c_data/04_associate.xlsx, model.

log2 abundance against leg_ext_1rm, within-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2



log2 abundance against leg_ext_1rm, within-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2



log2 abundance against leg_ext_1rm, within-person term

One line per subject from T1 to T2: within is the slope of the lines, between is the spread of subjects


HR LR T1 T2



log2 abundance against leg_ext_1rm, within-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects

HR LR T1 T2




122 of 1615 proteins reach $p < 0.05$ on this term; chance predicts 81. limma on 29 biopsies, subject blocked; q from BH within trait and term. Table: c_data/04_associate.xlsx, model.

log2 abundance against leg_ext_1rm, within-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2



log2 abundance against leg_ext_1rm, within-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2



log2 abundance against leg_ext_1rm, within-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects

HR LR T1 T2




122 of 1615 proteins reach $p < 0.05$ on this term; chance predicts 81. limma on 29 biopsies, subject blocked; q from BH within trait and term. Table: c_data/04_associate.xlsx, model.

log2 abundance against leg_ext_1rm, within-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2



log2 abundance against leg_ext_1rm, within-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2



log2 abundance against leg_ext_1rm, within-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2



log2 abundance against leg_press_1rm, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2



log2 abundance against leg_press_1rm, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2

107 of 1640 proteins reach $p < 0.05$ on this term; chance predicts 82. limma on 30 biopsies, subject blocked; q from BH within trait and term. Table: c_data/04_associate.xlsx, model.

log2 abundance against leg_press_1rm, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2



log2 abundance against leg_press_1rm, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2



log2 abundance against leg_press_1rm, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2



log2 abundance against leg_press_1rm, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2



log2 abundance against leg_press_1rm, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2



log2 abundance against leg_press_1rm, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2



log2 abundance against leg_press_1rm, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2



log2 abundance against leg_press_1rm, within-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2

82 of 1640 proteins reach $p < 0.05$ on this term; chance predicts 82. limma on 30 biopsies, subject blocked; q from BH within trait and term. Table: c_data/04_associate.xlsx, model.

log2 abundance against leg_press_1rm, within-person term

One line per subject from T1 to T2: within is the slope of the lines, between is the spread of subjects


HR LR T1 T2



log2 abundance against leg_press_1rm, within-person term


One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects

HR LR T1 T2




log2 abundance against leg_press_1rm, within-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects



log2 abundance against leg_press_1rm, within-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects




82 of 1640 proteins reach p < 0.05 on this term; chance predicts 82. limma on 30 biopsies, subject blocked; q from BH within trait and term. Table: c_data/04_associate.xlsx, model.

log2 abundance against leg_press_1rm, within-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2

82 of 1640 proteins reach $p < 0.05$ on this term; chance predicts 82. limma on 30 biopsies, subject blocked; q from BH within trait and term. Table: c_data/04_associate.xlsx, model.

log2 abundance against leg_press_1rm, within-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects

HR LR T1 T2




82 of 1640 proteins reach $p < 0.05$ on this term; chance predicts 82. limma on 30 biopsies, subject blocked; q from BH within trait and term. Table: c_data/04_associate.xlsx, model.

log2 abundance against mcsa, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2



log2 abundance against mcsa, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2



log2 abundance against mcsa, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2



log2 abundance against mcsa, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2



log2 abundance against mcsa, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2



log2 abundance against mcsa, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2



log2 abundance against mcsa, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2



log2 abundance against mcsa, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2



log2 abundance against mcsa, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2



log2 abundance against mcsa, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects

—●— HR —●— LR ○ T1 ● T2




206 of 1640 proteins reach $p < 0.05$ on this term; chance predicts 82. limma on 30 biopsies, subject blocked; q from BH within trait and term. Table: c_data/04_associate.xlsx, model.

log2 abundance against mcsa, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2



log2 abundance against mcsa, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


— HR — LR ○ T1 ● T2

206 of 1640 proteins reach $p < 0.05$ on this term; chance predicts 82. limma on 30 biopsies, subject blocked; q from BH within trait and term. Table: c_data/04_associate.xlsx, model.

log2 abundance against mcsa, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects

— HR — LR ○ T1 ● T2




206 of 1640 proteins reach $p < 0.05$ on this term; chance predicts 82. limma on 30 biopsies, subject blocked; q from BH within trait and term. Table: c_data/04_associate.xlsx, model.

log2 abundance against mcsa, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


—●— HR —●— LR ○ T1 ● T2

206 of 1640 proteins reach $p < 0.05$ on this term; chance predicts 82. limma on 30 biopsies, subject blocked; q from BH within trait and term. Table: c_data/04_associate.xlsx, model.

log2 abundance against mcsa, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2



log2 abundance against mcsa, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2

206 of 1640 proteins reach $p < 0.05$ on this term; chance predicts 82. limma on 30 biopsies, subject blocked; q from BH within trait and term. Table: c_data/04_associate.xlsx, model.

log2 abundance against mcsa, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


— HR — LR ○ T1 ● T2

206 of 1640 proteins reach $p < 0.05$ on this term; chance predicts 82. limma on 30 biopsies, subject blocked; q from BH within trait and term. Table: c_data/04_associate.xlsx, model.

log2 abundance against mcsa, between-person term


One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects

HR LR T1 T2



log2 abundance against mcsa, within-person term


One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects



log2 abundance against mcsa, within-person term


One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects

HR LR T1 T2



log2 abundance against mcsa, within-person term


One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects



log2 abundance against mcsa, within-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2




log2 abundance against mcsa, within-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects

HR LR T1 T2






log2 abundance against type1_share, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2



log2 abundance against type1_share, between-person term


One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects

HR LR ○ T1 ● T2

51 of 1640 proteins reach $p < 0.05$ on this term; chance predicts 82. limma on 30 biopsies, subject blocked; q from BH within trait and term. Table: c_data/04_associate.xlsx, model.


log2 abundance against type1_share, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects

51 of 1640 proteins reach $p < 0.05$ on this term; chance predicts 82. limma on 30 biopsies, subject blocked; q from BH within trait and term. Table: c_data/04_associate.xlsx, model.

log2 abundance against type1_share, between-person term


One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects

51 of 1640 proteins reach $p < 0.05$ on this term; chance predicts 82. limma on 30 biopsies, subject blocked; q from BH within trait and term. Table: c_data/04_associate.xlsx, model.

log2 abundance against type1_share, between-person term


One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects

HR LR T1 T2



log2 abundance against type1_share, within-person term
One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2



log2 abundance against type1_share, within-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2



log2 abundance against type1_share, within-person term


One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects

HR LR T1 T2



log2 abundance against type1_share, within-person term


One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects

56 of 1640 proteins reach $p < 0.05$ on this term; chance predicts 82. limma on 30 biopsies, subject blocked; q from BH within trait and term. Table: c_data/04_associate.xlsx, model.


log2 abundance against type1_share, within-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects

HR LR T1 T2




log2 abundance against phenotype, acute window




1683 of 29390 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window



1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: [c_data/04_associate.xlsx](#), [change_score](#) and [change_score_by_arm](#).


log2 abundance against phenotype, acute window



1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.


log2 abundance against phenotype, acute window

HR LR




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.

log2 abundance against phenotype, acute window




log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: [c_data/04_associate.xlsx](#), [change_score](#) and [change_score_by_arm](#).

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




1683 of 29390 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.

log2 abundance against phenotype, acute window




log2 abundance against phenotype, acute window



1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: [c_data/04_associate.xlsx](#), [change_score](#) and [change_score_by_arm](#).


log2 abundance against phenotype, acute window

HR LR




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.

log2 abundance against phenotype, acute window




log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




1683 of 29390 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




1683 of 29390 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.


log2 abundance against phenotype, acute window



log2 abundance against phenotype, acute window




log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: [c_data/04_associate.xlsx](#), [change_score](#) and [change_score_by_arm](#).

log2 abundance against phenotype, acute window




log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.

log2 abundance against phenotype, acute window




log2 abundance against phenotype, acute window




1683 of 29390 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.


log2 abundance against phenotype, acute window



log2 abundance against phenotype, acute window




log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: [c_data/04_associate.xlsx](#), [change_score](#) and [change_score_by_arm](#).


log2 abundance against phenotype, acute window



log2 abundance against phenotype, acute window




log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: [c_data/04_associate.xlsx](#), [change_score](#) and [change_score_by_arm](#).

log2 abundance against phenotype, acute window




log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.

log2 abundance against phenotype, acute window




log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




1683 of 29390 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




1683 of 29390 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




1683 of 29390 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.


log2 abundance against phenotype, acute window



log2 abundance against phenotype, acute window




log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




log2 abundance against phenotype, acute window




1683 of 29390 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.


log2 abundance against phenotype, acute window



log2 abundance against phenotype, acute window




log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p . Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.

log2 abundance against phenotype, acute window



1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.


log2 abundance against phenotype, acute window



1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.


log2 abundance against phenotype, acute window

HR LR




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: [c_data/04_associate.xlsx](#), [change_score](#) and [change_score_by_arm](#).

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: [c_data/04_associate.xlsx](#), [change_score](#) and [change_score_by_arm](#).

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.

log2 abundance against phenotype, acute window




log2 abundance against phenotype, acute window

HR LR




1683 of 29390 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.

log2 abundance against phenotype, acute window



1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.


log2 abundance against phenotype, acute window



1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.


log2 abundance against phenotype, acute window

HR LR



1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.


log2 abundance against phenotype, acute window



1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: [c_data/04_associate.xlsx](#), [change_score](#) and [change_score_by_arm](#).


log2 abundance against phenotype, acute window

HR LR




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: [c_data/04_associate.xlsx](#), [change_score](#) and [change_score_by_arm](#).

log2 abundance against phenotype, acute window



1683 of 29390 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.


log2 abundance against phenotype, acute window



1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.


log2 abundance against phenotype, acute window

HR LR




1683 of 29390 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




1683 of 29390 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window



1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.


log2 abundance against phenotype, acute window



1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.


log2 abundance against phenotype, acute window

HR LR




1683 of 29390 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: [c_data/04_associate.xlsx](#), [change_score](#) and [change_score_by_arm](#).

log2 abundance against phenotype, acute window




1683 of 29390 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.

log2 abundance against phenotype, baseline window




1548 of 31402 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.

log2 abundance against phenotype, baseline window




1548 of 31402 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: [c_data/04_associate.xlsx](#), [change_score](#) and [change_score_by_arm](#).

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.

log2 abundance against phenotype, baseline window




1548 of 31402 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.

log2 abundance against phenotype, baseline window




1548 of 31402 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.

log2 abundance against phenotype, baseline window




log2 abundance against phenotype, baseline window




1548 of 31402 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.


log2 abundance against phenotype, baseline window



log2 abundance against phenotype, baseline window




log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.

log2 abundance against phenotype, baseline window




1548 of 31402 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: [c_data/04_associate.xlsx](#), [change_score](#) and [change_score_by_arm](#).

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window



1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.


log2 abundance against phenotype, baseline window



1548 of 31402 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.


log2 abundance against phenotype, baseline window

HR LR




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: [c_data/04_associate.xlsx](#), [change_score](#) and [change_score_by_arm](#).

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window



1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.


log2 abundance against phenotype, baseline window



1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.


log2 abundance against phenotype, baseline window

HR LR




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.

log2 abundance against phenotype, baseline window




1548 of 31402 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




log2 abundance against phenotype, baseline window




1548 of 31402 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window



1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.


log2 abundance against phenotype, baseline window



1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.


log2 abundance against phenotype, baseline window

HR LR




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.

log2 abundance against phenotype, baseline window




log2 abundance against phenotype, baseline window



log2 abundance against phenotype, baseline window


HR LR



log2 abundance, baseline window, one point per subject


1548 of 31402 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window



log2 abundance against phenotype, baseline window


HR LR



log2 abundance, baseline window, one point per subject


1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.

log2 abundance against phenotype, baseline window




log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: [c_data/04_associate.xlsx](#), [change_score](#) and [change_score_by_arm](#).

log2 abundance against phenotype, baseline window



1548 of 31402 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.


log2 abundance against phenotype, baseline window



1548 of 31402 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window


HR LR



1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window


HR LR



1548 of 31402 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.


log2 abundance against phenotype, baseline window

HR LR



1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.


log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.

log2 abundance against phenotype, baseline window

HR LR




log2 abundance against phenotype, baseline window



1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.


log2 abundance against phenotype, baseline window

HR LR




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.

log2 abundance against phenotype, baseline window



1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.


log2 abundance against phenotype, baseline window



1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.


log2 abundance against phenotype, baseline window

HR LR




1548 of 31402 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




log2 abundance against phenotype, baseline window



1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, training window


HR LR



1145 of 28204 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, training window


HR LR



1145 of 28204 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.


log2 abundance against phenotype, training window

HR LR




1145 of 28204 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, training window




1145 of 28204 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, training window




log2 abundance against phenotype, training window




1145 of 28204 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, training window




1145 of 28204 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, training window




log2 abundance against phenotype, training window

HR LR




log2 abundance against phenotype, training window




1145 of 28204 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, training window




log2 abundance against phenotype, training window




1145 of 28204 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, training window




log2 abundance against phenotype, training window




1145 of 28204 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.

log2 abundance against phenotype, training window




log2 abundance against phenotype, training window




1145 of 28204 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, training window




1145 of 28204 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.

log2 abundance against phenotype, training window




1145 of 28204 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, training window




1145 of 28204 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, training window




1145 of 28204 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, training window




1145 of 28204 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, training window




1145 of 28204 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, training window




1145 of 28204 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, training window




1145 of 28204 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, training window




1145 of 28204 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.

log2 abundance against phenotype, training window




1145 of 28204 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, training window




1145 of 28204 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, training window




1145 of 28204 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, training window




1145 of 28204 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, training window




1145 of 28204 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, training window




1145 of 28204 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, training window




log2 abundance against phenotype, training window




1145 of 28204 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, training window




log2 abundance against phenotype, training window




1145 of 28204 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, training window




1145 of 28204 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.

log2 abundance against phenotype, training window




1145 of 28204 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, training window




1145 of 28204 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.

log2 abundance against phenotype, training window




1145 of 28204 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, training window




1145 of 28204 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, training window




1145 of 28204 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, training window




1145 of 28204 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, training window



1145 of 28204 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: [c_data/04_associate.xlsx](#), [change_score](#) and [change_score_by_arm](#).


log2 abundance against phenotype, training window



1145 of 28204 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.


log2 abundance against phenotype, training window

HR LR




1145 of 28204 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.

log2 abundance against phenotype, training window




log2 abundance against phenotype, training window




1145 of 28204 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, training window




1145 of 28204 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, training window




1145 of 28204 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.

log2 abundance against phenotype, training window




1145 of 28204 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, training window




1145 of 28204 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: [c_data/04_associate.xlsx](#), [change_score](#) and [change_score_by_arm](#).

log2 abundance against phenotype, training window



1145 of 28204 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.


log2 abundance against phenotype, training window



1145 of 28204 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.

log2 abundance against phenotype, training window


HR LR



1145 of 28204 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.

log2 abundance against phenotype, training window


HR LR




1145 of 28204 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, training window

HR LR




log2 abundance against phenotype, training window




1145 of 28204 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: [c_data/04_associate.xlsx](#), [change_score](#) and [change_score_by_arm](#).

log2 abundance against phenotype, training window




1145 of 28204 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, training window




log2 abundance against phenotype, training window




1145 of 28204 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.

log2 abundance against phenotype, training window




1145 of 28204 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: [c_data/04_associate.xlsx](#), [change_score](#) and [change_score_by_arm](#).

log2 abundance against phenotype, training window




1145 of 28204 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.

log2 abundance against phenotype, training window




1145 of 28204 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, training window




1145 of 28204 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, training window




1145 of 28204 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, training window




log2 abundance against phenotype, training window




1145 of 28204 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.

log2 abundance against phenotype, training window




1145 of 28204 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.

log2 abundance against phenotype, training window




1145 of 28204 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: [c_data/04_associate.xlsx](#), [change_score](#) and [change_score_by_arm](#).

log2 abundance against phenotype, training window




1145 of 28204 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.

log2 abundance against phenotype, training window




1145 of 28204 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, training window




1145 of 28204 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, training window




1145 of 28204 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, training window




1145 of 28204 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: [c_data/04_associate.xlsx](#), [change_score](#) and [change_score_by_arm](#).

log2 abundance against phenotype, training window




log2 abundance against phenotype, training window




1145 of 28204 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, training window




1145 of 28204 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, training window




1145 of 28204 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.




1145 of 28204 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, training window




1145 of 28204 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, training window




log2 abundance against phenotype, training window




1145 of 28204 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.

log2 abundance against phenotype, training window




1145 of 28204 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, training window




1145 of 28204 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: [c_data/04_associate.xlsx](#), [change_score](#) and [change_score_by_arm](#).

log2 abundance against phenotype, training window




1145 of 28204 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, training window




1145 of 28204 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, training window




1145 of 28204 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, training window




1145 of 28204 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, training window




1145 of 28204 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.


log2 abundance against phenotype, training window



1145 of 28204 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.




log2 abundance against phenotype, training window




1145 of 28204 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, training window



1145 of 28204 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.


log2 abundance against phenotype, training window



1145 of 28204 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, training window

HR LR



1145 of 28204 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.