



Nominal classifiers against chance




Chance is 5% of proteins tested. With 6 to 8 pairs the Wilcoxon p is coarse, so fewer than 5% of null tests reach 0.05 and a ratio below 1 is not worse than chance.

Training, HR (T1 to T2)


log2 abundance; AUC above 0.5 favours T2; 6 vs 6 paired samples; page 1 of 4





Training, HR (T1 to T2)


log2 abundance; AUC above 0.5 favours T2; 6 vs 6 paired samples; page 3 of 4



43 of 1555 proteins reach $p < 0.05$; chance predicts 78. Smallest attainable p: 0.031. p from the Wilcoxon signed-rank test, q from BH within task. Table: c_data/03_classify.xlsx, auc.

Training, HR (T1 to T2)


log2 abundance; AUC above 0.5 favours T2; 6 vs 6 paired samples; page 4 of 4



43 of 1555 proteins reach $p < 0.05$; chance predicts 78. Smallest attainable p : 0.031. p from the Wilcoxon signed-rank test, q from BH within task. Table: c_data/03_classify.xlsx, auc.


Training, LR (T1 to T2)

log2 abundance; AUC above 0.5 favours T2; 8 vs 8 paired samples; page 1 of 7




Training, LR (T1 to T2)

log2 abundance; AUC above 0.5 favours T2; 8 vs 8 paired samples; page 2 of 7




Training, LR (T1 to T2)

log2 abundance; AUC above 0.5 favours T2; 8 vs 8 paired samples; page 3 of 7




Training, LR (T1 to T2)

log2 abundance; AUC above 0.5 favours T2; 8 vs 8 paired samples; page 4 of 7




Training, LR (T1 to T2)

log2 abundance; AUC above 0.5 favours T2; 8 vs 8 paired samples; page 5 of 7




Training, LR (T1 to T2)

log2 abundance; AUC above 0.5 favours T2; 8 vs 8 paired samples; page 6 of 7



Training, LR (T1 to T2)


log2 abundance; AUC above 0.5 favours T2; 8 vs 8 paired samples; page 7 of 7



78 of 1771 proteins reach $p < 0.05$; chance predicts 89. Smallest attainable p: 0.0078. p from the Wilcoxon signed-rank test, q from BH within task. Table: c_data/03_classify.xlsx, auc.

Acute bout, HR (T2 to T3)


log2 abundance; AUC above 0.5 favours T3; 7 vs 7 paired samples; page 1 of 10



116 of 1703 proteins reach $p < 0.05$; chance predicts 85. Smallest attainable p: 0.016. p from the Wilcoxon signed-rank test, q from BH within task. Table: c_data/03_classify.xlsx, auc.


Acute bout, HR (T2 to T3)

log2 abundance; AUC above 0.5 favours T3; 7 vs 7 paired samples; page 2 of 10




Acute bout, HR (T2 to T3)


log2 abundance; AUC above 0.5 favours T3; 7 vs 7 paired samples; page 3 of 10



Acute bout, HR (T2 to T3)


log2 abundance; AUC above 0.5 favours T3; 7 vs 7 paired samples; page 4 of 10





Acute bout, HR (T2 to T3)


log2 abundance; AUC above 0.5 favours T3; 7 vs 7 paired samples; page 6 of 10



116 of 1703 proteins reach $p < 0.05$; chance predicts 85. Smallest attainable p : 0.016. p from the Wilcoxon signed-rank test, q from BH within task. Table: c_data/03_classify.xlsx, auc.


Acute bout, HR (T2 to T3)

log2 abundance; HR above 0.5 favours T3; 7 vs 7 paired samples; page 7 of 10




Acute bout, HR (T2 to T3)

log2 abundance; AUC above 0.5 favours T3; 7 vs 7 paired samples; page 8 of 10




Acute bout, HR (T2 to T3)

log2 abundance; AUC above 0.5 favours T3; 7 vs 7 paired samples; page 9 of 10




Acute bout, HR (T2 to T3)

log2 abundance; AUC above 0.5 favours T3; 7 vs 7 paired samples; page 10 of 10




Acute bout, LR (T2 to T3)

log2 abundance; AUC above 0.5 favours T3; 8 vs 8 paired samples; page 1 of 9




Acute bout, LR (T2 to T3)

log2 abundance; AUC above 0.5 favours T3; 8 vs 8 paired samples; page 2 of 9




Acute bout, LR (T2 to T3)

log2 abundance; AUC above 0.5 favours T3; 8 vs 8 paired samples; page 3 of 9




Acute bout, LR (T2 to T3)

log2 abundance; AUC above 0.5 favours T3; 8 vs 8 paired samples; page 4 of 9




Acute bout, LR (T2 to T3)

log2 abundance; AUC above 0.5 favours T3; 8 vs 8 paired samples; page 5 of 9




Acute bout, LR (T2 to T3)

log2 abundance; AUC above 0.5 favours T3; 8 vs 8 paired samples; page 6 of 9




Acute bout, LR (T2 to T3)

log2 abundance; AUC above 0.5 favours T3; 8 vs 8 paired samples; page 7 of 9




Acute bout, LR (T2 to T3)

log2 abundance; AUC above 0.5 favours T3; 8 vs 8 paired samples; page 8 of 9




Acute bout, LR (T2 to T3)

log2 abundance; AUC above 0.5 favours T3; 8 vs 8 paired samples; page 9 of 9




HR vs LR at T2

log2 abundance; AUC above 0.5 favours HR; 7 vs 8 subjects; page 1 of 6




HR vs LR at T2

log2 abundance; AUC above 0.5 favours HR; 7 vs 8 subjects; page 2 of 6




HR vs LR at T2

log2 abundance; AUC above 0.5 favours HR; 7 vs 8 subjects; page 3 of 6




HR vs LR at T2


log2 abundance; AUC above 0.5 favours HR; 7 vs 8 subjects; page 4 of 6



HR vs LR at T2


log2 abundance; AUC above 0.5 favours HR; 7 vs 8 subjects; page 5 of 6






HR vs LR, training change

log2 abundance; AUC above 0.5 favours HR; 6 vs 8 subjects; page 1 of 4




HR vs LR, training change

log2 abundance; AUC above 0.5 favours HR; 6 vs 8 subjects; page 2 of 4




HR vs LR, training change

log2 abundance; AUC above 0.5 favours HR; 6 vs 8 subjects; page 3 of 4



HR vs LR, training change


log2 abundance; AUC above 0.5 favours HR; 6 vs 8 subjects; page 4 of 4



39 of 1539 proteins reach $p < 0.05$; chance predicts 77. Smallest attainable p: 0.00067. p from the Wilcoxon rank-sum test, q from BH within task. Table: c_data/03_classify.xlsx, auc.


HR vs LR, acute change


log2 abundance; AUC above 0.5 favours HR; 7 vs 8 subjects; page 1 of 8



HR vs LR, acute change


log2 abundance; AUC above 0.5 favours HR; 7 vs 8 subjects; page 2 of 8






HR vs LR, acute change


log2 abundance; AUC above 0.5 favours HR; 7 vs 8 subjects; page 4 of 8






HR vs LR, acute change

log2 abundance; AUC above 0.5 favours HR; 7 vs 8 subjects; page 6 of 8




HR vs LR, acute change

log2 abundance; AUC above 0.5 favours HR; 7 vs 8 subjects; page 7 of 8



HR vs LR, acute change

log2 abundance; AUC above 0.5 favours HR; 7 vs 8 subjects; page 8 of 8



88 of 1645 proteins reach $p < 0.05$; chance predicts 82. Smallest attainable p : 0.00031. p from the Wilcoxon rank-sum test, q from BH within task. Table: c_data/03_classify.xlsx, auc.