


Nominal associations against chance




Chance is 5% of proteins tested per cell. Table: c_data/04_associate.xlsx, chance_expectation.

log2 abundance against fc_{sa}_mixed, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2



log2 abundance against fc_{sa}_mixed, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2



log2 abundance against fcsc_mixed, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects

HR LR T1 T2




67 of 1640 proteins reach $p < 0.05$ on this term; chance predicts 82. limma on 30 biopsies, subject blocked; q from BH within trait and term. Table: c_data/04_associate.xlsx, model.

log2 abundance against fcsa_mixed, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2



log2 abundance against fcsa_mixed, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2



log2 abundance against fc_{sa}_mixed, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2



log2 abundance against fcsc_mixed, within-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2



log2 abundance against fcsa_mixed, within-person term


One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects

HR LR T1 T2



log2 abundance against fc_{sa_mixed}, within-person term


One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects

61 of 1640 proteins reach $p < 0.05$ on this term; chance predicts 82. limma on 30 biopsies, subject blocked; q from BH within trait and term. Table: c_data/04_associate.xlsx, model.

log2 abundance against fc_{sa}_mixed, within-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects

HR LR T1 T2




61 of 1640 proteins reach p < 0.05 on this term; chance predicts 82. limma on 30 biopsies, subject blocked; q from BH within trait and term. Table: c_data/04_associate.xlsx, model.

log2 abundance against fcsa_mixed, within-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects

HR LR T1 T2




log2 abundance against fc_{sa}_mixed, within-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects

HR LR T1 T2


PARVA
within -0.000353 p 0.05 q 0.99



log2 abundance against fcsc_type1, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2



log2 abundance against fcsa_type1, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2



log2 abundance against fc_{sa}_type1, between-person term


One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects

HR LR T1 T2



log2 abundance against fc_{sa}_type1, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects




64 of 1640 proteins reach p < 0.05 on this term; chance predicts 82. limma on 30 biopsies, subject blocked; q from BH within trait and term. Table: c_data/04_associate.xlsx, model.

log2 abundance against fc_{sa}_type1, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2



log2 abundance against fc_{sa_type1}, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2



log2 abundance against fc_{sa}_type1, within-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2



log2 abundance against fcsc_type1, within-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2



log2 abundance against fc_{sa_type1}, within-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR ○ T1 ● T2



log2 abundance against fcsc_type1, within-person term


One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects

HR LR T1 T2




log2 abundance against fc_{sa_type1}, within-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects




log2 abundance against fcsa_type2, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects

64 of 1640 proteins reach $p < 0.05$ on this term; chance predicts 82. limma on 30 biopsies, subject blocked; q from BH within trait and term. Table: c_data/04_associate.xlsx, model.

log2 abundance against fcsa_type2, between-person term


One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects



64 of 1640 proteins reach $p < 0.05$ on this term; chance predicts 82. limma on 30 biopsies, subject blocked; q from BH within trait and term. Table: c_data/04_associate.xlsx, model.


log2 abundance against fcsc_type2, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects

64 of 1640 proteins reach $p < 0.05$ on this term; chance predicts 82. limma on 30 biopsies, subject blocked; q from BH within trait and term. Table: c_data/04_associate.xlsx, model.

log2 abundance against fc_{sa}_type2, between-person term


One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects

64 of 1640 proteins reach $p < 0.05$ on this term; chance predicts 82. limma on 30 biopsies, subject blocked; q from BH within trait and term. Table: c_data/04_associate.xlsx, model.

log2 abundance against fc_{sa_type2}, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2



log2 abundance against fcsa_type2, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2



log2 abundance against fc_{sa}_type2, within-person term


One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects

HR LR T1 T2



log2 abundance against fcsc_type2, within-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects




74 of 1640 proteins reach p < 0.05 on this term; chance predicts 82. limma on 30 biopsies, subject blocked; q from BH within trait and term. Table: c_data/04_associate.xlsx, model.

log2 abundance against fcsc_type2, within-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2



74 of 1640 proteins reach p < 0.05 on this term; chance predicts 82. limma on 30 biopsies, subject blocked; q from BH within trait and term. Table: c_data/04_associate.xlsx, model.

log2 abundance against fc_{sa}_type2, within-person term


One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects



74 of 1640 proteins reach p < 0.05 on this term; chance predicts 82. limma on 30 biopsies, subject blocked; q from BH within trait and term. Table: c_data/04_associate.xlsx, model.


log2 abundance against fcsa_type2, within-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects

74 of 1640 proteins reach $p < 0.05$ on this term; chance predicts 82. limma on 30 biopsies, subject blocked; q from BH within trait and term. Table: c_data/04_associate.xlsx, model.

log2 abundance against fc_{sa_type2}, within-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects



74 of 1640 proteins reach p < 0.05 on this term; chance predicts 82. limma on 30 biopsies, subject blocked; q from BH within trait and term. Table: c_data/04_associate.xlsx, model.



log2 abundance against fcsa_type2, within-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects

HR LR T1 T2

PLGRKT
within 0.000281 p 0.049 q 0.99


ACOT1
within -0.00054 p 0.05 q 0.99



log2 abundance against fibres_mixed, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2



log2 abundance against fibres_mixed, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects

HR LR T1 T2




83 of 1640 proteins reach p < 0.05 on this term; chance predicts 82. limma on 30 biopsies, subject blocked; q from BH within trait and term. Table: c_data/04_associate.xlsx, model.

log2 abundance against fibres_mixed, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects

HR LR T1 T2




83 of 1640 proteins reach p < 0.05 on this term; chance predicts 82. limma on 30 biopsies, subject blocked; q from BH within trait and term. Table: c_data/04_associate.xlsx, model.

log2 abundance against fibres_mixed, between-person term


One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects

HR LR T1 T2




log2 abundance against fibres_mixed, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects

83 of 1640 proteins reach $p < 0.05$ on this term; chance predicts 82. limma on 30 biopsies, subject blocked; q from BH within trait and term. Table: c_data/04_associate.xlsx, model.

log2 abundance against fibres_mixed, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects




83 of 1640 proteins reach p < 0.05 on this term; chance predicts 82. limma on 30 biopsies, subject blocked; q from BH within trait and term. Table: c_data/04_associate.xlsx, model.

log2 abundance against fibres_mixed, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2

83 of 1640 proteins reach $p < 0.05$ on this term; chance predicts 82. limma on 30 biopsies, subject blocked; q from BH within trait and term. Table: c_data/04_associate.xlsx, model.

log2 abundance against fibres_mixed, within-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2



log2 abundance against fibres_mixed, within-person term


One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects

HR LR T1 T2



log2 abundance against fibres_mixed, within-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects




46 of 1640 proteins reach p < 0.05 on this term; chance predicts 82. limma on 30 biopsies, subject blocked; q from BH within trait and term. Table: c_data/04_associate.xlsx, model.

log2 abundance against fibres_mixed, within-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2



log2 abundance against fibres_type1, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2



log2 abundance against fibres_type1, between-person term


One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects

HR LR T1 T2



log2 abundance against fibres_type1, between-person term


One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects

63 of 1640 proteins reach $p < 0.05$ on this term; chance predicts 82. limma on 30 biopsies, subject blocked; q from BH within trait and term. Table: c_data/04_associate.xlsx, model.

log2 abundance against fibres_type1, between-person term


One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects

HR LR T1 T2



log2 abundance against fibres_type1, between-person term


One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects

63 of 1640 proteins reach $p < 0.05$ on this term; chance predicts 82. limma on 30 biopsies, subject blocked; q from BH within trait and term. Table: c_data/04_associate.xlsx, model.

log2 abundance against fibres_type1, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2



log2 abundance against fibres_type1, within-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2

52 of 1640 proteins reach $p < 0.05$ on this term; chance predicts 82. limma on 30 biopsies, subject blocked; q from BH within trait and term. Table: c_data/04_associate.xlsx, model.

log2 abundance against fibres_type1, within-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2



log2 abundance against fibres_type1, within-person term


One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects

HR LR T1 T2



log2 abundance against fibres_type1, within-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects




52 of 1640 proteins reach $p < 0.05$ on this term; chance predicts 82. limma on 30 biopsies, subject blocked; q from BH within trait and term. Table: c_data/04_associate.xlsx, model.

log2 abundance against fibres_type1, within-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2



log2 abundance against leg_ext_1rm, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects

HR LR T1 T2




130 of 1615 proteins reach $p < 0.05$ on this term; chance predicts 81. limma on 29 biopsies, subject blocked; q from BH within trait and term. Table: c_data/04_associate.xlsx, model.

log2 abundance against leg_ext_1rm, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects

HR LR T1 T2




130 of 1615 proteins reach $p < 0.05$ on this term; chance predicts 81. limma on 29 biopsies, subject blocked; q from BH within trait and term. Table: c_data/04_associate.xlsx, model.

log2 abundance against leg_ext_1rm, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects

HR LR T1 T2




130 of 1615 proteins reach $p < 0.05$ on this term; chance predicts 81. limma on 29 biopsies, subject blocked; q from BH within trait and term. Table: c_data/04_associate.xlsx, model.

log2 abundance against leg_ext_1rm, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects

HR LR T1 T2




130 of 1615 proteins reach $p < 0.05$ on this term; chance predicts 81. limma on 29 biopsies, subject blocked; q from BH within trait and term. Table: c_data/04_associate.xlsx, model.

log2 abundance against leg_ext_1rm, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects

HR LR T1 T2




130 of 1615 proteins reach $p < 0.05$ on this term; chance predicts 81. limma on 29 biopsies, subject blocked; q from BH within trait and term. Table: c_data/04_associate.xlsx, model.

log2 abundance against leg_ext_1rm, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2



130 of 1615 proteins reach $p < 0.05$ on this term; chance predicts 81. limma on 29 biopsies, subject blocked; q from BH within trait and term. Table: c_data/04_associate.xlsx, model.

log2 abundance against leg_ext_1rm, between-person term


One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects



log2 abundance against leg_ext_1rm, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2



log2 abundance against leg_ext_1rm, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2



log2 abundance against leg_ext_1rm, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2



log2 abundance against leg_ext_1rm, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2



log2 abundance against leg_ext_1rm, within-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2



log2 abundance against leg_ext_1rm, within-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects

HR LR T1 T2




122 of 1615 proteins reach $p < 0.05$ on this term; chance predicts 81. limma on 29 biopsies, subject blocked; q from BH within trait and term. Table: c_data/04_associate.xlsx, model.

log2 abundance against leg_ext_1rm, within-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2



log2 abundance against leg_ext_1rm, within-person term

One line per subject from T1 to T2: within is the slope of the lines, between is the spread of subjects


HR LR T1 T2



log2 abundance against leg_ext_1rm, within-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects

HR LR T1 T2




122 of 1615 proteins reach $p < 0.05$ on this term; chance predicts 81. limma on 29 biopsies, subject blocked; q from BH within trait and term. Table: c_data/04_associate.xlsx, model.

log2 abundance against leg_ext_1rm, within-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2



log2 abundance against leg_ext_1rm, within-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2



log2 abundance against leg_ext_1rm, within-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects

HR LR T1 T2




122 of 1615 proteins reach $p < 0.05$ on this term; chance predicts 81. limma on 29 biopsies, subject blocked; q from BH within trait and term. Table: c_data/04_associate.xlsx, model.

log2 abundance against leg_ext_1rm, within-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2



log2 abundance against leg_ext_1rm, within-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2



log2 abundance against leg_ext_1rm, within-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2



log2 abundance against leg_press_1rm, between-person term


One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects

HR LR T1 T2



log2 abundance against leg_press_1rm, between-person term


One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


107 of 1640 proteins reach $p < 0.05$ on this term; chance predicts 82. limma on 30 biopsies, subject blocked; q from BH within trait and term. Table: c_data/04_associate.xlsx, model.

log2 abundance against leg_press_1rm, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2



log2 abundance against leg_press_1rm, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2



log2 abundance against leg_press_1rm, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2



log2 abundance against leg_press_1rm, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2



log2 abundance against leg_press_1rm, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2



log2 abundance against leg_press_1rm, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2



log2 abundance against leg_press_1rm, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2



log2 abundance against leg_press_1rm, within-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2



log2 abundance against leg_press_1rm, within-person term

One line per subject from T1 to T2: within is the slope of the lines, between is the spread of subjects


HR LR T1 T2



log2 abundance against leg_press_1rm, within-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2



log2 abundance against leg_press_1rm, within-person term


One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects

HR LR T1 T2



log2 abundance against leg_press_1rm, within-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects




82 of 1640 proteins reach p < 0.05 on this term; chance predicts 82. limma on 30 biopsies, subject blocked; q from BH within trait and term. Table: c_data/04_associate.xlsx, model.

log2 abundance against leg_press_1rm, within-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2



log2 abundance against leg_press_1rm, within-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects

HR LR T1 T2




leg_press_1rm at the biopsy

82 of 1640 proteins reach $p < 0.05$ on this term; chance predicts 82. limma on 30 biopsies, subject blocked; q from BH within trait and term. Table: c_data/04_associate.xlsx, model.

log2 abundance against mcsa, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2



log2 abundance against mcsa, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2



log2 abundance against mcsa, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2



log2 abundance against mcsa, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2



log2 abundance against mcsa, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2



log2 abundance against mcsa, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2



log2 abundance against mcsa, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2



log2 abundance against mcsa, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2



log2 abundance against mcsa, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2



log2 abundance against mcsa, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2



log2 abundance against mcsa, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2



log2 abundance against mcsa, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


— HR — LR ○ T1 ● T2

206 of 1640 proteins reach $p < 0.05$ on this term; chance predicts 82. limma on 30 biopsies, subject blocked; q from BH within trait and term. Table: c_data/04_associate.xlsx, model.

log2 abundance against mcsa, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


— HR — LR ○ T1 ● T2

206 of 1640 proteins reach $p < 0.05$ on this term; chance predicts 82. limma on 30 biopsies, subject blocked; q from BH within trait and term. Table: c_data/04_associate.xlsx, model.

log2 abundance against mcsa, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2



log2 abundance against mcsa, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


— HR — LR ○ T1 ● T2

206 of 1640 proteins reach $p < 0.05$ on this term; chance predicts 82. limma on 30 biopsies, subject blocked; q from BH within trait and term. Table: c_data/04_associate.xlsx, model.

log2 abundance against mcsa, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


—●— HR —●— LR ○ T1 ● T2

206 of 1640 proteins reach $p < 0.05$ on this term; chance predicts 82. limma on 30 biopsies, subject blocked; q from BH within trait and term. Table: c_data/04_associate.xlsx, model.

log2 abundance against mcsa, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


— HR — LR ○ T1 ● T2

206 of 1640 proteins reach $p < 0.05$ on this term; chance predicts 82. limma on 30 biopsies, subject blocked; q from BH within trait and term. Table: c_data/04_associate.xlsx, model.

log2 abundance against mcsa, between-person term


One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects

HR LR T1 T2



log2 abundance against mcsa, within-person term


One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects



log2 abundance against mcsa, within-person term


One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects

HR LR T1 T2



log2 abundance against mcsa, within-person term


One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects



log2 abundance against mcsa, within-person term


One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects

HR LR T1 T2



log2 abundance against mcsa, within-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects




63 of 1640 proteins reach $p < 0.05$ on this term; chance predicts 82. limma on 30 biopsies, subject blocked; q from BH within trait and term. Table: c_data/04_associate.xlsx, model.

log2 abundance against mcsa, within-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR O T1 ● T2



log2 abundance against type1_share, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2



log2 abundance against type1_share, between-person term


One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects

HR LR ○ T1 ● T2

51 of 1640 proteins reach $p < 0.05$ on this term; chance predicts 82. limma on 30 biopsies, subject blocked; q from BH within trait and term. Table: c_data/04_associate.xlsx, model.


log2 abundance against type1_share, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects

51 of 1640 proteins reach $p < 0.05$ on this term; chance predicts 82. limma on 30 biopsies, subject blocked; q from BH within trait and term. Table: c_data/04_associate.xlsx, model.

log2 abundance against type1_share, between-person term


One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects

51 of 1640 proteins reach $p < 0.05$ on this term; chance predicts 82. limma on 30 biopsies, subject blocked; q from BH within trait and term. Table: c_data/04_associate.xlsx, model.

log2 abundance against type1_share, between-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2



log2 abundance against type1_share, within-person term


One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects

HR LR T1 T2



log2 abundance against type1_share, within-person term


One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects

56 of 1640 proteins reach $p < 0.05$ on this term; chance predicts 82. limma on 30 biopsies, subject blocked; q from BH within trait and term. Table: c_data/04_associate.xlsx, model.

log2 abundance against type1_share, within-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects


HR LR T1 T2



log2 abundance against type1_share, within-person term

One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects

HR LR T1 T2




56 of 1640 proteins reach $p < 0.05$ on this term; chance predicts 82. limma on 30 biopsies, subject blocked; q from BH within trait and term. Table: c_data/04_associate.xlsx, model.

log2 abundance against type1_share, within-person term


One line per subject from T1 to T2: within is the slope of the lines, between the spread of subjects

HR LR T1 T2




log2 abundance against phenotype, acute window

HR LR




1683 of 29390 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




1683 of 29390 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.

log2 abundance against phenotype, acute window




log2 abundance against phenotype, acute window




1683 of 29390 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window



1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.


log2 abundance against phenotype, acute window



1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.


log2 abundance against phenotype, acute window

HR LR




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




1683 of 29390 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window



1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.


log2 abundance against phenotype, acute window



1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.


log2 abundance against phenotype, acute window

HR LR




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.

log2 abundance against phenotype, acute window




log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




1683 of 29390 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




log2 abundance against phenotype, acute window




1683 of 29390 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.


log2 abundance against phenotype, acute window



log2 abundance against phenotype, acute window




log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.


log2 abundance against phenotype, acute window



log2 abundance against phenotype, acute window




log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.

log2 abundance against phenotype, acute window




log2 abundance against phenotype, acute window




1683 of 29390 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: [c_data/04_associate.xlsx](#), [change_score](#) and [change_score_by_arm](#).

log2 abundance against phenotype, acute window




log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.


log2 abundance against phenotype, acute window




log2 abundance against phenotype, acute window



1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.




log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.

log2 abundance against phenotype, acute window




log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




1683 of 29390 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




1683 of 29390 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: [c_data/04_associate.xlsx](#), [change_score](#) and [change_score_by_arm](#).

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




1683 of 29390 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




1683 of 29390 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.

log2 abundance against phenotype, acute window




1683 of 29390 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




1683 of 29390 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.


log2 abundance against phenotype, acute window



log2 abundance against phenotype, acute window




log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.

log2 abundance against phenotype, acute window



1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.


log2 abundance against phenotype, acute window



1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.


log2 abundance against phenotype, acute window

HR LR




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window



1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: [c_data/04_associate.xlsx](#), [change_score](#) and [change_score_by_arm](#).


log2 abundance against phenotype, acute window



1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: [c_data/04_associate.xlsx](#), [change_score](#) and [change_score_by_arm](#).


log2 abundance against phenotype, acute window

HR LR




1683 of 29390 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




log2 abundance against phenotype, acute window

HR LR



1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.


log2 abundance against phenotype, acute window



1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.


log2 abundance against phenotype, acute window

HR LR



1683 of 29390 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.


log2 abundance against phenotype, acute window



1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.


log2 abundance against phenotype, acute window

HR LR



1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.


log2 abundance against phenotype, acute window



1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: [c_data/04_associate.xlsx](#), [change_score](#) and [change_score_by_arm](#).


log2 abundance against phenotype, acute window

HR LR




1683 of 29390 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window



1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.


log2 abundance against phenotype, acute window



1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.


log2 abundance against phenotype, acute window

HR LR




1683 of 29390 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window



1683 of 29390 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.


log2 abundance against phenotype, acute window



1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.


log2 abundance against phenotype, acute window

HR LR




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, acute window




1683 of 29390 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: [c_data/04_associate.xlsx](#), [change_score](#) and [change_score_by_arm](#).

log2 abundance against phenotype, acute window




1683 of 29390 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1470. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.

log2 abundance against phenotype, baseline window




1548 of 31402 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.

log2 abundance against phenotype, baseline window




1548 of 31402 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.

log2 abundance against phenotype, baseline window




1548 of 31402 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: [c_data/04_associate.xlsx](#), [change_score](#) and [change_score_by_arm](#).

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.

log2 abundance against phenotype, baseline window




1548 of 31402 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.

log2 abundance against phenotype, baseline window




1548 of 31402 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window



1548 of 31402 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.


log2 abundance against phenotype, baseline window



1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.


log2 abundance against phenotype, baseline window

HR LR




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: [c_data/04_associate.xlsx](#), [change_score](#) and [change_score_by_arm](#).

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p . Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: [c_data/04_associate.xlsx](#), [change_score](#) and [change_score_by_arm](#).

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: [c_data/04_associate.xlsx](#), [change_score](#) and [change_score_by_arm](#).

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.

log2 abundance against phenotype, baseline window




log2 abundance against phenotype, baseline window




1548 of 31402 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.


log2 abundance against phenotype, baseline window



log2 abundance against phenotype, baseline window




log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window



1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.


log2 abundance against phenotype, baseline window



1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.


log2 abundance against phenotype, baseline window

HR LR




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




log2 abundance against phenotype, baseline window



1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.

log2 abundance against phenotype, baseline window


HR LR



log2 abundance, baseline window, one point per subject


1548 of 31402 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




log2 abundance against phenotype, baseline window

HR LR




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.

log2 abundance against phenotype, baseline window




1548 of 31402 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window



1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.


log2 abundance against phenotype, baseline window



1548 of 31402 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window


HR LR



1548 of 31402 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.


log2 abundance against phenotype, baseline window

HR LR




1548 of 31402 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window



1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.


log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.

log2 abundance against phenotype, baseline window

HR LR




log2 abundance against phenotype, baseline window




1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window



1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.


log2 abundance against phenotype, baseline window



1548 of 31402 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window


HR LR



log2 abundance, baseline window, one point per subject


1548 of 31402 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window



1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: [c_data/04_associate.xlsx](#), [change_score](#) and [change_score_by_arm](#).


log2 abundance against phenotype, baseline window



1548 of 31402 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.


log2 abundance against phenotype, baseline window

HR LR




1548 of 31402 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, baseline window




log2 abundance against phenotype, baseline window



1548 of 31402 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1570. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, training window


HR LR



1145 of 28204 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, training window


HR LR



1145 of 28204 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.


log2 abundance against phenotype, training window

HR LR




1145 of 28204 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, training window



1145 of 28204 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, training window




log2 abundance against phenotype, training window




1145 of 28204 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, training window




1145 of 28204 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: [c_data/04_associate.xlsx](#), [change_score](#) and [change_score_by_arm](#).

log2 abundance against phenotype, training window




log2 abundance against phenotype, training window

HR LR




log2 abundance against phenotype, training window




1145 of 28204 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, training window




log2 abundance against phenotype, training window




1145 of 28204 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, training window




log2 abundance against phenotype, training window




1145 of 28204 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.

log2 abundance against phenotype, training window




1145 of 28204 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, training window




log2 abundance against phenotype, training window




1145 of 28204 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.

log2 abundance against phenotype, training window




1145 of 28204 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, training window




log2 abundance against phenotype, training window




1145 of 28204 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, training window




1145 of 28204 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, training window




1145 of 28204 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, training window




1145 of 28204 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, training window




1145 of 28204 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, training window




1145 of 28204 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.

log2 abundance against phenotype, training window




1145 of 28204 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, training window




1145 of 28204 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, training window




log2 abundance against phenotype, training window




1145 of 28204 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, training window




log2 abundance against phenotype, training window




1145 of 28204 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, training window




log2 abundance against phenotype, training window




1145 of 28204 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, training window




log2 abundance against phenotype, training window

HR LR




1145 of 28204 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, training window




1145 of 28204 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.

log2 abundance against phenotype, training window




1145 of 28204 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, training window




1145 of 28204 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, training window




1145 of 28204 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, training window




1145 of 28204 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, training window




1145 of 28204 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, training window




1145 of 28204 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, training window



1145 of 28204 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.


log2 abundance against phenotype, training window



1145 of 28204 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.


log2 abundance against phenotype, training window

HR LR




1145 of 28204 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.

log2 abundance against phenotype, training window




log2 abundance against phenotype, training window




1145 of 28204 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, training window




1145 of 28204 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, training window




1145 of 28204 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, training window




log2 abundance against phenotype, training window




1145 of 28204 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, training window



1145 of 28204 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.


log2 abundance against phenotype, training window



1145 of 28204 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.

log2 abundance against phenotype, training window


HR LR



1145 of 28204 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, training window


HR LR



1145 of 28204 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.


log2 abundance against phenotype, training window

HR LR




1145 of 28204 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, training window




1145 of 28204 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, training window




1145 of 28204 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, training window




log2 abundance against phenotype, training window




1145 of 28204 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, training window




1145 of 28204 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.

log2 abundance against phenotype, training window




1145 of 28204 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: [c_data/04_associate.xlsx](#), [change_score](#) and [change_score_by_arm](#).

log2 abundance against phenotype, training window




1145 of 28204 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, training window




1145 of 28204 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, training window




1145 of 28204 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.


log2 abundance against phenotype, training window



log2 abundance against phenotype, training window




log2 abundance against phenotype, training window




1145 of 28204 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.

log2 abundance against phenotype, training window




1145 of 28204 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.

log2 abundance against phenotype, training window




1145 of 28204 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.

log2 abundance against phenotype, training window




1145 of 28204 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: `c_data/04_associate.xlsx`, `change_score` and `change_score_by_arm`.

log2 abundance against phenotype, training window




1145 of 28204 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, training window




1145 of 28204 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, training window




1145 of 28204 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, training window




1145 of 28204 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, training window




1145 of 28204 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, training window




1145 of 28204 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: [c_data/04_associate.xlsx](#), [change_score](#) and [change_score_by_arm](#).

log2 abundance against phenotype, training window




1145 of 28204 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, training window




1145 of 28204 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, training window




1145 of 28204 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, training window




log2 abundance against phenotype, training window




1145 of 28204 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, training window




1145 of 28204 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, training window




1145 of 28204 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: [c_data/04_associate.xlsx](#), [change_score](#) and [change_score_by_arm](#).

log2 abundance against phenotype, training window




1145 of 28204 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, training window




1145 of 28204 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, training window




1145 of 28204 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, training window




1145 of 28204 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, training window




1145 of 28204 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.




1145 of 28204 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's ρ and exact p . Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, training window




1145 of 28204 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, training window



1145 of 28204 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.


log2 abundance against phenotype, training window



1145 of 28204 protein-outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

log2 abundance against phenotype, training window

HR LR



1145 of 28204 protein–outcome pairs reach pooled $p < 0.05$; chance predicts 1410. Header: pooled Spearman, q from BH within window and outcome. Corner: each arm's rho and exact p. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.