

Discrimination overview: proteins



All 1,900 proteins, one row each. Rows are ordered by average−linkage clustering (1 − Pearson r) of limpa's matrix, for display only. Grey: not tested. Table: c_data/01_proteins.xlsx, auc.

Hi vs Lo at T1


121 of 1900 proteins at p < 0.05, 0 at BH < 0.05. Page 1 of 2.



Hi (up to 8) against Lo (up to 6); AUC above 0.5 means higher in Hi. Header: AUC with its stratified bootstrap 95% interval (2000 resamples), Wilcoxon rank−sum p (smallest attainable 0.00067), BH q within task, and the fea

Hi vs Lo at T1


121 of 1900 proteins at p < 0.05, 0 at BH < 0.05. Page 2 of 2.



Hi (up to 8) against Lo (up to 6); AUC above 0.5 means higher in Hi. Header: AUC with its stratified bootstrap 95% interval (2000 resamples), Wilcoxon rank−sum p (smallest attainable 0.00067), BH q within task, and the fea

Hi vs Lo at T2


76 of 1900 proteins at p < 0.05, 0 at BH < 0.05. Page 1 of 2.



Hi (up to 7) against Lo (up to 7); AUC above 0.5 means higher in Hi. Header: AUC with its stratified bootstrap 95% interval (2000 resamples), Wilcoxon rank−sum p (smallest attainable 0.00058), BH q within task, and the fea

Hi vs Lo at T2


76 of 1900 proteins at p < 0.05, 0 at BH < 0.05. Page 2 of 2.



Hi (up to 7) against Lo (up to 7); AUC above 0.5 means higher in Hi. Header: AUC with its stratified bootstrap 95% interval (2000 resamples), Wilcoxon rank−sum p (smallest attainable 0.00058), BH q within task, and the fea

Hi vs Lo at T3


53 of 1900 proteins at p < 0.05, 0 at BH < 0.05. Page 1 of 2.



Hi (up to 7) against Lo (up to 7); AUC above 0.5 means higher in Hi. Header: AUC with its stratified bootstrap 95% interval (2000 resamples), Wilcoxon rank−sum p (smallest attainable 0.00058), BH q within task, and the fea

Hi vs Lo at T3


53 of 1900 proteins at p < 0.05, 0 at BH < 0.05. Page 2 of 2.



Hi (up to 7) against Lo (up to 7); AUC above 0.5 means higher in Hi. Header: AUC with its stratified bootstrap 95% interval (2000 resamples), Wilcoxon rank−sum p (smallest attainable 0.00058), BH q within task, and the fea

Hi vs Lo, training change


123 of 1900 proteins at p < 0.05, 0 at BH < 0.05. Page 1 of 2.



Hi (up to 7) against Lo (up to 6); AUC above 0.5 means higher in Hi. Header: AUC with its stratified bootstrap 95% interval (2000 resamples), Wilcoxon rank−sum p (smallest attainable 0.0012), BH q within task, and the featu

Hi vs Lo, training change


123 of 1900 proteins at p < 0.05, 0 at BH < 0.05. Page 2 of 2.



Hi (up to 7) against Lo (up to 6); AUC above 0.5 means higher in Hi. Header: AUC with its stratified bootstrap 95% interval (2000 resamples), Wilcoxon rank−sum p (smallest attainable 0.0012), BH q within task, and the featu

Hi vs Lo, acute change


26 of 1900 proteins at p < 0.05, 0 at BH < 0.05. Page 1 of 2.



Hi (up to 7) against Lo (up to 7); AUC above 0.5 means higher in Hi. Header: AUC with its stratified bootstrap 95% interval (2000 resamples), Wilcoxon rank−sum p (smallest attainable 0.00058), BH q within task, and the fea

Hi vs Lo, acute change

26 of 1900 proteins at p < 0.05, 0 at BH < 0.05. Page 2 of 2.



Hi (up to 7) against Lo (up to 7); AUC above 0.5 means higher in Hi. Header: AUC with its stratified bootstrap 95% interval (2000 resamples), Wilcoxon rank−sum p (smallest attainable 0.00058), BH q within task, and the fea