
fcsa_mixed, between−person term


67 of 1640 proteins at p < 0.05, 0 at BH < 0.05.



limma on 30 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

fcsa_mixed, between−person term


67 of 1640 proteins at p < 0.05, 0 at BH < 0.05.



limma on 30 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

fcsa_mixed, between−person term


67 of 1640 proteins at p < 0.05, 0 at BH < 0.05.



limma on 30 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

fcsa_mixed, between−person term


67 of 1640 proteins at p < 0.05, 0 at BH < 0.05.



limma on 30 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

fcsa_mixed, between−person term


67 of 1640 proteins at p < 0.05, 0 at BH < 0.05.



limma on 30 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

fcsa_mixed, between−person term


67 of 1640 proteins at p < 0.05, 0 at BH < 0.05.



limma on 30 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

fcsa_mixed, within−person term


61 of 1640 proteins at p < 0.05, 0 at BH < 0.05.



limma on 30 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

fcsa_mixed, within−person term


61 of 1640 proteins at p < 0.05, 0 at BH < 0.05.



limma on 30 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

fcsa_mixed, within−person term


61 of 1640 proteins at p < 0.05, 0 at BH < 0.05.



limma on 30 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

fcsa_mixed, within−person term


61 of 1640 proteins at p < 0.05, 0 at BH < 0.05.



limma on 30 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

fcsa_mixed, within−person term


61 of 1640 proteins at p < 0.05, 0 at BH < 0.05.



limma on 30 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

fcsa_mixed, within−person term


61 of 1640 proteins at p < 0.05, 0 at BH < 0.05.



limma on 30 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

fcsa_type1, between−person term


64 of 1640 proteins at p < 0.05, 0 at BH < 0.05.



limma on 30 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

fcsa_type1, between−person term


64 of 1640 proteins at p < 0.05, 0 at BH < 0.05.



limma on 30 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

fcsa_type1, between−person term


64 of 1640 proteins at p < 0.05, 0 at BH < 0.05.



limma on 30 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

fcsa_type1, between−person term


64 of 1640 proteins at p < 0.05, 0 at BH < 0.05.



limma on 30 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

fcsa_type1, between−person term


64 of 1640 proteins at p < 0.05, 0 at BH < 0.05.



limma on 30 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

fcsa_type1, between−person term


64 of 1640 proteins at p < 0.05, 0 at BH < 0.05.



limma on 30 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

fcsa_type1, within−person term


59 of 1640 proteins at p < 0.05, 0 at BH < 0.05.



limma on 30 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

fcsa_type1, within−person term


59 of 1640 proteins at p < 0.05, 0 at BH < 0.05.



limma on 30 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

fcsa_type1, within−person term


59 of 1640 proteins at p < 0.05, 0 at BH < 0.05.



limma on 30 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

fcsa_type1, within−person term


59 of 1640 proteins at p < 0.05, 0 at BH < 0.05.



limma on 30 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

fcsa_type1, within−person term


59 of 1640 proteins at p < 0.05, 0 at BH < 0.05.



limma on 30 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

fcsa_type2, between−person term


64 of 1640 proteins at p < 0.05, 0 at BH < 0.05.



limma on 30 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

fcsa_type2, between−person term


64 of 1640 proteins at p < 0.05, 0 at BH < 0.05.



limma on 30 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

fcsa_type2, between−person term


64 of 1640 proteins at p < 0.05, 0 at BH < 0.05.



limma on 30 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

fcsa_type2, between−person term


64 of 1640 proteins at p < 0.05, 0 at BH < 0.05.



limma on 30 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

fcsa_type2, between−person term


64 of 1640 proteins at p < 0.05, 0 at BH < 0.05.



limma on 30 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

fcsa_type2, between−person term


64 of 1640 proteins at p < 0.05, 0 at BH < 0.05.



limma on 30 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

fcsa_type2, within−person term


74 of 1640 proteins at p < 0.05, 0 at BH < 0.05.



limma on 30 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

fcsa_type2, within−person term


74 of 1640 proteins at p < 0.05, 0 at BH < 0.05.



limma on 30 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

fcsa_type2, within−person term


74 of 1640 proteins at p < 0.05, 0 at BH < 0.05.



limma on 30 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

fcsa_type2, within−person term


74 of 1640 proteins at p < 0.05, 0 at BH < 0.05.



limma on 30 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

fcsa_type2, within−person term


74 of 1640 proteins at p < 0.05, 0 at BH < 0.05.



limma on 30 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

fcsa_type2, within−person term


74 of 1640 proteins at p < 0.05, 0 at BH < 0.05.



limma on 30 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

fcsa_type2, within−person term


74 of 1640 proteins at p < 0.05, 0 at BH < 0.05.



limma on 30 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

fibres_mixed, between−person term


83 of 1640 proteins at p < 0.05, 0 at BH < 0.05.



limma on 30 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

fibres_mixed, between−person term


83 of 1640 proteins at p < 0.05, 0 at BH < 0.05.



limma on 30 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

fibres_mixed, between−person term


83 of 1640 proteins at p < 0.05, 0 at BH < 0.05.



limma on 30 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

fibres_mixed, between−person term


83 of 1640 proteins at p < 0.05, 0 at BH < 0.05.



limma on 30 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

fibres_mixed, between−person term


83 of 1640 proteins at p < 0.05, 0 at BH < 0.05.



limma on 30 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

fibres_mixed, between−person term


83 of 1640 proteins at p < 0.05, 0 at BH < 0.05.



limma on 30 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

fibres_mixed, between−person term


83 of 1640 proteins at p < 0.05, 0 at BH < 0.05.



limma on 30 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

fibres_mixed, within−person term


46 of 1640 proteins at p < 0.05, 0 at BH < 0.05.



limma on 30 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

fibres_mixed, within−person term


46 of 1640 proteins at p < 0.05, 0 at BH < 0.05.



limma on 30 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

fibres_mixed, within−person term


46 of 1640 proteins at p < 0.05, 0 at BH < 0.05.



limma on 30 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

fibres_mixed, within−person term


46 of 1640 proteins at p < 0.05, 0 at BH < 0.05.



limma on 30 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

fibres_type1, between−person term


63 of 1640 proteins at p < 0.05, 0 at BH < 0.05.



limma on 30 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

fibres_type1, between−person term


63 of 1640 proteins at p < 0.05, 0 at BH < 0.05.



limma on 30 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

fibres_type1, between−person term


63 of 1640 proteins at p < 0.05, 0 at BH < 0.05.



limma on 30 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

fibres_type1, between−person term


63 of 1640 proteins at p < 0.05, 0 at BH < 0.05.



limma on 30 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

fibres_type1, between−person term


63 of 1640 proteins at p < 0.05, 0 at BH < 0.05.



limma on 30 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

fibres_type1, between−person term


63 of 1640 proteins at p < 0.05, 0 at BH < 0.05.



limma on 30 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

fibres_type1, within−person term


52 of 1640 proteins at p < 0.05, 0 at BH < 0.05.



limma on 30 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

fibres_type1, within−person term


52 of 1640 proteins at p < 0.05, 0 at BH < 0.05.



limma on 30 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

fibres_type1, within−person term


52 of 1640 proteins at p < 0.05, 0 at BH < 0.05.



limma on 30 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

fibres_type1, within−person term


52 of 1640 proteins at p < 0.05, 0 at BH < 0.05.



limma on 30 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

fibres_type1, within−person term


52 of 1640 proteins at p < 0.05, 0 at BH < 0.05.



limma on 30 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

leg_ext_1rm, between−person term


130 of 1615 proteins at p < 0.05, 0 at BH < 0.05.



limma on 29 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

leg_ext_1rm, between−person term


130 of 1615 proteins at p < 0.05, 0 at BH < 0.05.



limma on 29 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

leg_ext_1rm, between−person term


130 of 1615 proteins at p < 0.05, 0 at BH < 0.05.



limma on 29 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

leg_ext_1rm, between−person term


130 of 1615 proteins at p < 0.05, 0 at BH < 0.05.



limma on 29 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

leg_ext_1rm, between−person term


130 of 1615 proteins at p < 0.05, 0 at BH < 0.05.



limma on 29 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

leg_ext_1rm, between−person term


130 of 1615 proteins at p < 0.05, 0 at BH < 0.05.



limma on 29 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

leg_ext_1rm, between−person term


130 of 1615 proteins at p < 0.05, 0 at BH < 0.05.



limma on 29 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

leg_ext_1rm, between−person term


130 of 1615 proteins at p < 0.05, 0 at BH < 0.05.



limma on 29 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

leg_ext_1rm, between−person term


130 of 1615 proteins at p < 0.05, 0 at BH < 0.05.



limma on 29 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

leg_ext_1rm, between−person term


130 of 1615 proteins at p < 0.05, 0 at BH < 0.05.



limma on 29 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

leg_ext_1rm, between−person term


130 of 1615 proteins at p < 0.05, 0 at BH < 0.05.



limma on 29 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

leg_ext_1rm, within−person term


122 of 1615 proteins at p < 0.05, 0 at BH < 0.05.



limma on 29 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

leg_ext_1rm, within−person term


122 of 1615 proteins at p < 0.05, 0 at BH < 0.05.



limma on 29 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

leg_ext_1rm, within−person term


122 of 1615 proteins at p < 0.05, 0 at BH < 0.05.



limma on 29 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

leg_ext_1rm, within−person term


122 of 1615 proteins at p < 0.05, 0 at BH < 0.05.



limma on 29 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

leg_ext_1rm, within−person term


122 of 1615 proteins at p < 0.05, 0 at BH < 0.05.



limma on 29 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

leg_ext_1rm, within−person term


122 of 1615 proteins at p < 0.05, 0 at BH < 0.05.



limma on 29 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

leg_ext_1rm, within−person term


122 of 1615 proteins at p < 0.05, 0 at BH < 0.05.



limma on 29 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

leg_ext_1rm, within−person term


122 of 1615 proteins at p < 0.05, 0 at BH < 0.05.



limma on 29 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

leg_ext_1rm, within−person term


122 of 1615 proteins at p < 0.05, 0 at BH < 0.05.



limma on 29 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

leg_ext_1rm, within−person term


122 of 1615 proteins at p < 0.05, 0 at BH < 0.05.



limma on 29 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

leg_ext_1rm, within−person term


122 of 1615 proteins at p < 0.05, 0 at BH < 0.05.



limma on 29 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

leg_press_1rm, between−person term


107 of 1640 proteins at p < 0.05, 0 at BH < 0.05.



limma on 30 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

leg_press_1rm, between−person term


107 of 1640 proteins at p < 0.05, 0 at BH < 0.05.



limma on 30 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

leg_press_1rm, between−person term


107 of 1640 proteins at p < 0.05, 0 at BH < 0.05.



limma on 30 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

leg_press_1rm, between−person term


107 of 1640 proteins at p < 0.05, 0 at BH < 0.05.



limma on 30 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

leg_press_1rm, between−person term


107 of 1640 proteins at p < 0.05, 0 at BH < 0.05.



limma on 30 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

leg_press_1rm, between−person term


107 of 1640 proteins at p < 0.05, 0 at BH < 0.05.



limma on 30 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

leg_press_1rm, between−person term


107 of 1640 proteins at p < 0.05, 0 at BH < 0.05.



limma on 30 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

leg_press_1rm, between−person term


107 of 1640 proteins at p < 0.05, 0 at BH < 0.05.



limma on 30 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

leg_press_1rm, between−person term


107 of 1640 proteins at p < 0.05, 0 at BH < 0.05.



limma on 30 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

leg_press_1rm, within−person term


82 of 1640 proteins at p < 0.05, 0 at BH < 0.05.



limma on 30 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

leg_press_1rm, within−person term


82 of 1640 proteins at p < 0.05, 0 at BH < 0.05.



limma on 30 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

leg_press_1rm, within−person term


82 of 1640 proteins at p < 0.05, 0 at BH < 0.05.



limma on 30 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

leg_press_1rm, within−person term


82 of 1640 proteins at p < 0.05, 0 at BH < 0.05.



limma on 30 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

leg_press_1rm, within−person term


82 of 1640 proteins at p < 0.05, 0 at BH < 0.05.



limma on 30 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

leg_press_1rm, within−person term


82 of 1640 proteins at p < 0.05, 0 at BH < 0.05.



limma on 30 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

leg_press_1rm, within−person term


82 of 1640 proteins at p < 0.05, 0 at BH < 0.05.



limma on 30 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

mcsa, between−person term


206 of 1640 proteins at p < 0.05, 0 at BH < 0.05.



limma on 30 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

mcsa, between−person term


206 of 1640 proteins at p < 0.05, 0 at BH < 0.05.



limma on 30 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

mcsa, between−person term


206 of 1640 proteins at p < 0.05, 0 at BH < 0.05.



limma on 30 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

mcsa, between−person term


206 of 1640 proteins at p < 0.05, 0 at BH < 0.05.



limma on 30 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

mcsa, between−person term


206 of 1640 proteins at p < 0.05, 0 at BH < 0.05.



limma on 30 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

mcsa, between−person term


206 of 1640 proteins at p < 0.05, 0 at BH < 0.05.



limma on 30 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

mcsa, between−person term


206 of 1640 proteins at p < 0.05, 0 at BH < 0.05.



limma on 30 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

mcsa, between−person term


206 of 1640 proteins at p < 0.05, 0 at BH < 0.05.



limma on 30 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

mcsa, between−person term


206 of 1640 proteins at p < 0.05, 0 at BH < 0.05.



limma on 30 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

mcsa, between−person term


206 of 1640 proteins at p < 0.05, 0 at BH < 0.05.



limma on 30 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

mcsa, between−person term


206 of 1640 proteins at p < 0.05, 0 at BH < 0.05.



limma on 30 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

mcsa, between−person term


206 of 1640 proteins at p < 0.05, 0 at BH < 0.05.



limma on 30 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

mcsa, between−person term


206 of 1640 proteins at p < 0.05, 0 at BH < 0.05.



limma on 30 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

mcsa, between−person term


206 of 1640 proteins at p < 0.05, 0 at BH < 0.05.



limma on 30 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

mcsa, between−person term


206 of 1640 proteins at p < 0.05, 0 at BH < 0.05.



limma on 30 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

mcsa, between−person term


206 of 1640 proteins at p < 0.05, 0 at BH < 0.05.



limma on 30 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

mcsa, between−person term


206 of 1640 proteins at p < 0.05, 0 at BH < 0.05.



limma on 30 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

mcsa, between−person term


206 of 1640 proteins at p < 0.05, 0 at BH < 0.05.



limma on 30 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

mcsa, within−person term


63 of 1640 proteins at p < 0.05, 0 at BH < 0.05.



limma on 30 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

mcsa, within−person term


63 of 1640 proteins at p < 0.05, 0 at BH < 0.05.



limma on 30 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

mcsa, within−person term


63 of 1640 proteins at p < 0.05, 0 at BH < 0.05.



limma on 30 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

mcsa, within−person term


63 of 1640 proteins at p < 0.05, 0 at BH < 0.05.



limma on 30 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

mcsa, within−person term


63 of 1640 proteins at p < 0.05, 0 at BH < 0.05.



limma on 30 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

mcsa, within−person term


63 of 1640 proteins at p < 0.05, 0 at BH < 0.05.



limma on 30 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

type1_share, between−person term


51 of 1640 proteins at p < 0.05, 0 at BH < 0.05.



limma on 30 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

type1_share, between−person term


51 of 1640 proteins at p < 0.05, 0 at BH < 0.05.



limma on 30 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

type1_share, between−person term


51 of 1640 proteins at p < 0.05, 0 at BH < 0.05.



limma on 30 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

type1_share, between−person term


51 of 1640 proteins at p < 0.05, 0 at BH < 0.05.



limma on 30 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

type1_share, between−person term


51 of 1640 proteins at p < 0.05, 0 at BH < 0.05.



limma on 30 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

type1_share, within−person term


56 of 1640 proteins at p < 0.05, 0 at BH < 0.05.



limma on 30 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

type1_share, within−person term


56 of 1640 proteins at p < 0.05, 0 at BH < 0.05.



limma on 30 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

type1_share, within−person term


56 of 1640 proteins at p < 0.05, 0 at BH < 0.05.



limma on 30 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

type1_share, within−person term


56 of 1640 proteins at p < 0.05, 0 at BH < 0.05.



limma on 30 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

type1_share, within−person term


56 of 1640 proteins at p < 0.05, 0 at BH < 0.05.



limma on 30 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 protein−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window

1145 of 28204 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window

1145 of 28204 protein−outcome pairs at p < 0.05, 1 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.