

a Can one feature tell HR from LR?
Best feature per task (smallest Wilcoxon p), AUC with 95% CI; right: that feature, then features at p < 0.05 / expected by chance
Proteins
Sets
Modules
Training
Acute
Between groups
T2 - T1
T3 - T2
T3
T2
T1
VDAC3 50 / 95
FSCN1 67 / 95
EIF4A2 100 / 95
EIF3G 57 / 95
HDHD2 44 / 95
Nucleoside phos... 49 / 69
Resolution of s... 32 / 69
Endoplasmicret... 52 / 69
Response to virus 12 / 69
Cytoskeleton de... 24 / 69
midnightblue 0 / 1
yellow 0 / 1
cyan 0 / 1
blue 0 / 1
midnightblue 0 / 1
0.0
0.5
1.0
AUC
higher in HR
higher in LR
95% CI
bootstrap
DeLong
b Group difference against the continuous interaction
leg press 1RM (% change), training window, 1900 proteins; Spearman across proteins 0.88 (descriptive)
t, growth x strength (continuous)
t, HR minus LR slope (group x trait)
TNNT1
TTN
TPM3
MYL2
MYBPC2
MYOT
MYH7
MYH1
HR - LR
FDR < 0.05
p < 0.05 only
c The same proteins, within each group
Spearman rho with leg press 1RM (% change), training window; 11 of 11 group-difference hits change sign
rho in LR
rho in HR
MYH1
MYOT
MYH7
MYL2
MYBPC2
MYOM1
MYL6
TPM3
TTN
TNNT1
NEB
d TPM3, person by person
Group difference in slope: t = -6.1, p < 0.001, FDR = 0.027
leg press 1RM (% change)
TPM3, training change (log2)
HR: rho -0.94 (n = 6)
LR: rho 0.60 (n = 8)