


Protein association hits, acute change (T3 – T2) (1/6)


Spearman against each phenotype; 435 proteins nominal in at least one column; rows 1–75



Rows: proteins at nominal p < 0.05 against at least one phenotype, most columns first, then by best p. Filled dot: nominal, fill = Spearman rho, size = -log10 p. Black ring: BH < 0.05. Grey speck: tested, not nominal. Table: c_data/03_phenotyp

Protein association hits, acute change (T3 – T2) (2/6)


Spearman against each phenotype; 435 proteins nominal in at least one column; rows 76–150



Rows: proteins at nominal $p < 0.05$ against at least one phenotype, most columns first, then by best p . Filled dot: nominal, fill = Spearman ρ , size = $-\log_{10} p$. Black ring: BH < 0.05 . Grey speck: tested, not nominal. Table: c_data/03_phenotype.


Protein association hits, acute change (T3 – T2) (3/6)

Spearman against each phenotype; 435 proteins nominal in at least one column; rows 151–225



Protein association hits, acute change (T3 – T2) (4/6)


Spearman against each phenotype; 435 proteins nominal in at least one column; rows 226–300




Rows: proteins at nominal p < 0.05 against at least one phenotype, most columns first, then by best p. Filled dot: nominal, fill = Spearman rho, size = -log10 p. Black ring: BH < 0.05. Grey speck: tested, not nominal. Table: c_data/03_phenotype

Protein association hits, acute change (T3 – T2) (5/6)

Spearman against each phenotype; 435 proteins nominal in at least one column; rows 301–375



Spearman against each phenotype; 435 proteins nominal in at least one column; rows 376–435



Rows: proteins at nominal $p < 0.05$ against at least one phenotype, most columns first, then by best p . Filled dot: nominal, fill = Spearman rho, size = $-\log_{10} p$. Black ring: BH < 0.05 . Grey speck: tested, not nominal. Table: c_data/03_phenotype.x