



NES concordance, all collections: Training_HR against Training_LR

fgsea NES per contrast, 1369 sets, BH within contrast


A



B



C




genes ● 50 ● 150 ● 300 significant in ● Both ● Training_HR ● Training_LR

One point per gene set, scored in both training contrasts. Dashed line is identity; grey points reach neither threshold. Discordant sets are significant in at least one arm and sit on opposite sides of zero. Panel C rescales them. Table: c_data/nescatter.csv.


NES concordance, Hallmark and GO Slim: Training_HR against Training_LR

fgsea NES per contrast, 90 non-nesting sets, BH within contrast


A



B



C




genes ● 50 ● 150 ● 300 significant in ● Both ● Training_HR ● Training_LR

Panel A is every Hallmark and GO Slim set; B and C rescale the significant sets with the same sign in both arms, so each can be named. Point size is gene count, colour the contrast a set reached FDR 0.05 in. Table: [c_data/nesc_scatter.csv](#).


NES concordance, all collections: Acute_HR against Acute_LR

fgsea NES per contrast, 1370 sets, BH within contrast


A



B



C




One point per gene set, scored in both acute contrasts. Dashed line is identity; grey points reach neither threshold. Discordant sets are significant in at least one arm and sit on opposite sides of zero. Panel C rescales them. Table: [c_data/nesc_scatter.csv](#).


NES concordance, Hallmark and GO Slim: Acute_HR against Acute_LR

fgsea NES per contrast, 90 non-nesting sets, BH within contrast


A



B



C



genes ● 50 ● 150 ● 300 significant in ● Both ● Acute_HR ● Acute_LR

Panel A is every Hallmark and GO Slim set; B and C rescale the significant sets with the same sign in both arms, so each can be named. Point size is gene count, colour the contrast a set reached FDR 0.05 in. Table: [c_data/nesc_scatter.csv](#).