


Training, HR (T1 to T2)


singscore per set, 6 paired subjects, AUC with direction fixed



All 50 sets reaching nominal p, by collection then p. Red separates higher in T2, blue lower. Shading is area under the curve, dashed line chance. p from the Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Training, HR (T1 to T2)


singscore per set, 6 paired subjects, AUC with direction fixed



All 50 sets reaching nominal p, by collection then p. Red separates higher in T2, blue lower. Shading is area under the curve, dashed line chance. p from the Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Training, HR (T1 to T2)


singscore per set, 6 paired subjects, AUC with direction fixed



All 50 sets reaching nominal p, by collection then p. Red separates higher in T2, blue lower. Shading is area under the curve, dashed line chance. p from the Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Training, HR (T1 to T2)


singscore per set, 6 paired subjects, AUC with direction fixed



All 50 sets reaching nominal p, by collection then p. Red separates higher in T2, blue lower. Shading is area under the curve, dashed line chance. p from the Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Training, LR (T1 to T2)


singscore per set, 8 paired subjects, AUC with direction fixed



All 30 sets reaching nominal p, by collection then p. Red separates higher in T2, blue lower. Shading is area under the curve, dashed line chance. p from the Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Training, LR (T1 to T2)


singscore per set, 8 paired subjects, AUC with direction fixed



All 30 sets reaching nominal p, by collection then p. Red separates higher in T2, blue lower. Shading is area under the curve, dashed line chance. p from the Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Acute bout, HR (T2 to T3)


singscore per set, 7 paired subjects, AUC with direction fixed



All 113 sets reaching nominal p, by collection then p. Red separates higher in T3, blue lower. Shading is area under the curve, dashed line chance. p from the Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Acute bout, HR (T2 to T3)


singscore per set, 7 paired subjects, AUC with direction fixed



All 113 sets reaching nominal p, by collection then p. Red separates higher in T3, blue lower. Shading is area under the curve, dashed line chance. p from the Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Acute bout, HR (T2 to T3)


singscore per set, 7 paired subjects, AUC with direction fixed



All 113 sets reaching nominal p, by collection then p. Red separates higher in T3, blue lower. Shading is area under the curve, dashed line chance. p from the Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Acute bout, HR (T2 to T3)


singscore per set, 7 paired subjects, AUC with direction fixed



All 113 sets reaching nominal p, by collection then p. Red separates higher in T3, blue lower. Shading is area under the curve, dashed line chance. p from the Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Acute bout, HR (T2 to T3)


singscore per set, 7 paired subjects, AUC with direction fixed



All 113 sets reaching nominal p, by collection then p. Red separates higher in T3, blue lower. Shading is area under the curve, dashed line chance. p from the Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Acute bout, HR (T2 to T3)


singscore per set, 7 paired subjects, AUC with direction fixed



All 113 sets reaching nominal p, by collection then p. Red separates higher in T3, blue lower. Shading is area under the curve, dashed line chance. p from the Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Acute bout, HR (T2 to T3)


singscore per set, 7 paired subjects, AUC with direction fixed



All 113 sets reaching nominal p, by collection then p. Red separates higher in T3, blue lower. Shading is area under the curve, dashed line chance. p from the Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Acute bout, HR (T2 to T3)


singscore per set, 7 paired subjects, AUC with direction fixed



All 113 sets reaching nominal p, by collection then p. Red separates higher in T3, blue lower. Shading is area under the curve, dashed line chance. p from the Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Acute bout, LR (T2 to T3)


singscore per set, 8 paired subjects, AUC with direction fixed



All 153 sets reaching nominal p, by collection then p. Red separates higher in T3, blue lower. Shading is area under the curve, dashed line chance, p from the Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Acute bout, LR (T2 to T3)


singscore per set, 8 paired subjects, AUC with direction fixed



All 153 sets reaching nominal p, by collection then p. Red separates higher in T3, blue lower. Shading is area under the curve, dashed line chance, p from the Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Acute bout, LR (T2 to T3)


singscore per set, 8 paired subjects, AUC with direction fixed



All 153 sets reaching nominal p, by collection then p. Red separates higher in T3, blue lower. Shading is area under the curve, dashed line chance, p from the Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Acute bout, LR (T2 to T3)


singscore per set, 8 paired subjects, AUC with direction fixed



All 153 sets reaching nominal p, by collection then p. Red separates higher in T3, blue lower. Shading is area under the curve, dashed line chance, p from the Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Acute bout, LR (T2 to T3)


singscore per set, 8 paired subjects, AUC with direction fixed



All 153 sets reaching nominal p, by collection then p. Red separates higher in T3, blue lower. Shading is area under the curve, dashed line chance, p from the Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Acute bout, LR (T2 to T3)


singscore per set, 8 paired subjects, AUC with direction fixed



All 153 sets reaching nominal p, by collection then p. Red separates higher in T3, blue lower. Shading is area under the curve, dashed line chance. p from the Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Acute bout, LR (T2 to T3)


singscore per set, 8 paired subjects, AUC with direction fixed



All 153 sets reaching nominal p, by collection then p. Red separates higher in T3, blue lower. Shading is area under the curve, dashed line chance. p from the Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Acute bout, LR (T2 to T3)


singscore per set, 8 paired subjects, AUC with direction fixed



All 153 sets reaching nominal p, by collection then p. Red separates higher in T3, blue lower. Shading is area under the curve, dashed line chance. p from the Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Acute bout, LR (T2 to T3)


singscore per set, 8 paired subjects, AUC with direction fixed



All 153 sets reaching nominal p, by collection then p. Red separates higher in T3, blue lower. Shading is area under the curve, dashed line chance. p from the Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Acute bout, LR (T2 to T3)


singscore per set, 8 paired subjects, AUC with direction fixed



All 153 sets reaching nominal p, by collection then p. Red separates higher in T3, blue lower. Shading is area under the curve, dashed line chance. p from the Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.


HR vs LR at T2

singscore per set, 7 subjects, AUC with direction fixed



HR vs LR, training change


singscore per set, 6 subjects, AUC with direction fixed



All 37 sets reaching nominal p, by collection then p. Red separates higher in HR, blue lower. Shading is area under the curve, dashed line chance, p from the Wilcoxon rank-sum test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

HR vs LR, training change


singscore per set, 6 subjects, AUC with direction fixed



All 37 sets reaching nominal p, by collection then p. Red separates higher in HR, blue lower. Shading is area under the curve, dashed line chance, p from the Wilcoxon rank-sum test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

HR vs LR, training change


singscore per set, 6 subjects, AUC with direction fixed



All 37 sets reaching nominal p, by collection then p. Red separates higher in HR, blue lower. Shading is area under the curve, dashed line chance. p from the Wilcoxon rank-sum test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

HR vs LR, acute change


singscore per set, 7 subjects, AUC with direction fixed



All 56 sets reaching nominal p, by collection then p. Red separates higher in HR, blue lower. Shading is area under the curve, dashed line chance, p from the Wilcoxon rank-sum test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

HR vs LR, acute change


singscore per set, 7 subjects, AUC with direction fixed



All 56 sets reaching nominal p, by collection then p. Red separates higher in HR, blue lower. Shading is area under the curve, dashed line chance, p from the Wilcoxon rank-sum test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

HR vs LR, acute change


singscore per set, 7 subjects, AUC with direction fixed



All 56 sets reaching nominal p, by collection then p. Red separates higher in HR, blue lower. Shading is area under the curve, dashed line chance, p from the Wilcoxon rank-sum test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

HR vs LR, acute change

singscore per set, 7 subjects, AUC with direction fixed




All 56 sets reaching nominal p, by collection then p. Red separates higher in HR, blue lower. Shading is area under the curve, dashed line chance. p from the Wilcoxon rank-sum test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, training change (T2 – T1)

Spearman, subjects pooled across arms

HR LR




All 518 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, training change (T2 – T1)

Spearman, subjects pooled across arms

HR LR



All 518 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, training change (T2 – T1)

Spearman, subjects pooled across arms

HR LR




All 518 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, training change (T2 – T1)

Spearman, subjects pooled across arms

HR LR



All 518 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, training change (T2 – T1)

Spearman, subjects pooled across arms


HR LR



All 518 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, training change (T2 – T1)



Spearman, subjects pooled across arms



All 518 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, training change (T2 – T1)



Spearman, subjects pooled across arms



All 518 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, training change (T2 – T1)



Spearman, subjects pooled across arms



All 518 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, training change (T2 – T1)

Spearman, subjects pooled across arms




All 518 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, training change (T2 – T1)

Spearman, subjects pooled across arms


HR LR



All 518 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, training change (T2 – T1)

Spearman, subjects pooled across arms




All 518 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, training change (T2 – T1)

Spearman, subjects pooled across arms

HR LR




All 518 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, training change (T2 – T1)

Spearman, subjects pooled across arms



HR LR



All 518 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, training change (T2 – T1)

Spearman, subjects pooled across arms




All 518 set–outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, training change (T2 – T1)

Spearman, subjects pooled across arms

HR LR




All 518 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, training change (T2 – T1)

Spearman, subjects pooled across arms

HR LR




All 518 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, training change (T2 – T1)

Spearman, subjects pooled across arms

HR LR




All 518 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, training change (T2 – T1)

Spearman, subjects pooled across arms



HR LR



All 518 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, training change (T2 – T1)



Spearman, subjects pooled across arms



All 518 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, training change (T2 – T1)

Spearman, subjects pooled across arms




All 518 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, training change (T2 – T1)

Spearman, subjects pooled across arms


HR LR



All 518 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, training change (T2 – T1)

Spearman, subjects pooled across arms



All 518 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, training change (T2 – T1)

Spearman, subjects pooled across arms

HR LR




All 518 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, training change (T2 – T1)

Spearman, subjects pooled across arms

HR LR




All 518 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, training change (T2 – T1)

Spearman, subjects pooled across arms

HR LR




All 518 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, training change (T2 – T1)

Spearman, subjects pooled across arms

HR LR




All 518 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, training change (T2 – T1)

Spearman, subjects pooled across arms



HR LR



All 518 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, training change (T2 – T1)

Spearman, subjects pooled across arms



All 518 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, training change (T2 – T1)

Spearman, subjects pooled across arms


HR LR



All 518 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, training change (T2 – T1)

Spearman, subjects pooled across arms




All 518 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, training change (T2 – T1)

Spearman, subjects pooled across arms


HR LR



All 518 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, training change (T2 – T1)

Spearman, subjects pooled across arms




All 518 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, training change (T2 – T1)

Spearman, subjects pooled across arms



HR LR



All 518 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, level at T1

Spearman, subjects pooled across arms




All 576 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, level at T1

Spearman, subjects pooled across arms


HR LR



All 576 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, level at T1

Spearman, subjects pooled across arms




All 576 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, level at T1

Spearman, subjects pooled across arms



HR LR



All 576 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, level at T1



Spearman, subjects pooled across arms



All 576 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, level at T1



Spearman, subjects pooled across arms



All 576 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, level at T1



Spearman, subjects pooled across arms



All 576 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, level at T1



Spearman, subjects pooled across arms



All 576 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, level at T1

Spearman, subjects pooled across arms




All 576 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, level at T1

Spearman, subjects pooled across arms

HR LR




All 576 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, level at T1

Spearman, subjects pooled across arms

HR LR




All 576 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, level at T1

Spearman, subjects pooled across arms

HR LR




All 576 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, level at T1

Spearman, subjects pooled across arms


HR LR



All 576 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, level at T1

Spearman, subjects pooled across arms



GOBP: Positive Reg.

Transmembrane

Transport vs d_nfbre_l

r = -0.55 p 0.033 q 0.99

GOBP: Reg. Kinase

Activity vs comp_hypertrophy

r = +0.55 p 0.034 q 0.99

GOBP: Synaptic

Vesicle

Recycling vs d_mcsa

r = +0.55 p 0.034 q 0.99

GOBP: Nuclear

Transcribed

mRNA Catabolism vs volume_load

r = +0.55 p 0.034 q 0.84

GOBP: Resp. to

Oxidative

Stress vs d_fcsc_l

r = -0.55 p 0.034 q 1

GOBP: Reg. Cardiac

Muscle

Contraction

By Reg. The

Release Of

Sequestered

Calcium Ion vs volume_load

r = -0.55 p 0.034 q 0.84

GOBP: Reg. Cell

Substrate

Adhesion vs d_1rm_ext

r = +0.57 p 0.034 q 0.45

GOBP: Vesicle

Localization vs d_1rm_ext

r = +0.57 p 0.034 q 0.45

GOBP: Reg. Monoatomic

Ion Transport vs d_mcsa

r = -0.55 p 0.035 q 0.99

GOBP: Golgi Org. vs d_1rm_ext

r = -0.56 p 0.036 q 0.45

GOBP: Mitophagy vs d_1rm_ext

r = -0.56 p 0.036 q 0.45

GOBP: Positive Reg.

WNT Signaling vs d_1rm_ext

r = +0.56 p 0.036 q 0.45

GOBP: Reg.

Cytoplasmic

Translation vs d_1rm_legpress

r = +0.54 p 0.036 q 0.53

GOBP: Cellular Resp.

to Chemical

Stress vs d_fcsc_l

r = -0.54 p 0.037 q 1

GOBP: Reg. Actomyosin

Structure Org. vs d_fcsc_l

r = +0.54 p 0.037 q 1

GOBP: Reg. Cell

Substrate

Junction Org. vs d_fcsc_mixed

r = +0.54 p 0.037 q 1

GOBP: Positive Reg.

Transmembrane

Transport vs d_nfbre_l

r = -0.55 p 0.033 q 0.99

GOBP: Reg. Kinase

Activity vs comp_hypertrophy

r = +0.55 p 0.034 q 0.99

GOBP: Synaptic

Vesicle

Recycling vs d_mcsa

r = +0.55 p 0.034 q 0.99

GOBP: Nuclear

Transcribed

mRNA Catabolism vs volume_load

r = +0.55 p 0.034 q 0.84

GOBP: Resp. to

Oxidative

Stress vs d_fcsc_l

r = -0.55 p 0.034 q 1

GOBP: Reg. Cardiac

Muscle

Contraction

By Reg. The

Release Of

Sequestered

Calcium Ion vs volume_load

r = -0.55 p 0.034 q 0.84

GOBP: Reg. Cell

Substrate

Adhesion vs d_1rm_ext

r = +0.57 p 0.034 q 0.45

GOBP: Vesicle

Localization vs d_1rm_ext

r = +0.57 p 0.034 q 0.45

GOBP: Reg. Monoatomic

Ion Transport vs d_mcsa

r = -0.55 p 0.035 q 0.99

GOBP: Golgi Org. vs d_1rm_ext

r = -0.56 p 0.036 q 0.45

GOBP: Mitophagy vs d_1rm_ext

r = -0.56 p 0.036 q 0.45

GOBP: Positive Reg.

WNT Signaling vs d_1rm_ext

r = +0.56 p 0.036 q 0.45

GOBP: Reg.

Cytoplasmic

Translation vs d_1rm_legpress

r = +0.54 p 0.036 q 0.53

GOBP: Cellular Resp.

to Chemical

Stress vs d_fcsc_l

r = -0.54 p 0.037 q 1

GOBP: Reg. Actomyosin

Structure Org. vs d_fcsc_l

r = +0.54 p 0.037 q 1

GOBP: Reg. Cell

Substrate

Junction Org. vs d_fcsc_mixed

r = +0.54 p 0.037 q 1

GOBP: Positive Reg.

Transmembrane

Transport vs d_nfbre_l

r = -0.55 p 0.033 q 0.99

GOBP: Reg. Kinase

Activity vs comp_hypertrophy

r = +0.55 p 0.034 q 0.99

GOBP: Synaptic

Vesicle

Recycling vs d_mcsa

r = +0.55 p 0.034 q 0.99

GOBP: Nuclear

Transcribed

mRNA Catabolism vs volume_load

r = +0.55 p 0.034 q 0.84

GOBP: Resp. to

Oxidative

Stress vs d_fcsc_l

r = -0.55 p 0.034 q 1

GOBP: Reg. Cardiac

Muscle

Contraction

By Reg. The

Release Of

Sequestered

Calcium Ion vs volume_load

r = -0.55 p 0.034 q 0.84

GOBP: Reg. Cell

Substrate

Adhesion vs d_1rm_ext

r = +0.57 p 0.034 q 0.45

GOBP: Vesicle

Localization vs d_1rm_ext

r = +0.57 p 0.034 q 0.45

GOBP: Reg. Monoatomic

Ion Transport vs d_mcsa

r = -0.55 p 0.035 q 0.99

GOBP: Golgi Org. vs d_1rm_ext

r = -0.56 p 0.036 q 0.45

GOBP: Mitophagy vs d_1rm_ext

r = -0.56 p 0.036 q 0.45

GOBP: Positive Reg.

WNT Signaling vs d_1rm_ext

r = +0.56 p 0.036 q 0.45

GOBP: Reg.

Cytoplasmic

Translation vs d_1rm_legpress

r = +0.54 p 0.036 q 0.53

GOBP: Cellular Resp.

to Chemical

Stress vs d_fcsc_l

r = -0.54 p 0.037 q 1

GOBP: Reg. Actomyosin

Structure Org. vs d_fcsc_l

r = +0.54 p 0.037 q 1

GOBP: Reg. Cell

Substrate

Junction Org. vs d_fcsc_mixed

r = +0.54 p 0.037 q 1

GOBP: Positive Reg.

Transmembrane

Transport vs d_nfbre_l

r = -0.55 p 0.033 q 0.99

GOBP: Reg. Kinase

Activity vs comp_hypertrophy

r = +0.55 p 0.034 q 0.99

GOBP: Synaptic

Vesicle

Recycling vs d_mcsa

r = +0.55 p 0.034 q 0.99

GOBP: Nuclear

Transcribed

mRNA Catabolism vs volume_load

r = +0.55 p 0.034 q 0.84

GOBP: Resp. to

Oxidative

Stress vs d_fcsc_l

r = -0.55 p 0.034 q 1

GOBP: Reg. Cardiac

Muscle

Contraction

By Reg. The

Release Of

Sequestered

Calcium Ion vs volume_load

r = -0.55 p 0.034 q 0.84

GOBP: Reg. Cell

Substrate

Adhesion vs d_1rm_ext

r = +0.57 p 0.034 q 0.45

GOBP: Vesicle

Localization vs d_1rm_ext

r = +0.57 p 0.034 q 0.45

GOBP: Reg. Monoatomic

Ion Transport vs d_mcsa

r = -0.55 p 0.035 q 0.99

GOBP: Golgi Org. vs d_1rm_ext

r = -0.56 p 0.036 q 0.45

GOBP: Mitophagy vs d_1rm_ext

r = -0.56 p 0.036 q 0.45

GOBP: Positive Reg.

WNT Signaling vs d_1rm_ext

r = +0.56 p 0.036 q 0.45

GOBP: Reg.

Cytoplasmic

Translation vs d_1rm_legpress

r = +0.54 p 0.036 q 0.53

GOBP: Cellular Resp.

to Chemical

Stress vs d_fcsc_l

r = -0.54 p 0.037 q 1

GOBP: Reg. Actomyosin

Structure Org. vs d_fcsc_l

r = +0.54 p 0.037 q 1

GOBP: Reg. Cell

Substrate

Junction Org. vs d_fcsc_mixed

r = +0.54 p 0.037 q 1

GOBP: Positive Reg.

Transmembrane

Transport vs d_nfbre_l

r = -0.55 p 0.033 q 0.99

GOBP: Reg. Kinase

Activity vs comp_hypertrophy

r = +0.55 p 0.034 q 0.99

GOBP: Synaptic

Vesicle

Recycling vs d_mcsa

r = +0.55 p 0.034 q 0.99

GOBP: Nuclear

Transcribed

mRNA Catabolism vs volume_load

r = +0.55 p 0.034 q 0.84

GOBP: Resp. to

Oxidative

Stress vs d_fcsc_l

r = -0.55 p 0.034 q 1

GOBP: Reg. Cardiac

Muscle

Contraction

By Reg. The

Release Of

Sequestered

Calcium Ion vs volume_load

r = -0.55 p 0.034 q 0.84

GOBP: Reg. Cell

Substrate

Adhesion vs d_1rm_ext

r = +0.57 p 0.034 q 0.45

GOBP: Vesicle

Localization vs d_1rm_ext

r = +0.57 p 0.034 q 0.45

GOBP: Reg. Monoatomic

Ion Transport vs d_mcsa

r = -0.55 p 0.035 q 0.99

GOBP: Golgi Org. vs d_1rm_ext

r = -0.56 p 0.036 q 0.45

GOBP: Mitophagy vs d_1rm_ext

r = -0.56 p 0.036 q 0.45

GOBP: Positive Reg.

WNT Signaling vs d_1rm_ext

r = +0.56 p 0.036 q 0.45

GOBP: Reg.

Cytoplasmic

Translation vs d_1rm_legpress

r = +0.54 p 0.036 q 0.53

GOBP: Cellular Resp.

to Chemical

Stress vs d_fcsc_l

r = -0.54 p 0.037 q 1

GOBP: Reg. Actomyosin

Structure Org. vs d_fcsc_l

r = +0.54 p 0.037 q 1

Set score against phenotype, level at T1

Spearman, subjects pooled across arms



HR LR



All 576 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, level at T1

Spearman, subjects pooled across arms




All 576 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, level at T1

Spearman, subjects pooled across arms

HR LR




All 576 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, level at T1

Spearman, subjects pooled across arms

HR LR




All 576 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, level at T1

Spearman, subjects pooled across arms

HR LR




All 576 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, level at T1

Spearman, subjects pooled across arms


HR LR



All 576 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, level at T1


Spearman, subjects pooled across arms



All 576 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, level at T1


Spearman, subjects pooled across arms



All 576 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, level at T1

Spearman, subjects pooled across arms




All 576 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, level at T1

Spearman, subjects pooled across arms



HR LR



All 576 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, level at T1


Spearman, subjects pooled across arms



All 576 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, level at T1



Spearman, subjects pooled across arms




All 576 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, level at T1

Spearman, subjects pooled across arms





All 576 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.



All 576 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, level at T1



Spearman, subjects pooled across arms



All 576 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, level at T1


Spearman, subjects pooled across arms




All 576 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, level at T1


Spearman, subjects pooled across arms




Reactome: Transcriptional Regulation By Runx3 vs d_1rm_ext
r = +0.61 p 0.02 q 0.097




Reactome: Asymmetric Localization Of Pcp Proteins vs d_1rm_ext
r = +0.61 p 0.02 q 0.097




Reactome: Degradation Of Axin vs d_1rm_ext
r = +0.61 p 0.02 q 0.097




Reactome: p53 Dependent G1 DNA Damage Response vs d_1rm_ext
r = +0.61 p 0.02 q 0.097




Reactome: Reg. Runx3 Expression & Activity vs d_1rm_ext
r = +0.61 p 0.02 q 0.097




Reactome: Stabilization Of p53 vs d_1rm_ext
r = +0.61 p 0.02 q 0.097




Reactome: Downstream Tcr Signaling vs d_1rm_ext
r = +0.61 p 0.021 q 0.1




Reactome: Complex Iv Assembly vs d_1rm_legpress
r = +0.59 p 0.021 q 0.56




Reactome: Metabolism Of Carbohydrates & Carbohydrate Derivatives vs d_1rm_legpress
r = -0.59 p 0.021 q 0.56




Reactome: Cell Junction Org. vs d_1rm_ext
r = +0.61 p 0.022 q 0.1




Reactome: Glycerophospholipid Biosynthesis vs volume_load
r = +0.59 p 0.022 q 0.68




Reactome: Ampk Induced Erad & Lysosome Mediated Degradation Of PD-L1 CD274 vs d_1rm_ext
r = +0.60 p 0.022 q 0.1




Reactome: Aggrephagy vs d_mcsa
r = +0.58 p 0.023 q 0.58




Reactome: Trans Golgi Network Vesicle Budding vs d_mcsa
r = +0.58 p 0.023 q 0.58



Reactome: Hsp90 Chaperone Cycle For Steroid Hormone Receptors Shr In The Presence Of Ligand vs volume_load
r = +0.58 p 0.023 q 0.68



Reactome: Degradation Of Beta Catenin By The Destruction Complex vs d_1rm_ext
r = +0.60 p 0.023 q 0.1




set score, level at T1, one point per subject

All 576 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, level at T1



Spearman, subjects pooled across arms



All 576 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, level at T1



Spearman, subjects pooled across arms



All 576 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, level at T1

Spearman, subjects pooled across arms




All 576 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, level at T1

Spearman, subjects pooled across arms



HR LR



All 576 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, level at T1



Spearman, subjects pooled across arms



All 576 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, acute change (T3 – T2)

Spearman, subjects pooled across arms




All 511 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, acute change (T3 – T2)

Spearman, subjects pooled across arms

HR LR




All 511 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, acute change (T3 – T2)

Spearman, subjects pooled across arms

HR LR




All 511 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, acute change (T3 – T2)

Spearman, subjects pooled across arms



HR LR



All 511 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, acute change (T3 – T2)



Spearman, subjects pooled across arms



All 511 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, acute change (T3 – T2)



Spearman, subjects pooled across arms



All 511 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, acute change (T3 – T2)


Spearman, subjects pooled across arms



All 511 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, acute change (T3 – T2)



Spearman, subjects pooled across arms



All 511 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, acute change (T3 – T2)

Spearman, subjects pooled across arms




All 511 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, acute change (T3 – T2)

Spearman, subjects pooled across arms

HR LR




All 511 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, acute change (T3 – T2)

Spearman, subjects pooled across arms

HR LR




All 511 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, acute change (T3 – T2)

Spearman, subjects pooled across arms

HR LR




All 511 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, acute change (T3 – T2)

Spearman, subjects pooled across arms

HR LR



All 511 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, acute change (T3 – T2)

Spearman, subjects pooled across arms

HR LR




All 511 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, acute change (T3 – T2)

Spearman, subjects pooled across arms



HR LR



All 511 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, acute change (T3 – T2)

Spearman, subjects pooled across arms




All 511 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, acute change (T3 – T2)

Spearman, subjects pooled across arms


HR LR



All 511 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, acute change (T3 – T2)

Spearman, subjects pooled across arms




All 511 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, acute change (T3 – T2)

Spearman, subjects pooled across arms

HR LR




All 511 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, acute change (T3 – T2)

Spearman, subjects pooled across arms

HR LR




All 511 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, acute change (T3 – T2)

Spearman, subjects pooled across arms

HR LR




All 511 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, acute change (T3 – T2)

Spearman, subjects pooled across arms

HR LR




All 511 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, acute change (T3 – T2)

Spearman, subjects pooled across arms

HR LR




All 511 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, acute change (T3 – T2)

Spearman, subjects pooled across arms

HR LR




All 511 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, acute change (T3 – T2)

Spearman, subjects pooled across arms

HR LR




All 511 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, acute change (T3 – T2)

Spearman, subjects pooled across arms

HR LR




All 511 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, acute change (T3 – T2)

Spearman, subjects pooled across arms

HR LR




All 511 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, acute change (T3 – T2)

Spearman, subjects pooled across arms



HR LR



All 511 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, acute change (T3 – T2)

Spearman, subjects pooled across arms




All 511 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, acute change (T3 – T2)

Spearman, subjects pooled across arms

HR LR




All 511 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, acute change (T3 – T2)

Spearman, subjects pooled across arms

HR LR




All 511 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, acute change (T3 – T2)

Spearman, subjects pooled across arms


HR LR



All 511 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Nominal hits relative to chance, by collection

Wilcoxon per set, 5 collections, uncorrected p



Bar length is observed nominal hits divided by the count that collection returns under the null. Red clears 1, grey does not. Labels give observed of tested. Table: [c_data/05_classify_and](#)