
Proteins that respond to training or the bout
Π < 0.05 in a group's training or acute contrast; top 10 per block by p
(shared: the weaker group's p); (n/48): biopsies where the protein was observed
Training
Acute
Between groups
Interaction
Shared
MPO (15/48)
CTSG (15/48)
CHDH (9/48)
MTX1 (36/48)
HNRNPC (35/48)
PPP1R3A (30/48)
MAN2C1 (35/48)
COPB1 (43/48)
HNRNPA2B1 (48/48)
LCN2 (8/48)
HR only
MAP1LC3B2 (45/48)
AMPD3 (48/48)
SLMAP (48/48)
NNMT (16/48)
GHITM (27/48)
RDX (48/48)
HK1 (48/48)
EZR (31/48)
CAMK2B (48/48)
COMP (9/48)
LR only
LRPPRC (48/48)
LAMTOR2 (25/48)
MRPL49 (30/48)
NAMPT (47/48)
TSFM (46/48)
UBE2L3 (48/48)
GCN1 (20/48)
OXSM (42/48)
SUCLA2 (48/48)
HSPE1 (48/48)
HR
LR
HR
LR
T1
T2
T3
Training
Acute
logFC (capped at ±1.25)
1.0
0.5
0.0
-0.5
-1.0
-log10 p
1
2
4+
Π < 0.05
Π ≥ 0.05
FDR < 0.05