



Nominal hits relative to chance, by outcome and window
Spearman per protein, subjects pooled across arms



Bar length is nominal hits ($p < 0.05$) over 5% of proteins tested. The vertical line is chance. Table: c_data/03_phenotype.xlsx, chance_expectation.

Protein association hits, training change (T2 – T1) (1/7)


Spearman against each phenotype; 453 proteins nominal in at least one column; rows 1–75



Rows: proteins at nominal p < 0.05 against at least one phenotype, most columns first, then by best p. Filled dot: nominal, fill = Spearman rho, size = -log10 p. Black ring: BH < 0.05. Grey speck: tested, not nominal. Table: c_data/03_phenotype.x

Protein association hits, training change (T2 – T1) (2/7)


Spearman against each phenotype; 453 proteins nominal in at least one column; rows 76–150



Rows: proteins at nominal p < 0.05 against at least one phenotype, most columns first, then by best p. Filled dot: nominal, fill = Spearman rho, size = -log10 p. Black ring: BH < 0.05. Grey speck: tested, not nominal. Table: c_data/03_phenotype


Protein association hits, training change (T2 – T1) (3/7)

Spearman against each phenotype; 453 proteins nominal in at least one column; rows 151–225



Protein association hits, training change (T2 – T1) (4/7)


Spearman against each phenotype; 453 proteins nominal in at least one column; rows 226–300



Rows: proteins at nominal $p < 0.05$ against at least one phenotype, most columns first, then by best p . Filled dot: nominal, fill = Spearman ρ , size = $-\log_{10} p$. Black ring: BH < 0.05 . Grey speck: tested, not nominal. Table: c_data/03_phenotype.

Protein association hits, training change (T2 – T1) (5/7)


Spearman against each phenotype; 453 proteins nominal in at least one column; rows 301–375



Rows: proteins at nominal $p < 0.05$ against at least one phenotype, most columns first, then by best p . Filled dot: nominal, fill = Spearman ρ , size = $-\log_{10} p$. Black ring: BH < 0.05. Grey speck: tested, not nominal. Table: c_data/03_phenotype.


Protein association hits, training change (T2 – T1) (6/7)

Spearman against each phenotype; 453 proteins nominal in at least one column; rows 376–450



Rows: proteins at nominal $p < 0.05$ against at least one phenotype, most columns first, then by best p . Filled dot: nominal, fill = Spearman rho, size = $-\log_{10} p$. Black ring: BH < 0.05. Grey speck: tested, not nominal. Table: c_data/03_phenotype.


Protein association hits, training change (T2 – T1) (7/7)
Spearman against each phenotype; 453 proteins nominal in at least one column; rows 451–453



Rows: proteins at nominal $p < 0.05$ against at least one phenotype, most columns first, then by best p . Filled dot: nominal, fill = Spearman ρ , size = $-\log_{10} p$. Black ring: BH < 0.05 . Grey speck: tested, not nominal. Table: c_data/03_phenotype.

Protein association hits, level at T1 (1/7)


Spearman against each phenotype; 519 proteins nominal in at least one column; rows 1–75



Rows: proteins at nominal $p < 0.05$ against at least one phenotype, most columns first, then by best p . Filled dot: nominal, fill = Spearman ρ , size = $-\log_{10} p$. Black ring: BH < 0.05. Grey speck: tested, not nominal. Table: c_data/03_phenotype.x

Protein association hits, level at T1 (2/7)


Spearman against each phenotype; 519 proteins nominal in at least one column; rows 76-150



Rows: proteins at nominal $p < 0.05$ against at least one phenotype, most columns first, then by best p . Filled dot: nominal, fill = Spearman ρ , size = $-\log_{10} p$. Black ring: BH < 0.05. Grey speck: tested, not nominal. Table: c_data/03_phenotype.

Protein association hits, level at T1 (3/7)


Spearman against each phenotype; 519 proteins nominal in at least one column; rows 151–225



Rows: proteins at nominal $p < 0.05$ against at least one phenotype, most columns first, then by best p. Filled dot: nominal, fill = Spearman rho, size = $-\log_{10} p$. Black ring: BH < 0.05 . Grey speck: tested, not nominal. Table: c_data/03_phenotype.

Protein association hits, level at T1 (4/7)


Spearman against each phenotype; 519 proteins nominal in at least one column; rows 226–300



Rows: proteins at nominal $p < 0.05$ against at least one phenotype, most columns first, then by best p . Filled dot: nominal, fill = Spearman ρ , size = $-\log_{10} p$. Black ring: BH < 0.05 . Grey speck: tested, not nominal. Table: c_data/03_phenotype.


Protein association hits, level at T1 (5/7)


Spearman against each phenotype; 519 proteins nominal in at least one column; rows 301–375



Rows: proteins at nominal $p < 0.05$ against at least one phenotype, most columns first, then by best p . Filled dot: nominal, fill = Spearman ρ , size = $-\log_{10} p$. Black ring: BH < 0.05 . Grey speck: tested, not nominal. Table: c_data/03_phenotype.x


Spearman association hits, level at T1 (6/7)
Spearman against each phenotype; 519 proteins nominal in at least one column; rows 376–450





Protein association hits, acute change (T3 – T2) (1/6)


Spearman against each phenotype; 435 proteins nominal in at least one column; rows 1–75



Rows: proteins at nominal p < 0.05 against at least one phenotype, most columns first, then by best p. Filled dot: nominal, fill = Spearman rho, size = -log10 p. Black ring: BH < 0.05. Grey speck: tested, not nominal. Table: c_data/03_phenotyp

Protein association hits, acute change (T3 – T2) (2/6)


Spearman against each phenotype; 435 proteins nominal in at least one column; rows 76–150



Rows: proteins at nominal $p < 0.05$ against at least one phenotype, most columns first, then by best p . Filled dot: nominal, fill = Spearman ρ , size = $-\log_{10} p$. Black ring: BH < 0.05 . Grey speck: tested, not nominal. Table: c_data/03_phenotype.


Protein association hits, acute change (T3 – T2) (3/6)

Spearman against each phenotype; 435 proteins nominal in at least one column; rows 151–225



Protein association hits, acute change (T3 – T2) (4/6)


Spearman against each phenotype; 435 proteins nominal in at least one column; rows 226–300



Rows: proteins at nominal $p < 0.05$ against at least one phenotype, most columns first, then by best p . Filled dot: nominal, fill = Spearman ρ , size = $-\log_{10} p$. Black ring: BH < 0.05. Grey speck: tested, not nominal. Table: c_data/03_phenotype

Protein association hits, acute change (T3 – T2) (5/6)

Spearman against each phenotype; 435 proteins nominal in at least one column; rows 301–375



Spearman against each phenotype; 435 proteins nominal in at least one column; rows 376–435

