
Proteins that respond to training or the bout
Π < 0.05 in a group's training or acute contrast
top 10 per block by Π (shared: the weaker group's)
(n/48): biopsies where the protein was observed
Training
Acute
Between groups
Interaction
Shared
HR only
LR only
MPO (15/48)
CTSG (15/48)
LCN2 (8/48)
HNRNPC (35/48)
CORO1A (8/48)
CHDH (9/48)
RRAD (42/48)
COPB1 (43/48)
CNN1 (17/48)
HNRNPA2B1 (48/48)
NNMT (16/48)
EZR (31/48)
COMP (9/48)
AMPD3 (48/48)
TPPP3 (27/48)
PLIN1 (9/48)
GFPT1 (15/48)
GHITM (27/48)
TMED9 (16/48)
MAP1LC3B2 (45/48)
GCN1 (20/48)
MNDA (8/48)
LAMTOR2 (25/48)
MRPL49 (30/48)
UBE2L3 (48/48)
UBE2D1 (34/48)
OXSM (42/48)
NAMPT (47/48)
PYURF (44/48)
PPM1A (39/48)
HR
LR
HR
LR
T1
T2
T3
Training
Acute
logFC (capped at ±1.5)
1.5
1.0
0.5
0.0
-0.5
-1.0
-1.5
-log10 p
1
2
4+
Π < 0.05
Π ≥ 0.05
FDR < 0.05