

a Filtering
2400 to 1900 proteins
raw input 2400
duplicate accession 2400
remove: complement 2388
remove: erythrocyte 2369
remove: globin 2362
remove: immunoglobulin 2321
remove: keratin 2303
remove: leukocyte 2298
remove: plasma 2178
outlier samples 2178
detection 1900
0 500 1000 1500 2000 2500
proteins remaining
b Detection per run
48 runs; 3 excluded as outliers
HR LR
% of proteins missing
run
T1 T2 T3 excluded kept
c Intensity per sample
median, 25-75% and 5-95% of proteins
before normalisation
after
log2 intensity
sample
T1 T2 T3
d Dynamic range
1900 proteins over 19.1 log2 units
MB
CKM ACTA1
MYH7
MYH2
ATP5F1A
NDUFA4
TTN
DES
PSMA1
mean log2 abundance
rank by mean abundance
e Missing because faint
12.3% missing; line: limpa detection curve, slope 0.67
values missing
mean log2 abundance where observed
f Sample correlation
median r: same person 0.17, different people -0.03
r
-0.5 0.0 0.5
g Blood in each biopsy
LR - HR at T3 0.84, permutation p = 0.23
blood index (log2, vs sample median)
T1 T2 T3
T1 T2 T3 HR LR
h Blood adjustment
nine contrasts pooled, r = 0.895
log2 FC with blood covariate
log2 FC without blood covariate
Acute HR - LR Interaction Training
i p-value calibration
dashed: flat null; median |t| is 0.67 under the null
Training HR median |t| 0.76
Training LR median |t| 0.67
Acute HR median |t| 0.88
Acute LR median |t| 0.61
HR - LR T1 median |t| 0.61
HR - LR T2 median |t| 0.57
HR - LR T3 median |t| 0.74
Int.training median |t| 0.58
Int.acute median |t| 0.69
proteins
p
j Power
99% of effects below the median detectable 0.95
proteins
|log2 FC|, training interaction
detectable at 80% power
median protein
90th percentile