
Discrimination overview: proteins



All 1,900 proteins, one row each. Rows are ordered by average−linkage clustering (1 − Pearson r) of limpa's matrix, for display only. Grey: not tested. Table: c_data/protein_classification.xlsx, auc.

C1 vs C2 at T1


370 of 1900 proteins at p < 0.05, 64 at BH < 0.05. Page 1 of 2.



C1 (up to 9) against C2 (up to 6); AUC above 0.5 means higher in C1. Header: AUC with its stratified bootstrap 95% interval (2000 resamples), Wilcoxon rank−sum p (smallest attainable 0.0004), BH q within task, and the fe

C1 vs C2 at T1


370 of 1900 proteins at p < 0.05, 64 at BH < 0.05. Page 2 of 2.



C1 (up to 9) against C2 (up to 6); AUC above 0.5 means higher in C1. Header: AUC with its stratified bootstrap 95% interval (2000 resamples), Wilcoxon rank−sum p (smallest attainable 0.0004), BH q within task, and the fe

C1 vs C2 at T2


127 of 1900 proteins at p < 0.05, 0 at BH < 0.05. Page 1 of 2.



C1 (up to 8) against C2 (up to 6); AUC above 0.5 means higher in C1. Header: AUC with its stratified bootstrap 95% interval (2000 resamples), Wilcoxon rank−sum p (smallest attainable 0.00067), BH q within task, and the f

C1 vs C2 at T2


127 of 1900 proteins at p < 0.05, 0 at BH < 0.05. Page 2 of 2.



C1 (up to 8) against C2 (up to 6); AUC above 0.5 means higher in C1. Header: AUC with its stratified bootstrap 95% interval (2000 resamples), Wilcoxon rank−sum p (smallest attainable 0.00067), BH q within task, and the f

C1 vs C2 at T3


164 of 1900 proteins at p < 0.05, 0 at BH < 0.05. Page 1 of 2.



C1 (up to 8) against C2 (up to 6); AUC above 0.5 means higher in C1. Header: AUC with its stratified bootstrap 95% interval (2000 resamples), Wilcoxon rank−sum p (smallest attainable 0.00067), BH q within task, and the f

C1 vs C2 at T3


164 of 1900 proteins at p < 0.05, 0 at BH < 0.05. Page 2 of 2.



C1 (up to 8) against C2 (up to 6); AUC above 0.5 means higher in C1. Header: AUC with its stratified bootstrap 95% interval (2000 resamples), Wilcoxon rank−sum p (smallest attainable 0.00067), BH q within task, and the f

C1 vs C2, training change


148 of 1900 proteins at p < 0.05, 0 at BH < 0.05. Page 1 of 2.



C1 (up to 8) against C2 (up to 6); AUC above 0.5 means higher in C1. Header: AUC with its stratified bootstrap 95% interval (2000 resamples), Wilcoxon rank−sum p (smallest attainable 0.00067), BH q within task, and the fo

C1 vs C2, training change


148 of 1900 proteins at p < 0.05, 0 at BH < 0.05. Page 2 of 2.



C1 (up to 8) against C2 (up to 6); AUC above 0.5 means higher in C1. Header: AUC with its stratified bootstrap 95% interval (2000 resamples), Wilcoxon rank−sum p (smallest attainable 0.00067), BH q within task, and the f

C1 vs C2, acute change


79 of 1900 proteins at p < 0.05, 0 at BH < 0.05. Page 1 of 2.



C1 (up to 8) against C2 (up to 6); AUC above 0.5 means higher in C1. Header: AUC with its stratified bootstrap 95% interval (2000 resamples), Wilcoxon rank−sum p (smallest attainable 0.00067), BH q within task, and the f

C1 vs C2, acute change

79 of 1900 proteins at p < 0.05, 0 at BH < 0.05. Page 2 of 2.



C1 (up to 8) against C2 (up to 6); AUC above 0.5 means higher in C1. Header: AUC with its stratified bootstrap 95% interval (2000 resamples), Wilcoxon rank−sum p (smallest attainable 0.00067), BH q within task, and the fo