
# Discrimination overview: proteins




All 1,900 proteins, one row each. Rows are ordered by average−linkage clustering (1 − Pearson r) of limpa's matrix, for display only. Grey: not tested. Table: c_data/protein_classification.xlsx, auc.

## C1 vs C2 at T1


37 of 1900 proteins at p < 0.05, 0 at BH < 0.05. Page 1 of 2.




C1 (up to 4) against C2 (up to 10); AUC above 0.5 means higher in C1. Header: AUC with its stratified bootstrap 95% interval (2000 resamples), Wilcoxon rank−sum p (smallest attainable 0.002), BH q within task, and the fe

## C1 vs C2 at T1


37 of 1900 proteins at p < 0.05, 0 at BH < 0.05. Page 2 of 2.




C1 (up to 4) against C2 (up to 10); AUC above 0.5 means higher in C1. Header: AUC with its stratified bootstrap 95% interval (2000 resamples), Wilcoxon rank−sum p (smallest attainable 0.002), BH q within task, and the fe

## C1 vs C2 at T2


108 of 1900 proteins at p < 0.05, 0 at BH < 0.05. Page 1 of 2.




C1 (up to 4) against C2 (up to 11); AUC above 0.5 means higher in C1. Header: AUC with its stratified bootstrap 95% interval (2000 resamples), Wilcoxon rank−sum p (smallest attainable 0.0015), BH q within task, and the f

## C1 vs C2 at T2


108 of 1900 proteins at p < 0.05, 0 at BH < 0.05. Page 2 of 2.




C1 (up to 4) against C2 (up to 11); AUC above 0.5 means higher in C1. Header: AUC with its stratified bootstrap 95% interval (2000 resamples), Wilcoxon rank−sum p (smallest attainable 0.0015), BH q within task, and the f

## C1 vs C2 at T3


171 of 1900 proteins at p < 0.05, 0 at BH < 0.05. Page 1 of 2.




C1 (up to 4) against C2 (up to 11); AUC above 0.5 means higher in C1. Header: AUC with its stratified bootstrap 95% interval (2000 resamples), Wilcoxon rank−sum p (smallest attainable 0.0015), BH q within task, and the fe

## C1 vs C2 at T3


171 of 1900 proteins at p < 0.05, 0 at BH < 0.05. Page 2 of 2.




C1 (up to 4) against C2 (up to 11); AUC above 0.5 means higher in C1. Header: AUC with its stratified bootstrap 95% interval (2000 resamples), Wilcoxon rank−sum p (smallest attainable 0.0015), BH q within task, and the fe

## C1 vs C2, training change


66 of 1900 proteins at p < 0.05, 0 at BH < 0.05. Page 1 of 2.




C1 (up to 4) against C2 (up to 10); AUC above 0.5 means higher in C1. Header: AUC with its stratified bootstrap 95% interval (2000 resamples), Wilcoxon rank−sum p (smallest attainable 0.002), BH q within task, and the fe

## C1 vs C2, training change


66 of 1900 proteins at p < 0.05, 0 at BH < 0.05. Page 2 of 2.




C1 (up to 4) against C2 (up to 10); AUC above 0.5 means higher in C1. Header: AUC with its stratified bootstrap 95% interval (2000 resamples), Wilcoxon rank−sum p (smallest attainable 0.002), BH q within task, and the fe

## C1 vs C2, acute change


322 of 1900 proteins at p < 0.05, 0 at BH < 0.05. Page 1 of 2.




C1 (up to 4) against C2 (up to 11); AUC above 0.5 means higher in C1. Header: AUC with its stratified bootstrap 95% interval (2000 resamples), Wilcoxon rank−sum p (smallest attainable 0.0015), BH q within task, and the fe

## C1 vs C2, acute change

322 of 1900 proteins at p < 0.05, 0 at BH < 0.05. Page 2 of 2.




C1 (up to 4) against C2 (up to 11); AUC above 0.5 means higher in C1. Header: AUC with its stratified bootstrap 95% interval (2000 resamples), Wilcoxon rank−sum p (smallest attainable 0.0015), BH q within task, and the fe