

























Samples included (N = 45): HR_T1 (n = 7), HR_T2 (n = 7), HR_T3 (n = 7), LR_T1 (n = 8), LR_T2 (n = 8), LR_T3 (n = 8)



MD plots (points + trend) − GLOBAL
log2
median
mean
vsn
quantile
cycloess
rlr
gi
log2FC
mean intensity (log2)
2
1
0
−1
−2
15
20
25
30


MD trend−only (loess fits) − GLOBAL
log2
median
mean
vsn
quantile
cycloess
rlr
gi
log2FC
mean intensity (log2)
0.3
0.2
0.1
0.0
−0.1
15
20
25
30


MD plots (2D−binned density + trend) − GLOBAL
log2
median
mean
vsn
quantile
cycloess
rlr
gi
log2FC
mean intensity (log2)
count
40
30
20
10
2
1
0
−1
−2
15
20
25
30


Delta−trend vs raw log2 (over−/under−correction) − GLOBAL
Delta loess(M) vs log2
0.3
0.2
0.1
0.0
−0.1
15
20
25
30
mean intensity (log2)
method
cycloess
gi
log2
mean
median
quantile
rlr
vsn


MD residual distributions by method − GLOBAL
cycloess
gi
log2
mean
median
quantile
rlr
vsn
Density
loess residual (M − M_hat)


log2FC distributions by normalization − GLOBAL
Normalization method
gi
rlr
cycloess
quantile
vsn
mean
median
log2
−3
−2
−1
0
1
2
3
log2FC