


Acute_HR


all collections, fgsea, BH within contrast



Ten strongest of 83 collapse survivors, ranked by adjusted p. Size is gene count, colour is $-\log_{10}$ FDR. Table: c_data/set_tests.csv.

Acute_Interaction


all collections, fgsea, BH within contrast



Ten strongest of 44 collapse survivors, ranked by adjusted p. Size is gene count, colour is $-\log_{10}$ FDR. Table: c_data/set_tests.csv.

Acute_LR


all collections, fgsea, BH within contrast



Ten strongest of 18 collapse survivors, ranked by adjusted p. Size is gene count, colour is $-\log_{10}$ FDR. Table: c_data/set_tests.csv.

Training_HR


all collections, fgsea, BH within contrast



Ten strongest of 23 collapse survivors, ranked by adjusted p. Size is gene count, colour is $-\log_{10}$ FDR. Table: c_data/set_tests.csv.

Training_Interaction


all collections, fgsea, BH within contrast



Ten strongest of 13 collapse survivors, ranked by adjusted p. Size is gene count, colour is $-\log_{10}$ FDR. Table: c_data/set_tests.csv.

Training_LR

all collections, fgsea, BH within contrast



Ten strongest of 18 collapse survivors, ranked by adjusted p. Size is gene count, colour is $-\log_{10}$ FDR. Table: c_data/set_tests.csv.

Significant sets before and after collapse


fgsea at FDR 0.05, then collapsePathways



collapsePathways re-tests each significant set conditioned on a stronger set's leading edge and keeps it only if it stays significant. Table: c_data/set_tests.csv.

Acute_HR


GO_Slim, fgsea, BH within contrast



All of 10 collapse survivors, ranked by adjusted p. Size is gene count, colour is $-\log_{10}$ FDR. Table: c_data/set_tests.csv.

Acute_Interaction

GO_Slim, fgsea, BH within contrast




All of 6 collapse survivors, ranked by adjusted p. Size is gene count, colour is $-\log_{10}$ FDR. Table: c_data/set_tests.csv.

Training_HR

GO_Slim, fgsea, BH within contrast

$-\log_{10}$ FDR



genes



Cell
Differentiation

-1.5

-1.0

-0.5

0.0

normalised enrichment score

Cytoskel. Org.

All of 2 collapse survivors, ranked by adjusted p. Size is gene count, colour is $-\log_{10}$ FDR. Table: c_data/set_tests.csv.

Training_Interaction

GO_Slim, fgsea, BH within contrast

Transmembrane
Transport

Cytoskel. Org.

-1

0

1

normalised enrichment score

genes

● 243

● 244

● 245

● 246

● 247




2.790272

All of 2 collapse survivors, ranked by adjusted p. Size is gene count, colour is $-\log_{10}$ FDR. Table: c_data/set_tests.csv.

Acute_HR


GOBP, fgsea, BH within contrast



Ten strongest of 42 collapse survivors, ranked by adjusted p. Size is gene count, colour is $-\log_{10}$ FDR. Table: c_data/set_tests.csv.

Acute_Interaction


GOBP, fgsea, BH within contrast



Ten strongest of 20 collapse survivors, ranked by adjusted p. Size is gene count, colour is $-\log_{10}$ FDR. Table: c_data/set_tests.csv.

Acute_LR


GOBP, fgsea, BH within contrast



All of 7 collapse survivors, ranked by adjusted p. Size is gene count, colour is $-\log_{10}$ FDR. Table: c_data/set_tests.csv.

Training_HR


GOBP, fgsea, BH within contrast



Ten strongest of 11 collapse survivors, ranked by adjusted p. Size is gene count, colour is $-\log_{10}$ FDR. Table: c_data/set_tests.csv.

Training_Interaction


GOBP, fgsea, BH within contrast



All of 6 collapse survivors, ranked by adjusted p. Size is gene count, colour is $-\log_{10}$ FDR. Table: c_data/set_tests.csv.

Training_LR


GOBP, fgsea, BH within contrast



All of 9 collapse survivors, ranked by adjusted p. Size is gene count, colour is $-\log_{10} \text{FDR}$. Table: c_data/set_tests.csv.

Acute_HR

Hallmark, fgsea, BH within contrast



All of 8 collapse survivors, ranked by adjusted p. Size is gene count, colour is $-\log_{10}$ FDR. Table: c_data/set_tests.csv.

Acute_Interaction


Hallmark, fgsea, BH within contrast



All of 8 collapse survivors, ranked by adjusted p. Size is gene count, colour is $-\log_{10}$ FDR. Table: c_data/set_tests.csv.

Acute_LR

Hallmark, fgsea, BH within contrast



All of 3 collapse survivors, ranked by adjusted p. Size is gene count, colour is $-\log_{10}$ FDR. Table: c_data/set_tests.csv.

Training_HR

Hallmark, fgsea, BH within contrast

OXPPOS

FA Metabolism

0


1

2


3

normalised enrichment score

genes




$-\log_{10}$ FDR



All of 2 collapse survivors, ranked by adjusted p. Size is gene count, colour is $-\log_{10}$ FDR. Table: c_data/set_tests.csv.

Training_Interaction

Hallmark, fgsea, BH within contrast



All of 1 collapse survivor, ranked by adjusted p. Size is gene count, colour is $-\log_{10}$ FDR. Table: c_data/set_tests.csv.

Training_LR

Hallmark, fgsea, BH within contrast

OXPPOS

$-\log_{10}$ FDR

2.897684

genes

153

0.0

0.5

1.0


1.5

normalised enrichment score

All of 1 collapse survivor, ranked by adjusted p. Size is gene count, colour is $-\log_{10}$ FDR. Table: c_data/set_tests.csv.


Set contrast hits, fry (1/6)

fry p per contrast, filled by fgsea NES; 445 sets nominal in at least one contrast




Set contrast hits, fry (2/6)

fry p per contrast, filled by fgsea NES; 445 sets nominal in at least one contrast



Set contrast hits, fry (3/6)


fry p per contrast, filled by fgsea NES; 445 sets nominal in at least one contrast



Rows: sets at fry p < 0.05 in at least one contrast, most contrasts first, then by best p. Filled dot: nominal, fill = fgsea NES, size = -log10 fry p. Black ring: fry FDR < 0.05. Grey speck: not nominal

Set contrast hits, fry (4/6)


fry p per contrast, filled by fgsea NES; 445 sets nominal in at least one contrast



Rows: sets at fry p < 0.05 in at least one contrast, most contrasts first, then by best p. Filled dot: nominal, fill = fgsea NES, size = $-\log_{10} p$. Black ring: fry FDR < 0.05. Grey speck: not nominal. Tab


Set contrast hits, fry (5/6)

fry p per contrast, filled by fgsea NES; 445 sets nominal in at least one contrast

Rows: sets at fry p < 0.05 in at least one contrast, most contrasts first, then by best p. Filled dot: nominal, fill = fgsea NES, size = $-\log_{10}$ fry p. Black ring: fry FDR < 0.05. Grey

Set contrast hits, fry (6/6)

fry p per contrast, filled by fgsea NES; 445 sets nominal in at least one contrast



Rows: sets at fry p < 0.05 in at least one contrast, most contrasts first, then by best p. Filled dot: nominal, fill = fgsea NES, size = -log₁₀ fry p. Black ring: fry FDR < 0.05. Grey speck: not nominal

Acute_HR

KEGG_Legacy, fgsea, BH within contrast

Fructose
& Mannose
Metabolism

Citrate Cycle
TCA Cycle

-2.5

-2.0


-1.5

-1.0

-0.5

0.0

normalised enrichment score



genes

16

18


20

22

24


All of 2 collapse survivors, ranked by adjusted p. Size is gene count, colour is $-\log_{10}$ FDR. Table: c_data/set_tests.csv.

KEGG_Legacy, fgsea, BH within contrast



Acute_LR

KEGG_Legacy, fgsea, BH within contrast



All of 1 collapse survivor, ranked by adjusted p. Size is gene count, colour is $-\log_{10}$ FDR. Table: c_data/set_tests.csv.

Training_HR

KEGG_Legacy, fgsea, BH within contrast

genes

● 88

$-\log_{10}$ FDR

7.083969

Huntingtons
Disease

0.0

0.5

1.0

1.5

2.0


2.5

normalised enrichment score

All of 1 collapse survivor, ranked by adjusted p. Size is gene count, colour is $-\log_{10}$ FDR. Table: c_data/set_tests.csv.

Training_Interaction


KEGG_Legacy, fgsea, BH within contrast



All of 2 collapse survivors, ranked by adjusted p. Size is gene count, colour is $-\log_{10}$ FDR. Table: c_data/set_tests.csv.

Training_LR


KEGG_Legacy, fgsea, BH within contrast



All of 1 collapse survivor, ranked by adjusted p. Size is gene count, colour is $-\log_{10}$ FDR. Table: c_data/set_tests.csv.

Acute_HR


Reactome, fgsea, BH within contrast



Ten strongest of 21 collapse survivors, ranked by adjusted p. Size is gene count, colour is $-\log_{10}$ FDR. Table: c_data/set_tests.csv.

Acute_Interaction


Reactome, fgsea, BH within contrast



All of 9 collapse survivors, ranked by adjusted p. Size is gene count, colour is $-\log_{10}$ FDR. Table: c_data/set_tests.csv.

Acute_LR


Reactome, fgsea, BH within contrast



All of 7 collapse survivors, ranked by adjusted p. Size is gene count, colour is $-\log_{10}$ FDR. Table: c_data/set_tests.csv.

Training_HR

Reactome, fgsea, BH within contrast



All of 7 collapse survivors, ranked by adjusted p. Size is gene count, colour is $-\log_{10}$ FDR. Table: c_data/set_tests.csv.

Training_Interaction

Reactome, fgsea, BH within contrast

Aerobic Resp. +
ETC

Striated Muscle
Contraction

-2

-1

0

1

2

normalised enrichment score

All of 2 collapse survivors, ranked by adjusted p. Size is gene count, colour is $-\log_{10}$ FDR. Table: c_data/set_tests.csv.

$-\log_{10}$ FDR

3.0

2.8

2.6

2.4

genes

30

60


90

120

150

Training_LR

Reactome, fgsea, BH within contrast



All of 7 collapse survivors, ranked by adjusted p. Size is gene count, colour is $-\log_{10}$ FDR. Table: c_data/set_tests.csv.