



Set score against phenotype, level at T1

Spearman, subjects pooled across arms




All 576 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, level at T1

Spearman, subjects pooled across arms



HR LR



All 576 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, level at T1



Spearman, subjects pooled across arms



All 576 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, level at T1



Spearman, subjects pooled across arms



All 576 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, level at T1



Spearman, subjects pooled across arms



All 576 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, level at T1



Spearman, subjects pooled across arms



All 576 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, level at T1



Spearman, subjects pooled across arms



All 576 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, level at T1



Spearman, subjects pooled across arms



All 576 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, level at T1

Spearman, subjects pooled across arms




All 576 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, level at T1

Spearman, subjects pooled across arms


HR LR



All 576 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, level at T1

Spearman, subjects pooled across arms




All 576 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, level at T1

Spearman, subjects pooled across arms

HR LR




All 576 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, level at T1

Spearman, subjects pooled across arms


HR LR



All 576 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, level at T1

Spearman, subjects pooled across arms



GOBP: Positive Reg.

Transmembrane

Transport vs d_nfbre_l

r = -0.55 p 0.033 q 0.99

GOBP: Reg. Kinase
Activity vs comp_hypertrophy

r = +0.55 p 0.034 q 0.99

GOBP: Synaptic

Vesicle

Recycling vs d_mcsa

r = +0.55 p 0.034 q 0.99

GOBP: Nuclear

Transcribed

mRNA Catabolism vs volume_load

r = +0.55 p 0.034 q 0.84

GOBP: Resp. to

Oxidative

Stress vs d_fcsc_l

r = -0.55 p 0.034 q 1

GOBP: Reg. Cardiac

Muscle

Contraction

By Reg. The

Release Of

Sequestered

Calcium Ion vs volume_load

r = -0.55 p 0.034 q 0.84

GOBP: Reg. Cell

Substrate

Adhesion vs d_1rm_ext

r = +0.57 p 0.034 q 0.45

GOBP: Vesicle

Localization vs d_1rm_ext

r = +0.57 p 0.034 q 0.45

GOBP: Reg. Monoatomic

Ion Transport vs d_mcsa

r = -0.55 p 0.035 q 0.99

GOBP: Golgi Org. vs d_1rm_ext

r = -0.56 p 0.036 q 0.45

GOBP: Mitophagy vs d_1rm_ext

r = -0.56 p 0.036 q 0.45

GOBP: Positive Reg.

WNT Signaling vs d_1rm_ext

r = +0.56 p 0.036 q 0.45

GOBP: Reg.

Cytoplasmic

Translation vs d_1rm_legpress

r = +0.54 p 0.036 q 0.53

GOBP: Cellular Resp.

to Chemical

Stress vs d_fcsc_l

r = -0.54 p 0.037 q 1

GOBP: Reg. Actomyosin

Structure Org. vs d_fcsc_l

r = +0.54 p 0.037 q 1

GOBP: Reg. Cell

Substrate

Junction Org. vs d_fcsc_mixed

r = +0.54 p 0.037 q 1

GOBP: Positive Reg.

Transmembrane

Transport vs d_nfbre_l

r = -0.55 p 0.033 q 0.99

GOBP: Reg. Kinase
Activity vs comp_hypertrophy

r = +0.55 p 0.034 q 0.99

GOBP: Synaptic

Vesicle

Recycling vs d_mcsa


r = +0.55 p 0.034 q 0.99

GOBP: Nuclear

Transcribed

mRNA Catabolism vs volume_load


r = +0.55 p 0.034 q 0.84



All 576 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, level at T1

Spearman, subjects pooled across arms




All 576 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, level at T1

Spearman, subjects pooled across arms

HR LR




All 576 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, level at T1

Spearman, subjects pooled across arms



HR LR



All 576 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, level at T1

Spearman, subjects pooled across arms




All 576 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, level at T1

Spearman, subjects pooled across arms

HR LR




All 576 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, level at T1

Spearman, subjects pooled across arms


HR LR



All 576 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, level at T1

Spearman, subjects pooled across arms



All 576 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, level at T1

Spearman, subjects pooled across arms

HR LR



All 576 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, level at T1

Spearman, subjects pooled across arms

HR LR




All 576 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, level at T1

Spearman, subjects pooled across arms



HR LR



All 576 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, level at T1


Spearman, subjects pooled across arms



All 576 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, level at T1



Spearman, subjects pooled across arms



All 576 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, level at T1

Spearman, subjects pooled across arms



All 576 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

HR



LR



All 576 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, level at T1



Spearman, subjects pooled across arms



All 576 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, level at T1


Spearman, subjects pooled across arms



All 576 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, level at T1


Spearman, subjects pooled across arms



All 576 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, level at T1



Spearman, subjects pooled across arms



All 576 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, level at T1

Spearman, subjects pooled across arms




All 576 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, level at T1

Spearman, subjects pooled across arms

HR LR




All 576 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, level at T1

Spearman, subjects pooled across arms



HR LR



All 576 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, level at T1

Spearman, subjects pooled across arms



All 576 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.