


Training, HR (T1 to T2)


singscore per set, 6 paired subjects, AUC with direction fixed



All 50 sets reaching nominal p, by collection then p. Red separates higher in T2, blue lower. Shading is area under the curve, dashed line chance. p from the Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Training, HR (T1 to T2)


singscore per set, 6 paired subjects, AUC with direction fixed



All 50 sets reaching nominal p, by collection then p. Red separates higher in T2, blue lower. Shading is area under the curve, dashed line chance. p from the Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Training, HR (T1 to T2)


singscore per set, 6 paired subjects, AUC with direction fixed



All 50 sets reaching nominal p, by collection then p. Red separates higher in T2, blue lower. Shading is area under the curve, dashed line chance. p from the Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Training, HR (T1 to T2)


singscore per set, 6 paired subjects, AUC with direction fixed



All 50 sets reaching nominal p, by collection then p. Red separates higher in T2, blue lower. Shading is area under the curve, dashed line chance. p from the Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Training, LR (T1 to T2)


singscore per set, 8 paired subjects, AUC with direction fixed



All 30 sets reaching nominal p, by collection then p. Red separates higher in T2, blue lower. Shading is area under the curve, dashed line chance. p from the Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Training, LR (T1 to T2)


singscore per set, 8 paired subjects, AUC with direction fixed



All 30 sets reaching nominal p, by collection then p. Red separates higher in T2, blue lower. Shading is area under the curve, dashed line chance. p from the Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Acute bout, HR (T2 to T3)


singscore per set, 7 paired subjects, AUC with direction fixed



All 113 sets reaching nominal p, by collection then p. Red separates higher in T3, blue lower. Shading is area under the curve, dashed line chance. p from the Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Acute bout, HR (T2 to T3)


singscore per set, 7 paired subjects, AUC with direction fixed



All 113 sets reaching nominal p, by collection then p. Red separates higher in T3, blue lower. Shading is area under the curve, dashed line chance. p from the Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Acute bout, HR (T2 to T3)


singscore per set, 7 paired subjects, AUC with direction fixed



All 113 sets reaching nominal p, by collection then p. Red separates higher in T3, blue lower. Shading is area under the curve, dashed line chance. p from the Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Acute bout, HR (T2 to T3)


singscore per set, 7 paired subjects, AUC with direction fixed



All 113 sets reaching nominal p, by collection then p. Red separates higher in T3, blue lower. Shading is area under the curve, dashed line chance. p from the Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Acute bout, HR (T2 to T3)


singscore per set, 7 paired subjects, AUC with direction fixed



All 113 sets reaching nominal p, by collection then p. Red separates higher in T3, blue lower. Shading is area under the curve, dashed line chance. p from the Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Acute bout, HR (T2 to T3)


singscore per set, 7 paired subjects, AUC with direction fixed



All 113 sets reaching nominal p, by collection then p. Red separates higher in T3, blue lower. Shading is area under the curve, dashed line chance. p from the Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Acute bout, HR (T2 to T3)


singscore per set, 7 paired subjects, AUC with direction fixed



All 113 sets reaching nominal p, by collection then p. Red separates higher in T3, blue lower. Shading is area under the curve, dashed line chance. p from the Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Acute bout, HR (T2 to T3)


singscore per set, 7 paired subjects, AUC with direction fixed



All 113 sets reaching nominal p, by collection then p. Red separates higher in T3, blue lower. Shading is area under the curve, dashed line chance. p from the Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Acute bout, LR (T2 to T3)


singscore per set, 8 paired subjects, AUC with direction fixed



All 153 sets reaching nominal p, by collection then p. Red separates higher in T3, blue lower. Shading is area under the curve, dashed line chance, p from the Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Acute bout, LR (T2 to T3)


singscore per set, 8 paired subjects, AUC with direction fixed



All 153 sets reaching nominal p, by collection then p. Red separates higher in T3, blue lower. Shading is area under the curve, dashed line chance, p from the Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Acute bout, LR (T2 to T3)


singscore per set, 8 paired subjects, AUC with direction fixed



All 153 sets reaching nominal p, by collection then p. Red separates higher in T3, blue lower. Shading is area under the curve, dashed line chance, p from the Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Acute bout, LR (T2 to T3)


singscore per set, 8 paired subjects, AUC with direction fixed



All 153 sets reaching nominal p, by collection then p. Red separates higher in T3, blue lower. Shading is area under the curve, dashed line chance, p from the Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Acute bout, LR (T2 to T3)


singscore per set, 8 paired subjects, AUC with direction fixed



All 153 sets reaching nominal p, by collection then p. Red separates higher in T3, blue lower. Shading is area under the curve, dashed line chance, p from the Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Acute bout, LR (T2 to T3)


singscore per set, 8 paired subjects, AUC with direction fixed



All 153 sets reaching nominal p, by collection then p. Red separates higher in T3, blue lower. Shading is area under the curve, dashed line chance, p from the Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Acute bout, LR (T2 to T3)


singscore per set, 8 paired subjects, AUC with direction fixed



All 153 sets reaching nominal p, by collection then p. Red separates higher in T3, blue lower. Shading is area under the curve, dashed line chance. p from the Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Acute bout, LR (T2 to T3)


singscore per set, 8 paired subjects, AUC with direction fixed



All 153 sets reaching nominal p, by collection then p. Red separates higher in T3, blue lower. Shading is area under the curve, dashed line chance. p from the Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Acute bout, LR (T2 to T3)


singscore per set, 8 paired subjects, AUC with direction fixed



All 153 sets reaching nominal p, by collection then p. Red separates higher in T3, blue lower. Shading is area under the curve, dashed line chance. p from the Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Acute bout, LR (T2 to T3)


singscore per set, 8 paired subjects, AUC with direction fixed



All 153 sets reaching nominal p, by collection then p. Red separates higher in T3, blue lower. Shading is area under the curve, dashed line chance. p from the Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

HR vs LR at T2


singscore per set, 7 subjects, AUC with direction fixed



All 14 sets reaching nominal p, by collection then p. Red separates higher in HR, blue lower. Shading is area under the curve, dashed line chance. p from the Wilcoxon rank-sum test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

HR vs LR, training change


singscore per set, 6 subjects, AUC with direction fixed



All 37 sets reaching nominal p, by collection then p. Red separates higher in HR, blue lower. Shading is area under the curve, dashed line chance, p from the Wilcoxon rank-sum test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

HR vs LR, training change


singscore per set, 6 subjects, AUC with direction fixed



All 37 sets reaching nominal p, by collection then p. Red separates higher in HR, blue lower. Shading is area under the curve, dashed line chance, p from the Wilcoxon rank-sum test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

HR vs LR, training change


singscore per set, 6 subjects, AUC with direction fixed



All 37 sets reaching nominal p, by collection then p. Red separates higher in HR, blue lower. Shading is area under the curve, dashed line chance. p from the Wilcoxon rank-sum test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

HR vs LR, acute change


singscore per set, 7 subjects, AUC with direction fixed



All 56 sets reaching nominal p, by collection then p. Red separates higher in HR, blue lower. Shading is area under the curve, dashed line chance, p from the Wilcoxon rank-sum test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

HR vs LR, acute change


singscore per set, 7 subjects, AUC with direction fixed



All 56 sets reaching nominal p, by collection then p. Red separates higher in HR, blue lower. Shading is area under the curve, dashed line chance, p from the Wilcoxon rank-sum test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

HR vs LR, acute change


singscore per set, 7 subjects, AUC with direction fixed



All 56 sets reaching nominal p, by collection then p. Red separates higher in HR, blue lower. Shading is area under the curve, dashed line chance, p from the Wilcoxon rank-sum test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

HR vs LR, acute change

singscore per set, 7 subjects, AUC with direction fixed



All 56 sets reaching nominal p, by collection then p. Red separates higher in HR, blue lower. Shading is area under the curve, dashed line chance. p from the Wilcoxon rank-sum test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, training change (T2 – T1)

Spearman, subjects pooled across arms

HR LR




All 518 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, training change (T2 – T1)

Spearman, subjects pooled across arms

HR LR



All 518 set–outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, training change (T2 – T1)

Spearman, subjects pooled across arms

HR LR




All 518 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, training change (T2 – T1)

Spearman, subjects pooled across arms

HR LR




All 518 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, training change (T2 – T1)

Spearman, subjects pooled across arms


HR LR



All 518 set–outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, training change (T2 – T1)



Spearman, subjects pooled across arms



All 518 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, training change (T2 – T1)



Spearman, subjects pooled across arms



All 518 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, training change (T2 – T1)



Spearman, subjects pooled across arms



All 518 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, training change (T2 – T1)

Spearman, subjects pooled across arms




All 518 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, training change (T2 – T1)

Spearman, subjects pooled across arms


HR LR



All 518 set–outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, training change (T2 – T1)

Spearman, subjects pooled across arms




All 518 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, training change (T2 – T1)

Spearman, subjects pooled across arms

HR LR




All 518 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, training change (T2 – T1)

Spearman, subjects pooled across arms

HR LR




All 518 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, training change (T2 – T1)

Spearman, subjects pooled across arms

HR LR




All 518 set–outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, training change (T2 – T1)

Spearman, subjects pooled across arms

HR LR




All 518 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, training change (T2 – T1)

Spearman, subjects pooled across arms

HR LR




All 518 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, training change (T2 – T1)

Spearman, subjects pooled across arms

HR LR




All 518 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, training change (T2 – T1)

Spearman, subjects pooled across arms

HR LR




All 518 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, training change (T2 – T1)

Spearman, subjects pooled across arms



HR LR



All 518 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, training change (T2 – T1)

Spearman, subjects pooled across arms




All 518 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, training change (T2 – T1)

Spearman, subjects pooled across arms

HR LR




All 518 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, training change (T2 – T1)

Spearman, subjects pooled across arms

HR LR



All 518 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, training change (T2 – T1)

Spearman, subjects pooled across arms

HR LR




All 518 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, training change (T2 – T1)

Spearman, subjects pooled across arms

HR LR




All 518 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, training change (T2 – T1)

Spearman, subjects pooled across arms

HR LR




All 518 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, training change (T2 – T1)

Spearman, subjects pooled across arms

HR LR




All 518 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, training change (T2 – T1)


Spearman, subjects pooled across arms

HR LR


Reactome: Sars Cov 1 Host Interactions vs d_1rm_ext
r = +0.74 p 0.004 q 0.27




Reactome: Sars Cov 1 Infection vs d_1rm_ext
r = +0.73 p 0.0048 q 0.27




Reactome: Pre-mRNA Processing vs d_1rm_legpress
r = +0.70 p 0.0053 q 0.79




Reactome: Adherens Junctions Interactions vs d_1rm_ext
r = -0.72 p 0.0059 q 0.27




Reactome: mRNA Splicing vs d_1rm_legpress
r = +0.69 p 0.0062 q 0.79




Reactome: Transmission Across Chemical Synapses vs volume_load
r = -0.68 p 0.007 q 0.77




Reactome: mTOR Signalling vs d_1rm_legpress
r = +0.68 p 0.0072 q 0.79




Reactome: Factors Involved In Megakaryocyte Development & Platelet Production vs d_1rm_ext
r = -0.70 p 0.0081 q 0.27




Reactome: Dengue Virus Host Interactions vs d_1rm_ext
r = -0.69 p 0.0085 q 0.27




Reactome: Chromatin Org. vs d_1rm_ext
r = -0.68 p 0.011 q 0.27




Reactome: Cellular Responses To Mechanical Stimuli vs d_fcsl_II
r = -0.65 p 0.011 q 0.93




Reactome: Hcmv Early Events vs volume_load
r = -0.65 p 0.011 q 0.93




Reactome: Intrinsic Pathway For Apoptosis vs d_1rm_ext
r = +0.67 p 0.012 q 0.27




Reactome: Runx1 Regulates Transcription Of Genes Involved In Differentiation Of Hscs vs d_1rm_ext
r = -0.67 p 0.012 q 0.27



Reactome: Rac3 GTPase Cycle vs d_1rm_ext
r = +0.67 p 0.013 q 0.27



Reactome: Transcriptional Regulation By Runx1 vs d_1rm_ext
r = -0.67 p 0.013 q 0.27





set score, training change (T2 – T1), one point per subject

All 518 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, training change (T2 – T1)

Spearman, subjects pooled across arms



All 518 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, training change (T2 – T1)

Spearman, subjects pooled across arms

HR LR




All 518 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, training change (T2 – T1)

Spearman, subjects pooled across arms

HR LR




All 518 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, training change (T2 – T1)

Spearman, subjects pooled across arms


HR LR



All 518 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, training change (T2 – T1)

Spearman, subjects pooled across arms




All 518 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, training change (T2 – T1)

Spearman, subjects pooled across arms



HR LR



All 518 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, level at T1

Spearman, subjects pooled across arms




All 576 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, level at T1

Spearman, subjects pooled across arms


HR LR



All 576 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, level at T1



Spearman, subjects pooled across arms



All 576 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, level at T1



Spearman, subjects pooled across arms



All 576 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, level at T1



Spearman, subjects pooled across arms



All 576 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, level at T1



Spearman, subjects pooled across arms



All 576 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, level at T1



Spearman, subjects pooled across arms



All 576 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, level at T1


Spearman, subjects pooled across arms



All 576 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, level at T1

Spearman, subjects pooled across arms




All 576 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, level at T1

Spearman, subjects pooled across arms

HR LR




All 576 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, level at T1

Spearman, subjects pooled across arms

HR LR




All 576 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, level at T1

Spearman, subjects pooled across arms

HR LR




All 576 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, level at T1

Spearman, subjects pooled across arms


HR LR



All 576 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, level at T1

Spearman, subjects pooled across arms



GOBP: Positive Reg.

Transmembrane

Transport vs d_nfbre_l

r = -0.55 p 0.033 q 0.99

GOBP: Reg. Kinase

Activity vs comp_hypertrophy

r = +0.55 p 0.034 q 0.99

GOBP: Synaptic

Vesicle

Recycling vs d_mcsa

r = +0.55 p 0.034 q 0.99

GOBP: Nuclear

Transcribed

mRNA Catabolism vs volume_load

r = +0.55 p 0.034 q 0.84

GOBP: Resp. to

Oxidative

Stress vs d_fcsc_l

r = -0.55 p 0.034 q 1

GOBP: Reg. Cardiac

Muscle

Contraction

By Reg. The

Release Of

Sequestered

Calcium Ion vs volume_load

r = -0.55 p 0.034 q 0.84

GOBP: Reg. Cell

Substrate

Adhesion vs d_1rm_ext

r = +0.57 p 0.034 q 0.45

GOBP: Vesicle

Localization vs d_1rm_ext

r = +0.57 p 0.034 q 0.45

GOBP: Reg. Monoatomic

Ion Transport vs d_mcsa

r = -0.55 p 0.035 q 0.99

GOBP: Golgi Org. vs d_1rm_ext

r = -0.56 p 0.036 q 0.45

GOBP: Mitophagy vs d_1rm_ext

r = -0.56 p 0.036 q 0.45

GOBP: Positive Reg.

WNT Signaling vs d_1rm_ext

r = +0.56 p 0.036 q 0.45

GOBP: Reg. Cytoplasmic

Translation vs d_1rm_legpress

r = +0.54 p 0.036 q 0.53

GOBP: Cellular Resp.

to Chemical

Stress vs d_fcsc_l

r = -0.54 p 0.037 q 1

GOBP: Reg. Actomyosin

Structure Org. vs d_fcsc_l

r = +0.54 p 0.037 q 1

GOBP: Reg. Cell

Substrate

Junction Org. vs d_fcsc_mixed

r = +0.54 p 0.037 q 1

GOBP: Positive Reg.

Transmembrane

Transport vs d_nfbre_l

r = -0.55 p 0.033 q 0.99

GOBP: Reg. Kinase

Activity vs comp_hypertrophy

r = +0.55 p 0.034 q 0.99

GOBP: Synaptic

Vesicle

Recycling vs d_mcsa

r = +0.55 p 0.034 q 0.99

GOBP: Nuclear

Transcribed

mRNA Catabolism vs volume_load

r = +0.55 p 0.034 q 0.84

GOBP: Resp. to

Oxidative

Stress vs d_fcsc_l

r = -0.55 p 0.034 q 1

GOBP: Reg. Cardiac

Muscle

Contraction

By Reg. The

Release Of

Sequestered

Calcium Ion vs volume_load

r = -0.55 p 0.034 q 0.84

GOBP: Reg. Cell

Substrate

Adhesion vs d_1rm_ext

r = +0.57 p 0.034 q 0.45

GOBP: Vesicle

Localization vs d_1rm_ext

r = +0.57 p 0.034 q 0.45

GOBP: Reg. Monoatomic

Ion Transport vs d_mcsa

r = -0.55 p 0.035 q 0.99

GOBP: Golgi Org. vs d_1rm_ext

r = -0.56 p 0.036 q 0.45

GOBP: Mitophagy vs d_1rm_ext

r = -0.56 p 0.036 q 0.45

GOBP: Positive Reg.

WNT Signaling vs d_1rm_ext

r = +0.56 p 0.036 q 0.45

GOBP: Reg. Cytoplasmic

Translation vs d_1rm_legpress

r = +0.54 p 0.036 q 0.53

GOBP: Cellular Resp.

to Chemical

Stress vs d_fcsc_l

r = -0.54 p 0.037 q 1

GOBP: Reg. Actomyosin

Structure Org. vs d_fcsc_l

r = +0.54 p 0.037 q 1

GOBP: Reg. Cell

Substrate

Junction Org. vs d_fcsc_mixed

r = +0.54 p 0.037 q 1

All 576 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, level at T1

Spearman, subjects pooled across arms

HR LR




All 576 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, level at T1

Spearman, subjects pooled across arms

HR LR




All 576 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, level at T1

Spearman, subjects pooled across arms

HR LR




All 576 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, level at T1

Spearman, subjects pooled across arms

HR LR




All 576 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, level at T1

Spearman, subjects pooled across arms

HR LR




All 576 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, level at T1

Spearman, subjects pooled across arms


HR LR



All 576 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, level at T1


Spearman, subjects pooled across arms



All 576 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, level at T1


Spearman, subjects pooled across arms



All 576 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, level at T1

Spearman, subjects pooled across arms




All 576 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, level at T1

Spearman, subjects pooled across arms

HR LR



All 576 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, level at T1

Spearman, subjects pooled across arms


HR LR



All 576 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, level at T1



Spearman, subjects pooled across arms



All 576 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, level at T1

Spearman, subjects pooled across arms



All 576 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

HR



LR



All 576 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, level at T1

Spearman, subjects pooled across arms




All 576 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, level at T1

Spearman, subjects pooled across arms


HR LR



All 576 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, level at T1


Spearman, subjects pooled across arms



All 576 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, level at T1

Spearman, subjects pooled across arms




All 576 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, level at T1

Spearman, subjects pooled across arms



HR LR



All 576 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, level at T1

Spearman, subjects pooled across arms




All 576 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, level at T1

Spearman, subjects pooled across arms



HR LR



All 576 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, level at T1

Spearman, subjects pooled across arms




All 576 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, acute change (T3 – T2)

Spearman, subjects pooled across arms

HR LR




All 511 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, acute change (T3 – T2)

Spearman, subjects pooled across arms

HR LR




All 511 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, acute change (T3 – T2)

Spearman, subjects pooled across arms

HR LR




All 511 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, acute change (T3 – T2)

Spearman, subjects pooled across arms



HR LR



All 511 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, acute change (T3 – T2)



Spearman, subjects pooled across arms



All 511 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, acute change (T3 – T2)



Spearman, subjects pooled across arms



All 511 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, acute change (T3 – T2)


Spearman, subjects pooled across arms



All 511 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, acute change (T3 – T2)



Spearman, subjects pooled across arms



All 511 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, acute change (T3 – T2)

Spearman, subjects pooled across arms




All 511 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, acute change (T3 – T2)

Spearman, subjects pooled across arms

HR LR




All 511 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, acute change (T3 – T2)

Spearman, subjects pooled across arms

HR LR




All 511 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, acute change (T3 – T2)

Spearman, subjects pooled across arms

HR LR




All 511 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, acute change (T3 – T2)

Spearman, subjects pooled across arms


HR LR



All 511 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, acute change (T3 – T2)

Spearman, subjects pooled across arms




All 511 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, acute change (T3 – T2)

Spearman, subjects pooled across arms



HR LR



All 511 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, acute change (T3 – T2)

Spearman, subjects pooled across arms




All 511 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, acute change (T3 – T2)

Spearman, subjects pooled across arms


HR LR



All 511 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, acute change (T3 – T2)

Spearman, subjects pooled across arms




All 511 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, acute change (T3 – T2)

Spearman, subjects pooled across arms

HR LR




All 511 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, acute change (T3 – T2)

Spearman, subjects pooled across arms

HR LR




All 511 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, acute change (T3 – T2)

Spearman, subjects pooled across arms

HR LR




All 511 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, acute change (T3 – T2)

Spearman, subjects pooled across arms


HR LR



All 511 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, acute change (T3 – T2)

Spearman, subjects pooled across arms




All 511 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, acute change (T3 – T2)

Spearman, subjects pooled across arms

HR LR




All 511 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, acute change (T3 – T2)

Spearman, subjects pooled across arms

HR LR




All 511 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, acute change (T3 – T2)

Spearman, subjects pooled across arms

HR LR




All 511 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, acute change (T3 – T2)

Spearman, subjects pooled across arms

HR LR




All 511 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, acute change (T3 – T2)

Spearman, subjects pooled across arms

HR LR




All 511 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, acute change (T3 – T2)

Spearman, subjects pooled across arms

HR LR




All 511 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, acute change (T3 – T2)

Spearman, subjects pooled across arms

HR LR




All 511 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, acute change (T3 – T2)

Spearman, subjects pooled across arms

HR LR




All 511 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Set score against phenotype, acute change (T3 – T2)

Spearman, subjects pooled across arms


HR LR



All 511 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection, window and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 44.9 of 897, GO_Slim 2.9 of 58, Hallmark 1.7 of 33, KEGG_Legacy 3.1 of 62, Reactome 16.4 of 328.

Nominal hits relative to chance, by collection

Wilcoxon per set, 5 collections, uncorrected p



Bar length is observed nominal hits divided by the count that collection returns under the null. Red clears 1, grey does not. Labels give observed of tested. Table: [c_data/05_classify_and](#)