

Sample model overview: proteins




All 1,900 proteins, one row each. Moderated t of the within and between terms, abundance ~ timepoint + between + within, subject blocked. Colours stop at the 99th percentile of |t|, 2.8.
Rows are ordered by average−linkage clustering (1 − Pearson r) of the imputed matrix, for display only. Grey: not tested. Table: c_data/04_associate.xlsx, model.

Change score overview, training window: proteins




All 1,900 proteins, one row each. Spearman rho between the log2 abundance (change, T1 to T2) and each outcome. Rows are ordered by average−linkage clustering (1 − Pearson r) of the imputed matrix, for display only. Grey: not tested. Table: c_data/04_associate.xlsx, change_score.

Change score overview, baseline window: proteins



All 1,900 proteins, one row each. Spearman rho between the log2 abundance (T1 level) and each outcome. Rows are ordered by average−linkage clustering (1 − Pearson r) of the imputed matrix, for display only. Grey: not tested. Table: c_data/04_associate.xlsx, change_score.


Change score overview, acute window: proteins



All 1,900 proteins, one row each. Spearman rho between the log2 abundance (change, T2 to T3) and each outcome. Rows are ordered by average−linkage clustering (1 − Pearson r) of the imputed matrix, for display only. Grey: not tested. Table: c_data/04_associate.xlsx, change_score.

fcsa_mixed, within−person term


61 of 1640 proteins at p < 0.05, 0 at BH < 0.05. Page 1 of 6.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

fcsa_mixed, within−person term


61 of 1640 proteins at p < 0.05, 0 at BH < 0.05. Page 2 of 6.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

fcsa_mixed, within−person term


61 of 1640 proteins at p < 0.05, 0 at BH < 0.05. Page 3 of 6.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

fcsa_mixed, within−person term


61 of 1640 proteins at p < 0.05, 0 at BH < 0.05. Page 4 of 6.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

fcsa_mixed, within−person term


61 of 1640 proteins at p < 0.05, 0 at BH < 0.05. Page 5 of 6.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

fcsa_mixed, within−person term


61 of 1640 proteins at p < 0.05, 0 at BH < 0.05. Page 6 of 6.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

fcsa_mixed, between−person term


67 of 1640 proteins at p < 0.05, 0 at BH < 0.05. Page 1 of 6.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

fcsa_mixed, between−person term

67 of 1640 proteins at p < 0.05, 0 at BH < 0.05. Page 2 of 6.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

fcsa_mixed, between−person term


67 of 1640 proteins at p < 0.05, 0 at BH < 0.05. Page 3 of 6.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

fcsa_mixed, between−person term


67 of 1640 proteins at p < 0.05, 0 at BH < 0.05. Page 4 of 6.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

fcsa_mixed, between−person term


67 of 1640 proteins at p < 0.05, 0 at BH < 0.05. Page 5 of 6.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

fcsa_mixed, between−person term


67 of 1640 proteins at p < 0.05, 0 at BH < 0.05. Page 6 of 6.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

fcsa_type1, within−person term


59 of 1640 proteins at p < 0.05, 0 at BH < 0.05. Page 1 of 5.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

fcsa_type1, within−person term


59 of 1640 proteins at p < 0.05, 0 at BH < 0.05. Page 2 of 5.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

fcsa_type1, within−person term


59 of 1640 proteins at p < 0.05, 0 at BH < 0.05. Page 3 of 5.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

fcsa_type1, within−person term


59 of 1640 proteins at p < 0.05, 0 at BH < 0.05. Page 4 of 5.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

fcsa_type1, within−person term


59 of 1640 proteins at p < 0.05, 0 at BH < 0.05. Page 5 of 5.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

fcsa_type1, between−person term


64 of 1640 proteins at p < 0.05, 0 at BH < 0.05. Page 1 of 6.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

fcsa_type1, between−person term


64 of 1640 proteins at p < 0.05, 0 at BH < 0.05. Page 2 of 6.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

fcsa_type1, between−person term


64 of 1640 proteins at p < 0.05, 0 at BH < 0.05. Page 3 of 6.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

fcsa_type1, between−person term


64 of 1640 proteins at p < 0.05, 0 at BH < 0.05. Page 4 of 6.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

fcsa_type1, between−person term


64 of 1640 proteins at p < 0.05, 0 at BH < 0.05. Page 5 of 6.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

fcsa_type1, between−person term


64 of 1640 proteins at p < 0.05, 0 at BH < 0.05. Page 6 of 6.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

fcsa_type2, within−person term


74 of 1640 proteins at p < 0.05, 0 at BH < 0.05. Page 1 of 7.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

fcsa_type2, within−person term


74 of 1640 proteins at p < 0.05, 0 at BH < 0.05. Page 2 of 7.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

fcsa_type2, within−person term


74 of 1640 proteins at p < 0.05, 0 at BH < 0.05. Page 3 of 7.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

fcsa_type2, within−person term


74 of 1640 proteins at p < 0.05, 0 at BH < 0.05. Page 4 of 7.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

fcsa_type2, within−person term


74 of 1640 proteins at p < 0.05, 0 at BH < 0.05. Page 5 of 7.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

fcsa_type2, within−person term


74 of 1640 proteins at p < 0.05, 0 at BH < 0.05. Page 6 of 7.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

fcsa_type2, within−person term


74 of 1640 proteins at p < 0.05, 0 at BH < 0.05. Page 7 of 7.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

fcsa_type2, between−person term


64 of 1640 proteins at p < 0.05, 0 at BH < 0.05. Page 1 of 6.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

fcsa_type2, between−person term

64 of 1640 proteins at p < 0.05, 0 at BH < 0.05. Page 2 of 6.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

fcsa_type2, between−person term


64 of 1640 proteins at p < 0.05, 0 at BH < 0.05. Page 3 of 6.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

fcsa_type2, between−person term


64 of 1640 proteins at p < 0.05, 0 at BH < 0.05. Page 4 of 6.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

fcsa_type2, between−person term


64 of 1640 proteins at p < 0.05, 0 at BH < 0.05. Page 5 of 6.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

fcsa_type2, between−person term


64 of 1640 proteins at p < 0.05, 0 at BH < 0.05. Page 6 of 6.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

fibres_mixed, within−person term


46 of 1640 proteins at p < 0.05, 0 at BH < 0.05. Page 1 of 4.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

fibres_mixed, within−person term


46 of 1640 proteins at p < 0.05, 0 at BH < 0.05. Page 2 of 4.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

fibres_mixed, within−person term


46 of 1640 proteins at p < 0.05, 0 at BH < 0.05. Page 3 of 4.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

fibres_mixed, within−person term


46 of 1640 proteins at p < 0.05, 0 at BH < 0.05. Page 4 of 4.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

fibres_mixed, between−person term


83 of 1640 proteins at p < 0.05, 0 at BH < 0.05. Page 1 of 7.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

fibres_mixed, between−person term


83 of 1640 proteins at p < 0.05, 0 at BH < 0.05. Page 2 of 7.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

fibres_mixed, between−person term


83 of 1640 proteins at p < 0.05, 0 at BH < 0.05. Page 3 of 7.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

fibres_mixed, between−person term


83 of 1640 proteins at p < 0.05, 0 at BH < 0.05. Page 4 of 7.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

fibres_mixed, between−person term


83 of 1640 proteins at p < 0.05, 0 at BH < 0.05. Page 5 of 7.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

fibres_mixed, between−person term


83 of 1640 proteins at p < 0.05, 0 at BH < 0.05. Page 6 of 7.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

fibres_mixed, between−person term


83 of 1640 proteins at p < 0.05, 0 at BH < 0.05. Page 7 of 7.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

fibres_type1, within−person term


52 of 1640 proteins at p < 0.05, 0 at BH < 0.05. Page 1 of 5.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

fibres_type1, within−person term


52 of 1640 proteins at p < 0.05, 0 at BH < 0.05. Page 2 of 5.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

fibres_type1, within−person term


52 of 1640 proteins at p < 0.05, 0 at BH < 0.05. Page 3 of 5.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

fibres_type1, within−person term


52 of 1640 proteins at p < 0.05, 0 at BH < 0.05. Page 4 of 5.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

fibres_type1, within−person term


52 of 1640 proteins at p < 0.05, 0 at BH < 0.05. Page 5 of 5.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

fibres_type1, between−person term


63 of 1640 proteins at p < 0.05, 0 at BH < 0.05. Page 1 of 6.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

fibres_type1, between−person term


63 of 1640 proteins at p < 0.05, 0 at BH < 0.05. Page 2 of 6.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

fibres_type1, between−person term


63 of 1640 proteins at p < 0.05, 0 at BH < 0.05. Page 3 of 6.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

fibres_type1, between−person term


63 of 1640 proteins at p < 0.05, 0 at BH < 0.05. Page 4 of 6.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

fibres_type1, between−person term


63 of 1640 proteins at p < 0.05, 0 at BH < 0.05. Page 5 of 6.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

fibres_type1, between−person term


63 of 1640 proteins at p < 0.05, 0 at BH < 0.05. Page 6 of 6.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

mcsa, within−person term


63 of 1640 proteins at p < 0.05, 0 at BH < 0.05. Page 1 of 6.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

mcsa, within−person term


63 of 1640 proteins at p < 0.05, 0 at BH < 0.05. Page 2 of 6.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

mcsa, within−person term

63 of 1640 proteins at p < 0.05, 0 at BH < 0.05. Page 3 of 6.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

mcsa, within−person term


63 of 1640 proteins at p < 0.05, 0 at BH < 0.05. Page 4 of 6.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

mcsa, within−person term

63 of 1640 proteins at p < 0.05, 0 at BH < 0.05. Page 5 of 6.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

mcsa, within−person term


63 of 1640 proteins at p < 0.05, 0 at BH < 0.05. Page 6 of 6.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

mcsa, between−person term


206 of 1640 proteins at p < 0.05, 0 at BH < 0.05. Page 1 of 18.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

mcsa, between−person term


206 of 1640 proteins at p < 0.05, 0 at BH < 0.05. Page 2 of 18.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

mcsa, between−person term


206 of 1640 proteins at p < 0.05, 0 at BH < 0.05. Page 3 of 18.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

mcsa, between−person term


206 of 1640 proteins at p < 0.05, 0 at BH < 0.05. Page 4 of 18.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

mcsa, between−person term


206 of 1640 proteins at p < 0.05, 0 at BH < 0.05. Page 5 of 18.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

mcsa, between−person term


206 of 1640 proteins at p < 0.05, 0 at BH < 0.05. Page 6 of 18.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

mcsa, between−person term


206 of 1640 proteins at p < 0.05, 0 at BH < 0.05. Page 7 of 18.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

mcsa, between−person term


206 of 1640 proteins at p < 0.05, 0 at BH < 0.05. Page 8 of 18.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

mcsa, between−person term


206 of 1640 proteins at p < 0.05, 0 at BH < 0.05. Page 9 of 18.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

mcsa, between−person term


206 of 1640 proteins at p < 0.05, 0 at BH < 0.05. Page 10 of 18.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

mcsa, between−person term


206 of 1640 proteins at p < 0.05, 0 at BH < 0.05. Page 11 of 18.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

mcsa, between−person term


206 of 1640 proteins at p < 0.05, 0 at BH < 0.05. Page 12 of 18.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

mcsa, between−person term


206 of 1640 proteins at p < 0.05, 0 at BH < 0.05. Page 13 of 18.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

mcsa, between−person term


206 of 1640 proteins at p < 0.05, 0 at BH < 0.05. Page 14 of 18.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

mcsa, between−person term


206 of 1640 proteins at p < 0.05, 0 at BH < 0.05. Page 15 of 18.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

mcsa, between−person term


206 of 1640 proteins at p < 0.05, 0 at BH < 0.05. Page 16 of 18.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

mcsa, between−person term


206 of 1640 proteins at p < 0.05, 0 at BH < 0.05. Page 17 of 18.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

mcsa, between−person term


206 of 1640 proteins at p < 0.05, 0 at BH < 0.05. Page 18 of 18.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

leg_press_1rm, within−person term


82 of 1640 proteins at p < 0.05, 0 at BH < 0.05. Page 1 of 7.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

leg_press_1rm, within−person term


82 of 1640 proteins at p < 0.05, 0 at BH < 0.05. Page 2 of 7.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

leg_press_1rm, within−person term


82 of 1640 proteins at p < 0.05, 0 at BH < 0.05. Page 3 of 7.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

leg_press_1rm, within−person term


82 of 1640 proteins at p < 0.05, 0 at BH < 0.05. Page 4 of 7.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

leg_press_1rm, within−person term


82 of 1640 proteins at p < 0.05, 0 at BH < 0.05. Page 5 of 7.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

leg_press_1rm, within−person term


82 of 1640 proteins at p < 0.05, 0 at BH < 0.05. Page 6 of 7.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

leg_press_1rm, within−person term


82 of 1640 proteins at p < 0.05, 0 at BH < 0.05. Page 7 of 7.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

leg_press_1rm, between−person term


107 of 1640 proteins at p < 0.05, 0 at BH < 0.05. Page 1 of 9.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

leg_press_1rm, between−person term


107 of 1640 proteins at p < 0.05, 0 at BH < 0.05. Page 2 of 9.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

leg_press_1rm, between−person term


107 of 1640 proteins at p < 0.05, 0 at BH < 0.05. Page 3 of 9.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

leg_press_1rm, between−person term


107 of 1640 proteins at p < 0.05, 0 at BH < 0.05. Page 4 of 9.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

leg_press_1rm, between−person term


107 of 1640 proteins at p < 0.05, 0 at BH < 0.05. Page 5 of 9.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

leg_press_1rm, between−person term


107 of 1640 proteins at p < 0.05, 0 at BH < 0.05. Page 6 of 9.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

leg_press_1rm, between−person term


107 of 1640 proteins at p < 0.05, 0 at BH < 0.05. Page 7 of 9.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

leg_press_1rm, between−person term


107 of 1640 proteins at p < 0.05, 0 at BH < 0.05. Page 8 of 9.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

leg_press_1rm, between−person term


107 of 1640 proteins at p < 0.05, 0 at BH < 0.05. Page 9 of 9.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

leg_ext_1rm, within−person term


122 of 1615 proteins at p < 0.05, 0 at BH < 0.05. Page 1 of 11.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

leg_ext_1rm, within−person term


122 of 1615 proteins at p < 0.05, 0 at BH < 0.05. Page 2 of 11.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

leg_ext_1rm, within−person term


122 of 1615 proteins at p < 0.05, 0 at BH < 0.05. Page 3 of 11.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

leg_ext_1rm, within−person term


122 of 1615 proteins at p < 0.05, 0 at BH < 0.05. Page 4 of 11.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

leg_ext_1rm, within−person term


122 of 1615 proteins at p < 0.05, 0 at BH < 0.05. Page 5 of 11.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

leg_ext_1rm, within−person term


122 of 1615 proteins at p < 0.05, 0 at BH < 0.05. Page 6 of 11.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

leg_ext_1rm, within−person term


122 of 1615 proteins at p < 0.05, 0 at BH < 0.05. Page 7 of 11.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

leg_ext_1rm, within−person term


122 of 1615 proteins at p < 0.05, 0 at BH < 0.05. Page 8 of 11.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

leg_ext_1rm, within−person term


122 of 1615 proteins at p < 0.05, 0 at BH < 0.05. Page 9 of 11.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

leg_ext_1rm, within−person term


122 of 1615 proteins at p < 0.05, 0 at BH < 0.05. Page 10 of 11.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

leg_ext_1rm, within−person term


122 of 1615 proteins at p < 0.05, 0 at BH < 0.05. Page 11 of 11.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

leg_ext_1rm, between−person term


130 of 1615 proteins at p < 0.05, 0 at BH < 0.05. Page 1 of 11.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

leg_ext_1rm, between−person term


130 of 1615 proteins at p < 0.05, 0 at BH < 0.05. Page 2 of 11.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

leg_ext_1rm, between−person term


130 of 1615 proteins at p < 0.05, 0 at BH < 0.05. Page 3 of 11.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

leg_ext_1rm, between−person term


130 of 1615 proteins at p < 0.05, 0 at BH < 0.05. Page 4 of 11.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

leg_ext_1rm, between−person term


130 of 1615 proteins at p < 0.05, 0 at BH < 0.05. Page 5 of 11.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

leg_ext_1rm, between−person term


130 of 1615 proteins at p < 0.05, 0 at BH < 0.05. Page 6 of 11.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

leg_ext_1rm, between−person term


130 of 1615 proteins at p < 0.05, 0 at BH < 0.05. Page 7 of 11.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

leg_ext_1rm, between−person term


130 of 1615 proteins at p < 0.05, 0 at BH < 0.05. Page 8 of 11.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

leg_ext_1rm, between−person term


130 of 1615 proteins at p < 0.05, 0 at BH < 0.05. Page 9 of 11.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

leg_ext_1rm, between−person term


130 of 1615 proteins at p < 0.05, 0 at BH < 0.05. Page 10 of 11.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

leg_ext_1rm, between−person term


130 of 1615 proteins at p < 0.05, 0 at BH < 0.05. Page 11 of 11.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

type1_share, within−person term


56 of 1640 proteins at p < 0.05, 0 at BH < 0.05. Page 1 of 5.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

type1_share, within−person term


56 of 1640 proteins at p < 0.05, 0 at BH < 0.05. Page 2 of 5.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

type1_share, within−person term


56 of 1640 proteins at p < 0.05, 0 at BH < 0.05. Page 3 of 5.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

type1_share, within−person term


56 of 1640 proteins at p < 0.05, 0 at BH < 0.05. Page 4 of 5.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

type1_share, within−person term


56 of 1640 proteins at p < 0.05, 0 at BH < 0.05. Page 5 of 5.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

type1_share, between−person term


51 of 1640 proteins at p < 0.05, 0 at BH < 0.05. Page 1 of 5.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

type1_share, between−person term


51 of 1640 proteins at p < 0.05, 0 at BH < 0.05. Page 2 of 5.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

type1_share, between−person term


51 of 1640 proteins at p < 0.05, 0 at BH < 0.05. Page 3 of 5.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

type1_share, between−person term


51 of 1640 proteins at p < 0.05, 0 at BH < 0.05. Page 4 of 5.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

type1_share, between−person term


51 of 1640 proteins at p < 0.05, 0 at BH < 0.05. Page 5 of 5.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

Change score, training window


1145 of 28204 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 1 of 96.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 2 of 96.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 3 of 96.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 4 of 96.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window

1145 of 28204 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 5 of 96.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 6 of 96.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 7 of 96.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 8 of 96.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 9 of 96.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 10 of 96.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 11 of 96.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 12 of 96.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 13 of 96.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 14 of 96.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 15 of 96.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 16 of 96.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 17 of 96.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 18 of 96.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 19 of 96.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 20 of 96.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 21 of 96.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 22 of 96.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 23 of 96.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 24 of 96.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 25 of 96.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 26 of 96.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 27 of 96.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 28 of 96.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 29 of 96.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 30 of 96.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 31 of 96.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 32 of 96.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window

1145 of 28204 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 33 of 96.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 34 of 96.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 35 of 96.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 36 of 96.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 37 of 96.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 38 of 96.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 39 of 96.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 40 of 96.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 41 of 96.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 42 of 96.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 43 of 96.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 44 of 96.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 45 of 96.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 46 of 96.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 47 of 96.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 48 of 96.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 49 of 96.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 50 of 96.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 51 of 96.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 52 of 96.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 53 of 96.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 54 of 96.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 55 of 96.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 56 of 96.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 57 of 96.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 58 of 96.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 59 of 96.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 60 of 96.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window

1145 of 28204 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 61 of 96.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 62 of 96.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 63 of 96.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 64 of 96.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 65 of 96.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 66 of 96.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 67 of 96.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 68 of 96.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 69 of 96.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 70 of 96.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 71 of 96.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 72 of 96.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 73 of 96.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 74 of 96.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 75 of 96.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 76 of 96.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 77 of 96.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 78 of 96.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 79 of 96.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 80 of 96.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 81 of 96.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 82 of 96.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 83 of 96.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 84 of 96.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 85 of 96.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 86 of 96.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 87 of 96.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 88 of 96.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 89 of 96.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 90 of 96.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 91 of 96.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 92 of 96.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 93 of 96.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 94 of 96.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 95 of 96.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window


1145 of 28204 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 96 of 96.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 1 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 2 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 3 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 4 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 5 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 6 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 7 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 8 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 9 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 10 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 11 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 12 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 13 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 14 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 15 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 16 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 17 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 18 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 19 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 20 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 21 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 22 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 23 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 24 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 25 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window

1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 26 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 27 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 28 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 29 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 30 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 31 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 32 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 33 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 34 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 35 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 36 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 37 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 38 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 39 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 40 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 41 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 42 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 43 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 44 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 45 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 46 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 47 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 48 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 49 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 50 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 51 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window

1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 52 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 53 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 54 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 55 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 56 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 57 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 58 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 59 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 60 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 61 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 62 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 63 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 64 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 65 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 66 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 67 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 68 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 69 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 70 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 71 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 72 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 73 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 74 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 75 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 76 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 77 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 78 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 79 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 80 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 81 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 82 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 83 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 84 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 85 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 86 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 87 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 88 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 89 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 90 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 91 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 92 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window

1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 93 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 94 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window

1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 95 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 96 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 97 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 98 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 99 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 100 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 101 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 102 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 103 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 104 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 105 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 106 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 107 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 108 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 109 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 110 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 111 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 112 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 113 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 114 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 115 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 116 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 117 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 118 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 119 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 120 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 121 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 122 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 123 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 124 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 125 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 126 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 127 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 128 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


1548 of 31402 feature−outcome pairs at p < 0.05, 1 at BH < 0.05. Page 129 of 129.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 1 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 2 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 3 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 4 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 5 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 6 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 7 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 8 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 9 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 10 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 11 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 12 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 13 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 14 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 15 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 16 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 17 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 18 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 19 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 20 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 21 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 22 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 23 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 24 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 25 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 26 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 27 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 28 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 29 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 30 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window

1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 31 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 32 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 33 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 34 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 35 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 36 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 37 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 38 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 39 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 40 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 41 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 42 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 43 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 44 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 45 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window

1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 46 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 47 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 48 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 49 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 50 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 51 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 52 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 53 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 54 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 55 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 56 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 57 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 58 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 59 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 60 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 61 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 62 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 63 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 64 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 65 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 66 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 67 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 68 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 69 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 70 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 71 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 72 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 73 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 74 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 75 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 76 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window

1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 77 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 78 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window

1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 79 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 80 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 81 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 82 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 83 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 84 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 85 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 86 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 87 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 88 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window

1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 89 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 90 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 91 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 92 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 93 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window

1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 94 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 95 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 96 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 97 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 98 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 99 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 100 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 101 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 102 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 103 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 104 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 105 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 106 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 107 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 108 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 109 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 110 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 111 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 112 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 113 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 114 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 115 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 116 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 117 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 118 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 119 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 120 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 121 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window

1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 122 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 123 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 124 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 125 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 126 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 127 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 128 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 129 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 130 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 131 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window

1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 132 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 133 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window

1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 134 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 135 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 136 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 137 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 138 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 139 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window


1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 140 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window

1683 of 29390 feature−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 141 of 141.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.