



Nominal hits relative to chance, by outcome and window
Spearman per protein, subjects pooled across arms



Bar length is nominal hits ($p < 0.05$) over 5% of proteins tested. The vertical line is chance. Table: c_data/03_phenotype.xlsx, chance_expectation.


Protein association hits, training change (T2 – T1) (1/7)

Spearman against each phenotype; 453 proteins nominal in at least one column; rows 1–75



Protein association hits, training change (T2 – T1) (2/7)


Spearman against each phenotype; 453 proteins nominal in at least one column; rows 76–150



Rows: proteins at nominal p < 0.05 against at least one phenotype, most columns first, then by best p. Filled dot: nominal, fill = Spearman rho, size = -log10 p. Black ring: BH < 0.05. Grey speck: tested, not nominal. Table: c_data/03_phenotype

Protein association hits, training change (T2 – T1) (3/7)


Spearman against each phenotype; 453 proteins nominal in at least one column; rows 151–225



Rows: proteins at nominal $p < 0.05$ against at least one phenotype, most columns first, then by best p . Filled dot: nominal, fill = Spearman rho, size = $-\log_{10} p$. Black ring: BH < 0.05. Grey speck: tested, not nominal. Table: c_data/03_phenotype.x

Protein association hits, training change (T2 – T1) (4/7)


Spearman against each phenotype; 453 proteins nominal in at least one column; rows 226–300



Rows: proteins at nominal p < 0.05 against at least one phenotype, most columns first, then by best p. Filled dot: nominal, fill = Spearman rho, size = -log₁₀ p. Black ring: BH < 0.05. Grey speck: tested, not nominal. Table: c_data/03_phenotype.

Protein association hits, training change (T2 – T1) (5/7)


Spearman against each phenotype; 453 proteins nominal in at least one column; rows 301–375



Rows: proteins at nominal $p < 0.05$ against at least one phenotype, most columns first, then by best p . Filled dot: nominal, fill = Spearman ρ , size = $-\log_{10} p$. Black ring: BH < 0.05. Grey speck: tested, not nominal. Table: c_data/03_phenotype.


Protein association hits, training change (T2 – T1) (6/7)

Spearman against each phenotype; 453 proteins nominal in at least one column; rows 376–450



Rows: proteins at nominal $p < 0.05$ against at least one phenotype, most columns first, then by best p . Filled dot: nominal, fill = Spearman ρ , size = $-\log_{10} p$. Black ring: BH < 0.05. Grey speck: tested, not nominal. Table: c_data/03_phenotype.


Protein association hits, training change (T2 – T1) (7/7)
Spearman against each phenotype; 453 proteins nominal in at least one column; rows 451–453



Rows: proteins at nominal p < 0.05 against at least one phenotype, most columns first, then by best p. Filled dot: nominal, fill = Spearman rho, size = -log10 p. Black ring: BH < 0.05. Grey speck: tested, not nominal. Table: c_data/03_phenotype.

Protein association hits, level at T1 (1/7)


Spearman against each phenotype; 519 proteins nominal in at least one column; rows 1–75



Rows: proteins at nominal p < 0.05 against at least one phenotype, most columns first, then by best p. Filled dot: nominal, fill = Spearman rho, size = -log10 p. Black ring: BH < 0.05. Grey speck: tested, not nominal. Table: c_data/03_phenotype.p

Protein association hits, level at T1 (2/7)


Spearman against each phenotype; 519 proteins nominal in at least one column; rows 76-150



Rows: proteins at nominal $p < 0.05$ against at least one phenotype, most columns first, then by best p. Filled dot: nominal, fill = Spearman rho, size = $-\log_{10} p$. Black ring: BH < 0.05. Grey speck: tested, not nominal. Table: c_data/03_phenotype.


Protein association hits, level at T1 (3/7)

Spearman against each phenotype; 519 proteins nominal in at least one column; rows 151–225




Rows: proteins at nominal $p < 0.05$ against at least one phenotype, most columns first, then by best p. Filled dot: nominal, fill = Spearman rho, size = $-\log_{10} p$. Black ring: BH < 0.05 . Grey speck: tested, not nominal. Table: c_data/03_phenotype.

Protein association hits, level at T1 (4/7)
 Spearman against each phenotype; 519 proteins nominal in at least one column; rows 226–300





Spearman against each phenotype; 519 proteins nominal in at least one column; rows 301–375



Rows: proteins at nominal $p < 0.05$ against at least one phenotype, most columns first, then by best p . Filled dot: nominal, fill = Spearman rho, size = $-\log_{10} p$. Black ring: BH < 0.05 . Grey speck: tested, not nominal. Table: c_data/03_phenotype


Spearman association hits, level at T1 (6/7)
Spearman against each phenotype; 519 proteins nominal in at least one column; rows 376–450






Protein association hits, acute change (T3 – T2) (1/6)

Spearman against each phenotype; 435 proteins nominal in at least one column; rows 1–75



Protein association hits, acute change (T3 – T2) (2/6)


Spearman against each phenotype; 435 proteins nominal in at least one column; rows 76–150



Rows: proteins at nominal $p < 0.05$ against at least one phenotype, most columns first, then by best p. Filled dot: nominal, fill = Spearman rho, size = $-\log_{10} p$. Black ring: BH < 0.05 . Grey speck: tested, not nominal. Table: c_data/03_phenotype.

Protein association hits, acute change (T3 – T2) (3/6)


Spearman against each phenotype; 435 proteins nominal in at least one column; rows 151–225



Rows: proteins at nominal $p < 0.05$ against at least one phenotype, most columns first, then by best p. Filled dot: nominal, fill = Spearman rho, size = $-\log_{10} p$. Black ring: BH < 0.05. Grey speck: tested, not nominal. Table: c_data/03_phenotype

Protein association hits, acute change (T3 – T2) (4/6)


Spearman against each phenotype; 435 proteins nominal in at least one column; rows 226–300



Rows: proteins at nominal p < 0.05 against at least one phenotype, most columns first, then by best p. Filled dot: nominal, fill = Spearman rho, size = $-\log_{10} p$. Black ring: BH < 0.05. Grey speck: tested, not nominal. Table: c_data/03_phenotype


Protein association hits, acute change (T3 – T2) (5/6)

Spearman against each phenotype; 435 proteins nominal in at least one column; rows 301–375



Rows: proteins at nominal $p < 0.05$ against at least one phenotype, most columns first, then by best p . Filled dot: nominal, fill = Spearman rho, size = $-\log_{10} p$. Black ring: BH < 0.05. Grey speck: tested, not nominal. Table: c_data/03_phenotype.

Spearman against each phenotype; 435 proteins nominal in at least one column; rows 376–435




Rows: proteins at nominal $p < 0.05$ against at least one phenotype, most columns first, then by best p . Filled dot: nominal, fill = Spearman rho, size = $-\log_{10} p$. Black ring: BH < 0.05 . Grey speck: tested, not nominal. Table: c_data/03_phenotype.x

Protein against phenotype, training change (T2 – T1)

Spearman, subjects pooled across arms

HR LR




All 633 protein–outcome pairs at nominal $p < 0.05$, by p. Lines are fitted within each arm; the header ρ is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, training change (T2 – T1)

Spearman, subjects pooled across arms

HR LR




All 633 protein–outcome pairs at nominal $p < 0.05$, by p . Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, training change (T2 – T1)

Spearman, subjects pooled across arms

HR LR




All 633 protein–outcome pairs at nominal $p < 0.05$, by p . Lines are fitted within each arm; the header ρ is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, training change (T2 – T1)

Spearman, subjects pooled across arms

HR LR




All 633 protein–outcome pairs at nominal $p < 0.05$, by p . Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, training change (T2 – T1)

Spearman, subjects pooled across arms

— HR — LR




All 633 protein–outcome pairs at nominal p < 0.05, by p. Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, training change (T2 – T1)

Spearman, subjects pooled across arms


HR LR



All 633 protein–outcome pairs at nominal $p < 0.05$, by p . Lines are fitted within each arm; the header ρ is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, training change (T2 – T1)


Spearman, subjects pooled across arms



All 633 protein–outcome pairs at nominal p < 0.05, by p. Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, training change (T2 – T1)


Spearman, subjects pooled across arms



All 633 protein–outcome pairs at nominal $p < 0.05$, by p . Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, training change (T2 – T1)

Spearman, subjects pooled across arms




All 633 protein–outcome pairs at nominal $p < 0.05$, by p . Lines are fitted within each arm; the header ρ is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, training change (T2 – T1)

Spearman, subjects pooled across arms

HR LR




All 633 protein–outcome pairs at nominal p < 0.05, by p. Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, training change (T2 – T1)

Spearman, subjects pooled across arms

HR LR




All 633 protein–outcome pairs at nominal $p < 0.05$, by p . Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, training change (T2 – T1)

Spearman, subjects pooled across arms

HR LR




All 633 protein–outcome pairs at nominal $p < 0.05$, by p . Lines are fitted within each arm; the header ρ is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, training change (T2 – T1)

Spearman, subjects pooled across arms

HR LR




All 633 protein–outcome pairs at nominal $p < 0.05$, by p. Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, training change (T2 – T1)

Spearman, subjects pooled across arms


HR LR



All 633 protein–outcome pairs at nominal $p < 0.05$, by p . Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, training change (T2 – T1)

Spearman, subjects pooled across arms




All 633 protein–outcome pairs at nominal p < 0.05, by p. Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, training change (T2 – T1)

Spearman, subjects pooled across arms

HR LR




All 633 protein–outcome pairs at nominal p < 0.05, by p. Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, training change (T2 – T1)


Spearman, subjects pooled across arms

HR LR



Protein against phenotype, training change (T2 – T1)

Spearman, subjects pooled across arms




All 633 protein–outcome pairs at nominal p < 0.05, by p. Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, training change (T2 – T1)

Spearman, subjects pooled across arms


HR LR



All 633 protein–outcome pairs at nominal $p < 0.05$, by p . Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, training change (T2 – T1)

Spearman, subjects pooled across arms




All 633 protein–outcome pairs at nominal $p < 0.05$, by p . Lines are fitted within each arm; the header ρ is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, training change (T2 – T1)

Spearman, subjects pooled across arms

HR LR




All 633 protein–outcome pairs at nominal p < 0.05, by p. Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, training change (T2 – T1)

Spearman, subjects pooled across arms


HR LR



All 633 protein–outcome pairs at nominal p < 0.05, by p. Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, training change (T2 – T1)

Spearman, subjects pooled across arms




All 633 protein–outcome pairs at nominal p < 0.05, by p. Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, training change (T2 – T1)

Spearman, subjects pooled across arms

HR LR




All 633 protein–outcome pairs at nominal $p < 0.05$, by p . Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, training change (T2 – T1)

Spearman, subjects pooled across arms

HR LR




All 633 protein–outcome pairs at nominal $p < 0.05$, by p . Lines are fitted within each arm; the header ρ is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, training change (T2 – T1)

Spearman, subjects pooled across arms


HR LR



All 633 protein–outcome pairs at nominal p < 0.05, by p. Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, training change (T2 – T1)

Spearman, subjects pooled across arms




All 633 protein–outcome pairs at nominal p < 0.05, by p. Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, training change (T2 – T1)

Spearman, subjects pooled across arms

HR LR















All 633 protein–outcome pairs at nominal $p < 0.05$, by p . Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, training change (T2 – T1)

Spearman, subjects pooled across arms

HR LR




log2 abundance, training change (T2 – T1), one point per subject

Protein against phenotype, training change (T2 – T1)

Spearman, subjects pooled across arms

HR LR




All 633 protein–outcome pairs at nominal p < 0.05, by p. Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, training change (T2 – T1)

Spearman, subjects pooled across arms

HR LR




All 633 protein–outcome pairs at nominal p < 0.05, by p. Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, training change (T2 – T1)

Spearman, subjects pooled across arms


HR LR



All 633 protein-outcome pairs at nominal p < 0.05, by p. Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, training change (T2 – T1)


Spearman, subjects pooled across arms



All 633 protein–outcome pairs at nominal p < 0.05, by p. Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, training change (T2 – T1)

Spearman, subjects pooled across arms




All 633 protein–outcome pairs at nominal p < 0.05, by p. Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, training change (T2 – T1)

Spearman, subjects pooled across arms

HR LR




All 633 protein–outcome pairs at nominal p < 0.05, by p. Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, training change (T2 – T1)

Spearman, subjects pooled across arms

HR LR




All 633 protein–outcome pairs at nominal p < 0.05, by p. Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, training change (T2 – T1)

Spearman, subjects pooled across arms

HR LR




All 633 protein-outcome pairs at nominal p < 0.05, by p. Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, training change (T2 – T1)

Spearman, subjects pooled across arms


HR LR



All 633 protein–outcome pairs at nominal p < 0.05, by p. Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, training change (T2 – T1)

Spearman, subjects pooled across arms




All 633 protein–outcome pairs at nominal p < 0.05, by p. Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, training change (T2 – T1)

Spearman, subjects pooled across arms

HR LR




All 633 protein–outcome pairs at nominal $p < 0.05$, by p . Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, training change (T2 – T1)

Spearman, subjects pooled across arms

HR LR




All 633 protein–outcome pairs at nominal $p < 0.05$, by p. Lines are fitted within each arm; the header ρ is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, training change (T2 – T1)

Spearman, subjects pooled across arms


HR LR



All 633 protein-outcome pairs at nominal p < 0.05, by p. Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, training change (T2 – T1)

Spearman, subjects pooled across arms




All 633 protein–outcome pairs at nominal $p < 0.05$, by p. Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, training change (T2 – T1)

Spearman, subjects pooled across arms


HR LR



All 633 protein–outcome pairs at nominal p < 0.05, by p. Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, training change (T2 – T1)


Spearman, subjects pooled across arms



All 633 protein–outcome pairs at nominal p < 0.05, by p. Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, training change (T2 – T1)

Spearman, subjects pooled across arms




All 633 protein–outcome pairs at nominal $p < 0.05$, by p. Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, training change (T2 – T1)

Spearman, subjects pooled across arms

HR LR




All 633 protein–outcome pairs at nominal $p < 0.05$, by p . Lines are fitted within each arm; the header ρ is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, training change (T2 – T1)

Spearman, subjects pooled across arms

HR LR




All 633 protein–outcome pairs at nominal $p < 0.05$, by p . Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, training change (T2 – T1)

Spearman, subjects pooled across arms

HR LR




All 633 protein–outcome pairs at nominal $p < 0.05$, by p . Lines are fitted within each arm; the header ρ is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, training change (T2 – T1)

Spearman, subjects pooled across arms

HR LR















All 633 protein–outcome pairs at nominal p < 0.05, by p. Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, training change (T2 – T1)

Spearman, subjects pooled across arms

HR LR




log2 abundance, training change (T2 – T1), one point per subject

Protein against phenotype, training change (T2 – T1)

Spearman, subjects pooled across arms

HR LR




All 633 protein–outcome pairs at nominal $p < 0.05$, by p . Lines are fitted within each arm; the header ρ is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, training change (T2 – T1)

Spearman, subjects pooled across arms

HR LR




log2 abundance, training change (T2 – T1), one point per subject

All 633 protein–outcome pairs at nominal p < 0.05, by p. Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, level at T1

Spearman, subjects pooled across arms

HR LR




All 783 protein-outcome pairs at nominal $p < 0.05$, by p . Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, level at T1

Spearman, subjects pooled across arms

HR LR




All 783 protein–outcome pairs at nominal $p < 0.05$, by p . Lines are fitted within each arm; the header ρ is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, level at T1

Spearman, subjects pooled across arms

HR LR




All 783 protein–outcome pairs at nominal $p < 0.05$, by p . Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, level at T1

Spearman, subjects pooled across arms

HR LR




All 783 protein-outcome pairs at nominal $p < 0.05$, by p . Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, level at T1

Spearman, subjects pooled across arms

HR LR




All 783 protein-outcome pairs at nominal $p < 0.05$, by p. Lines are fitted within each arm; the header ρ is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, level at T1

Spearman, subjects pooled across arms

HR LR




All 783 protein-outcome pairs at nominal $p < 0.05$, by p. Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, level at T1

Spearman, subjects pooled across arms

HR LR




All 783 protein-outcome pairs at nominal $p < 0.05$, by p. Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, level at T1

Spearman, subjects pooled across arms

HR LR




All 783 protein-outcome pairs at nominal p < 0.05, by p. Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, level at T1

Spearman, subjects pooled across arms


HR LR



All 783 protein-outcome pairs at nominal p < 0.05, by p. Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, level at T1

Spearman, subjects pooled across arms




All 783 protein–outcome pairs at nominal $p < 0.05$, by p. Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, level at T1

Spearman, subjects pooled across arms

HR LR




All 783 protein-outcome pairs at nominal p < 0.05, by p. Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, level at T1

Spearman, subjects pooled across arms


HR LR



All 783 protein-outcome pairs at nominal p < 0.05, by p. Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, level at T1

Spearman, subjects pooled across arms




All 783 protein–outcome pairs at nominal p < 0.05, by p. Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, level at T1

Spearman, subjects pooled across arms


HR LR



All 783 protein–outcome pairs at nominal p < 0.05, by p. Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, level at T1


Spearman, subjects pooled across arms



All 783 protein-outcome pairs at nominal p < 0.05, by p. Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, level at T1


Spearman, subjects pooled across arms



All 783 protein-outcome pairs at nominal $p < 0.05$, by p. Lines are fitted within each arm; the header ρ is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, level at T1


Spearman, subjects pooled across arms



All 783 protein–outcome pairs at nominal $p < 0.05$, by p . Lines are fitted within each arm; the header ρ is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, level at T1

Spearman, subjects pooled across arms




All 783 protein-outcome pairs at nominal $p < 0.05$, by p. Lines are fitted within each arm; the header ρ is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, level at T1

Spearman, subjects pooled across arms


HR LR



All 783 protein-outcome pairs at nominal $p < 0.05$, by p. Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, level at T1

Spearman, subjects pooled across arms




All 783 protein-outcome pairs at nominal $p < 0.05$, by p. Lines are fitted within each arm; the header ρ is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, level at T1

Spearman, subjects pooled across arms


HR LR



All 783 protein-outcome pairs at nominal p < 0.05, by p. Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, level at T1


Spearman, subjects pooled across arms



All 783 protein–outcome pairs at nominal $p < 0.05$, by p . Lines are fitted within each arm; the header ρ is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, level at T1

Spearman, subjects pooled across arms




All 783 protein-outcome pairs at nominal $p < 0.05$, by p. Lines are fitted within each arm; the header ρ is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, level at T1

Spearman, subjects pooled across arms


HR LR



All 783 protein-outcome pairs at nominal p < 0.05, by p. Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, level at T1


Spearman, subjects pooled across arms



All 783 protein-outcome pairs at nominal $p < 0.05$, by p. Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, level at T1

Spearman, subjects pooled across arms




All 783 protein-outcome pairs at nominal $p < 0.05$, by p. Lines are fitted within each arm; the header ρ is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, level at T1

Spearman, subjects pooled across arms


HR LR



All 783 protein-outcome pairs at nominal $p < 0.05$, by p. Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, level at T1

Spearman, subjects pooled across arms




All 783 protein–outcome pairs at nominal p < 0.05, by p. Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, level at T1

Spearman, subjects pooled across arms

HR LR




All 783 protein–outcome pairs at nominal $p < 0.05$, by p. Lines are fitted within each arm; the header ρ is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, level at T1

Spearman, subjects pooled across arms


HR LR



All 783 protein-outcome pairs at nominal p < 0.05, by p. Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, level at T1


Spearman, subjects pooled across arms



All 783 protein–outcome pairs at nominal $p < 0.05$, by p . Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, level at T1

Spearman, subjects pooled across arms




All 783 protein-outcome pairs at nominal p < 0.05, by p. Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, level at T1

Spearman, subjects pooled across arms


HR LR



All 783 protein-outcome pairs at nominal p < 0.05, by p. Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, level at T1

Spearman, subjects pooled across arms




All 783 protein–outcome pairs at nominal $p < 0.05$, by p. Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, level at T1

Spearman, subjects pooled across arms


HR LR



All 783 protein–outcome pairs at nominal $p < 0.05$, by p . Lines are fitted within each arm; the header ρ is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, level at T1

Spearman, subjects pooled across arms




All 783 protein-outcome pairs at nominal p < 0.05, by p. Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, level at T1

Spearman, subjects pooled across arms

HR LR




All 783 protein-outcome pairs at nominal p < 0.05, by p. Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, level at T1

Spearman, subjects pooled across arms


HR LR



All 783 protein-outcome pairs at nominal $p < 0.05$, by p. Lines are fitted within each arm; the header ρ is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, level at T1


Spearman, subjects pooled across arms



All 783 protein-outcome pairs at nominal p < 0.05, by p. Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, level at T1


Spearman, subjects pooled across arms



All 783 protein–outcome pairs at nominal $p < 0.05$, by p . Lines are fitted within each arm; the header ρ is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, level at T1


Spearman, subjects pooled across arms



All 783 protein–outcome pairs at nominal $p < 0.05$, by p . Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, level at T1


Spearman, subjects pooled across arms



All 783 protein-outcome pairs at nominal p < 0.05, by p. Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, level at T1


Spearman, subjects pooled across arms



All 783 protein–outcome pairs at nominal p < 0.05, by p. Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, level at T1


Spearman, subjects pooled across arms



Protein against phenotype, level at T1

Spearman, subjects pooled across arms


HR LR



All 783 protein-outcome pairs at nominal $p < 0.05$, by p. Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, level at T1

Spearman, subjects pooled across arms




All 783 protein-outcome pairs at nominal p < 0.05, by p. Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, level at T1

Spearman, subjects pooled across arms


HR LR



All 783 protein-outcome pairs at nominal p < 0.05, by p. Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, level at T1


Spearman, subjects pooled across arms



All 783 protein-outcome pairs at nominal $p < 0.05$, by p. Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, level at T1

Spearman, subjects pooled across arms




All 783 protein-outcome pairs at nominal $p < 0.05$, by p. Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, level at T1

Spearman, subjects pooled across arms


HR LR



All 783 protein-outcome pairs at nominal $p < 0.05$, by p. Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, level at T1

Spearman, subjects pooled across arms




All 783 protein-outcome pairs at nominal p < 0.05, by p. Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, level at T1

Spearman, subjects pooled across arms

HR LR




All 783 protein–outcome pairs at nominal $p < 0.05$, by p. Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, level at T1

Spearman, subjects pooled across arms

HR LR




All 783 protein-outcome pairs at nominal p < 0.05, by p. Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, level at T1

Spearman, subjects pooled across arms


HR LR



All 783 protein–outcome pairs at nominal p < 0.05, by p. Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, level at T1

Spearman, subjects pooled across arms




All 783 protein-outcome pairs at nominal p < 0.05, by p. Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, level at T1

Spearman, subjects pooled across arms


HR LR



All 783 protein-outcome pairs at nominal $p < 0.05$, by p. Lines are fitted within each arm; the header ρ is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, level at T1

Spearman, subjects pooled across arms




All 783 protein-outcome pairs at nominal p < 0.05, by p. Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, level at T1

Spearman, subjects pooled across arms


HR LR



All 783 protein-outcome pairs at nominal p < 0.05, by p. Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, level at T1


Spearman, subjects pooled across arms



All 783 protein-outcome pairs at nominal p < 0.05, by p. Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, level at T1


Spearman, subjects pooled across arms



All 783 protein-outcome pairs at nominal p < 0.05, by p. Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, level at T1

Spearman, subjects pooled across arms




All 783 protein-outcome pairs at nominal $p < 0.05$, by p. Lines are fitted within each arm; the header ρ is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, level at T1

Spearman, subjects pooled across arms


HR LR



All 783 protein-outcome pairs at nominal $p < 0.05$, by p. Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, level at T1

Spearman, subjects pooled across arms




All 783 protein-outcome pairs at nominal p < 0.05, by p. Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, level at T1

Spearman, subjects pooled across arms


HR LR



All 783 protein–outcome pairs at nominal $p < 0.05$, by p. Lines are fitted within each arm; the header ρ is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, level at T1

Spearman, subjects pooled across arms




All 783 protein–outcome pairs at nominal p < 0.05, by p. Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, level at T1

Spearman, subjects pooled across arms


HR LR



All 783 protein-outcome pairs at nominal p < 0.05, by p. Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, acute change (T3 – T2)


Spearman, subjects pooled across arms



All 750 protein–outcome pairs at nominal $p < 0.05$, by p . Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, acute change (T3 – T2)


Spearman, subjects pooled across arms



All 750 protein–outcome pairs at nominal $p < 0.05$, by p . Lines are fitted within each arm; the header ρ is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, acute change (T3 – T2)

Spearman, subjects pooled across arms




All 750 protein–outcome pairs at nominal p < 0.05, by p. Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, acute change (T3 – T2)

Spearman, subjects pooled across arms


HR LR



All 750 protein–outcome pairs at nominal $p < 0.05$, by p . Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, acute change (T3 – T2)


Spearman, subjects pooled across arms



All 750 protein–outcome pairs at nominal $p < 0.05$, by p . Lines are fitted within each arm; the header ρ is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, acute change (T3 – T2)

Spearman, subjects pooled across arms




All 750 protein–outcome pairs at nominal $p < 0.05$, by p . Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, acute change (T3 – T2)

Spearman, subjects pooled across arms

HR LR




All 750 protein–outcome pairs at nominal $p < 0.05$, by p . Lines are fitted within each arm; the header ρ is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, acute change (T3 – T2)

Spearman, subjects pooled across arms

HR LR




All 750 protein–outcome pairs at nominal $p < 0.05$, by p . Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, acute change (T3 – T2)

Spearman, subjects pooled across arms

HR LR















All 750 protein–outcome pairs at nominal $p < 0.05$, by p . Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, acute change (T3 – T2)

Spearman, subjects pooled across arms

HR LR




log2 abundance, acute change (T3 – T2), one point per subject

All 750 protein–outcome pairs at nominal p < 0.05, by p. Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, acute change (T3 – T2)

Spearman, subjects pooled across arms




All 750 protein–outcome pairs at nominal p < 0.05, by p. Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, acute change (T3 – T2)

Spearman, subjects pooled across arms

HR LR




All 750 protein–outcome pairs at nominal p < 0.05, by p. Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, acute change (T3 – T2)

Spearman, subjects pooled across arms

HR LR




All 750 protein-outcome pairs at nominal p < 0.05, by p. Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, acute change (T3 – T2)

Spearman, subjects pooled across arms


HR LR



All 750 protein–outcome pairs at nominal p < 0.05, by p. Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, acute change (T3 – T2)


Spearman, subjects pooled across arms



All 750 protein–outcome pairs at nominal $p < 0.05$, by p . Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, acute change (T3 – T2)

Spearman, subjects pooled across arms




All 750 protein–outcome pairs at nominal p < 0.05, by p. Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, acute change (T3 – T2)

Spearman, subjects pooled across arms


HR LR



All 750 protein–outcome pairs at nominal $p < 0.05$, by p . Lines are fitted within each arm; the header ρ is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, acute change (T3 – T2)


Spearman, subjects pooled across arms



All 750 protein–outcome pairs at nominal p < 0.05, by p. Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, acute change (T3 – T2)

Spearman, subjects pooled across arms




All 750 protein–outcome pairs at nominal $p < 0.05$, by p . Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, acute change (T3 – T2)

Spearman, subjects pooled across arms


HR LR



All 750 protein–outcome pairs at nominal $p < 0.05$, by p . Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, acute change (T3 – T2)

Spearman, subjects pooled across arms




All 750 protein–outcome pairs at nominal p < 0.05, by p. Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, acute change (T3 – T2)

Spearman, subjects pooled across arms

HR LR




All 750 protein–outcome pairs at nominal p < 0.05, by p. Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, acute change (T3 – T2)

Spearman, subjects pooled across arms


HR LR



All 750 protein–outcome pairs at nominal $p < 0.05$, by p. Lines are fitted within each arm; the header ρ is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, acute change (T3 – T2)


Spearman, subjects pooled across arms



All 750 protein–outcome pairs at nominal $p < 0.05$, by p . Lines are fitted within each arm; the header ρ is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, acute change (T3 – T2)

Spearman, subjects pooled across arms















All 750 protein–outcome pairs at nominal p < 0.05, by p. Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, acute change (T3 – T2)

Spearman, subjects pooled across arms

HR LR




log2 abundance, acute change (T3 – T2), one point per subject

All 750 protein–outcome pairs at nominal p < 0.05, by p. Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.


Protein against phenotype, acute change (T3 – T2)

Spearman, subjects pooled across arms

HR LR




All 750 protein–outcome pairs at nominal $p < 0.05$, by p. Lines are fitted within each arm; the header ρ is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.



Protein against phenotype, acute change (T3 – T2)

Spearman, subjects pooled across arms


HR LR



All 750 protein–outcome pairs at nominal $p < 0.05$, by p . Lines are fitted within each arm; the header ρ is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, acute change (T3 – T2)


Spearman, subjects pooled across arms



All 750 protein–outcome pairs at nominal $p < 0.05$, by p . Lines are fitted within each arm; the header ρ is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, acute change (T3 – T2)

Spearman, subjects pooled across arms




All 750 protein–outcome pairs at nominal $p < 0.05$, by p . Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, acute change (T3 – T2)

Spearman, subjects pooled across arms

HR LR




All 750 protein-outcome pairs at nominal p < 0.05, by p. Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, acute change (T3 – T2)

Spearman, subjects pooled across arms


HR LR



All 750 protein–outcome pairs at nominal p < 0.05, by p. Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, acute change (T3 – T2)


Spearman, subjects pooled across arms



All 750 protein–outcome pairs at nominal p < 0.05, by p. Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, acute change (T3 – T2)


Spearman, subjects pooled across arms



All 750 protein–outcome pairs at nominal $p < 0.05$, by p . Lines are fitted within each arm; the header ρ is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, acute change (T3 – T2)


Spearman, subjects pooled across arms



All 750 protein–outcome pairs at nominal p < 0.05, by p. Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, acute change (T3 – T2)

Spearman, subjects pooled across arms




All 750 protein–outcome pairs at nominal p < 0.05, by p. Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, acute change (T3 – T2)

Spearman, subjects pooled across arms

HR LR




All 750 protein–outcome pairs at nominal p < 0.05, by p. Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, acute change (T3 – T2)

Spearman, subjects pooled across arms

HR LR















All 750 protein–outcome pairs at nominal $p < 0.05$, by p . Lines are fitted within each arm; the header ρ is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, acute change (T3 – T2)

Spearman, subjects pooled across arms

HR LR




log2 abundance, acute change (T3 – T2), one point per subject

All 750 protein–outcome pairs at nominal p < 0.05, by p. Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, acute change (T3 – T2)


Spearman, subjects pooled across arms



All 750 protein–outcome pairs at nominal $p < 0.05$, by p . Lines are fitted within each arm; the header ρ is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, acute change (T3 – T2)

Spearman, subjects pooled across arms




All 750 protein–outcome pairs at nominal p < 0.05, by p. Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, acute change (T3 – T2)

Spearman, subjects pooled across arms


HR LR



All 750 protein-outcome pairs at nominal $p < 0.05$, by p. Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, acute change (T3 – T2)

Spearman, subjects pooled across arms




All 750 protein–outcome pairs at nominal $p < 0.05$, by p . Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, acute change (T3 – T2)

Spearman, subjects pooled across arms


HR LR



All 750 protein–outcome pairs at nominal p < 0.05, by p. Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, acute change (T3 – T2)

Spearman, subjects pooled across arms




All 750 protein–outcome pairs at nominal $p < 0.05$, by p . Lines are fitted within each arm; the header ρ is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, acute change (T3 – T2)

Spearman, subjects pooled across arms

HR LR




All 750 protein-outcome pairs at nominal p < 0.05, by p. Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, acute change (T3 – T2)

Spearman, subjects pooled across arms


HR LR



All 750 protein–outcome pairs at nominal $p < 0.05$, by p . Lines are fitted within each arm; the header ρ is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, acute change (T3 – T2)


Spearman, subjects pooled across arms



All 750 protein–outcome pairs at nominal p < 0.05, by p. Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, acute change (T3 – T2)

Spearman, subjects pooled across arms




All 750 protein–outcome pairs at nominal p < 0.05, by p. Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, acute change (T3 – T2)

Spearman, subjects pooled across arms


HR LR



All 750 protein–outcome pairs at nominal p < 0.05, by p. Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, acute change (T3 – T2)

Spearman, subjects pooled across arms




All 750 protein–outcome pairs at nominal $p < 0.05$, by p . Lines are fitted within each arm; the header ρ is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, acute change (T3 – T2)

Spearman, subjects pooled across arms

HR LR




All 750 protein–outcome pairs at nominal $p < 0.05$, by p . Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.



Protein against phenotype, acute change (T3 – T2)

Spearman, subjects pooled across arms




All 750 protein–outcome pairs at nominal p < 0.05, by p. Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, acute change (T3 – T2)

Spearman, subjects pooled across arms

HR LR




All 750 protein–outcome pairs at nominal $p < 0.05$, by p . Lines are fitted within each arm; the header ρ is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, acute change (T3 – T2)

Spearman, subjects pooled across arms

HR LR




All 750 protein-outcome pairs at nominal p < 0.05, by p. Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, acute change (T3 – T2)

Spearman, subjects pooled across arms


HR LR



All 750 protein–outcome pairs at nominal p < 0.05, by p. Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, acute change (T3 – T2)


Spearman, subjects pooled across arms



All 750 protein–outcome pairs at nominal $p < 0.05$, by p . Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, acute change (T3 – T2)


Spearman, subjects pooled across arms



All 750 protein–outcome pairs at nominal p < 0.05, by p. Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, acute change (T3 – T2)


Spearman, subjects pooled across arms



All 750 protein–outcome pairs at nominal p < 0.05, by p. Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, acute change (T3 – T2)


Spearman, subjects pooled across arms



All 750 protein–outcome pairs at nominal p < 0.05, by p. Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.

Protein against phenotype, acute change (T3 – T2)

Spearman, subjects pooled across arms



All 750 protein–outcome pairs at nominal p < 0.05, by p. Lines are fitted within each arm; the header rho is pooled, q from BH within window and outcome. Table: c_data/03_phenotype.xlsx, protein_association and protein_by_arm.