
a Where HR and LR responses differ
six strongest fgsea sets per contrast
Anchoring of the basal body to the plasma…
Cellular response to heat
Antigen processing and presentation
Cardiac conduction
Striated muscle contraction
Calcium signaling pathway
Mitochondrial biogenesis
Cristae formation
Mitochondrion organization
Parkinsons disease
Aerobic respiration and respiratory elect…
Training
Acute
interaction contrast
-log10 FDR 2 6 12+
NES -3 -2 -1 0 1 2 3
b OXPHOS catch-up
HR - LR at T1 NES -2.60; training int. +2.25
OXPHOS score (Hallmark, singscore)
0.27
0.25
0.23
T1
T2
T3
HR LR
c Acute OXPHOS response
interaction NES -2.70; fry p = 0.15
OXPHOS score, T3 - T2
0.02
0.00
-0.02
-0.04
-0.06
HR
LR
HR LR
d Sarcomere-strength link by arm
rho HR -1.00, LR +0.48; FDR 0.0045
black (sarcomere) module change, T2 - T1
0.25
0.00
-0.25
-0.50
0
30
60
90
leg-press 1RM gain (%)
HR LR
e HR and LR cannot be told apart
grey: permutation median to 95th percentile
modules T1
modules T2
modules T3
modules training
modules acute
proteins T1
proteins T2
proteins T3
proteins training
proteins acute
DIABLOT1
DIABLOT1 + phenotype
DIABLOT2
DIABLOT2 + phenotype
DIABLOT3
DIABLO training
DIABLO training + phenotype
DIABLO acute
0.0
0.5
1.0
cross-validated AUC