





All 1,900 proteins by 45 biopsies, grouped by arm and time point; column labels are subjects. Fill: log2 abundance minus the protein's mean over biopsies. Colours stop at the 99th percentile of |value|, 1.9. Rows are ordered by average−linkage clustering (1 − Pearson r) of the imputed matrix, for display only. Grey: missing.

# Contrast overview: proteins




All 1,900 proteins by the nine contrasts. Fill: log2 fold change from the limma fit; status: p and BH q within contrast. Colours stop at the 99th percentile of |value|, 1.4. Rows are ordered by average−linkage clustering (1 − Pearson r) of the imputed matrix, for display only. Grey: not tested. Table: c_data/02_contrasts.xlsx, DEP_matrix.




Dashed line: a flat null (tested / 20). A spike at zero on a flat background is signal; a slope up toward 1 is a conservative test. Table: c_data/02_contrasts.xlsx, DEP_matrix.

## Volcano per contrast


49 to 179 proteins at p < 0.05 per contrast; 0 at BH < 0.05 across all nine.




limma via proteoDA on the unimputed matrix, subject as a random effect. Dashed line: p = 0.05. BH within contrast. Table: c_data/02_contrasts.xlsx, DEP_matrix.

ANCOVA: HR minus LR at the later time point, at equal starting value

Training: 62 of 1539 proteins at p < 0.05. Acute: 173 of 1645.




lm(later ~ earlier + arm) per protein; training is T2 on T1, acute T3 on T2. Dashed line: p = 0.05. BH within window. Table: c_data/02_contrasts.xlsx, ancova.