




Female: phenotype effects of PAS
% change against vehicle, Welch t on log values, n = 5 v 5
CSA
Capillary density
Satellite cells / fibre
Soleus pyruvate
Plantaris pyruvate
Soleus octanoyl
p > 0.05
p ≤ 0.05
Female
-50%
+0%
+100%
+300%
+700%
PAS vs vehicle, % change (95% CI, log scale)


Female: phenotype associations by level


This sex's ten mice, pooled across PAS and vehicle or within treatment; p ≤ 0.05 of tests; BH within level, phenotype and model





Check: protein associations with and without pellet mitochondrial share
Share centred within group, added to the association model
FDR ≤ 0.05
p ≤ 0.05
CSA
Capillary density
Satellite cells / fibre
Soleus pyruvate
Plantaris pyruvate
Soleus octanoyl
0
20
40
60
0
100
200
300
Proteins
with pellet mito share
without


Female: Modules, treatment effect and phenotype associations
Pearson r across this sex's ten mice (treatment: PAS = 1); box: FDR ≤ 0.05 (limma, BH within level)
Phenotype associations
Treatment effect
Yellow 0.27 -0.10 0.34 0.47 0.38 -0.57 0.21
Turquoise 0.36 0.27 0.55 -0.05 0.06 -0.33 0.42
Tan -0.43 -0.27 -0.28 -0.33 -0.39 -0.30 -0.43
Salmon 0.19 -0.02 0.14 -0.31 0.01 -0.17 -0.08
Red 0.84 0.66 0.70 0.09 0.26 0.52 0.73
Purple -0.15 0.10 -0.21 -0.37 -0.23 0.05 -0.11
Pink -0.18 -0.14 -0.11 -0.39 -0.06 -0.57 -0.20
Magenta -0.57 -0.56 -0.57 -0.08 -0.34 -0.53 -0.73
Greenyellow -0.21 -0.17 -0.44 -0.04 0.10 0.12 -0.24
Green -0.12 0.15 -0.29 -0.50 -0.16 -0.01 -0.09
Cyan -0.76 -0.69 -0.84 -0.30 -0.30 -0.13 -0.89
Brown -0.29 0.12 -0.34 -0.64 -0.39 0.09 -0.19
Blue -0.21 -0.28 -0.43 -0.08 0.08 0.05 -0.38
Black 0.45 0.18 0.31 0.27 0.33 0.07 0.35
CSA
Capillary density
Satellite cells / fibre
Soleus pyruvate
Plantaris pyruvate
Soleus octanoyl
PAS vs vehicle
r
1.0
0.5
0.0
-0.5
-1.0



Female: Complexes, treatment effect and phenotype associations
Pearson r across this sex's ten mice (treatment: PAS = 1); box: FDR ≤ 0.05 (limma, BH within level)
Phenotype associations
Treatment effect
Tnfalpha Nfkappa B Signaling Complex 5232 0.04 0.29 0.17 -0.39 -0.26 0.12 0.11
Spliceosome E Complex 8391 -0.21 -0.36 -0.37 0.09 0.01 -0.19 -0.44
Spliceosome A Complex 8372 -0.29 -0.43 -0.40 -0.08 -0.09 -0.25 -0.58
Spermatoproteasome 8532 -0.52 -0.23 -0.61 -0.32 -0.24 0.53 -0.53
Sarcoglycan Sarcospan…trobrevin Complex 349 -0.37 -0.49 -0.17 0.10 -0.04 -0.66 -0.39
Sarcoglycan Sarcospan…troglycan Complex 346 -0.27 -0.38 -0.11 0.21 0.06 -0.73 -0.21
Sarcoglycan Sarcospan Complex Sg Spn 343 -0.31 -0.43 -0.15 0.36 0.04 -0.65 -0.25
Respiratory Chain Com…eta Mitochondrial 402 -0.35 -0.55 -0.27 0.40 0.04 -0.50 -0.32
Respiratory Chain Com…it Mitochondrial 2929 -0.36 -0.58 -0.26 0.43 0.08 -0.56 -0.32
Respiratory Chain Com…it Mitochondrial 2914 -0.36 -0.56 -0.28 0.36 0.01 -0.50 -0.35
Pa700 Complex 32 -0.77 -0.54 -0.69 0.03 -0.40 -0.23 -0.75
Pa700 20s Pa28 Complex 193 -0.68 -0.38 -0.69 -0.32 -0.37 0.23 -0.70
Oligosaccharyltransferase Ostc Iii 1209 0.46 0.56 0.57 -0.13 -0.04 0.28 0.58
Oligosaccharyltransferase Ostc Ii 1208 0.40 0.52 0.51 -0.13 -0.16 0.27 0.50
Oligosaccharyltransferase Ostc I 1203 0.34 0.49 0.44 0.03 -0.10 0.22 0.53
1200 0.24 0.43 0.29 0.03 -0.07 0.29 0.48
1199 0.24 0.48 0.30 0.08 -0.09 0.38 0.51
8918 -0.35 -0.36 -0.31 -0.02 -0.21 -0.32 -0.35
Mitrac Complex Mitochondrial 8569 -0.71 -0.55 -0.80 -0.13 -0.08 -0.03 -0.69
Large Drosha Complex 1332 -0.86 -0.76 -0.79 0.04 -0.07 -0.17 -0.74
Kinase Maturation Complex 1 5199 -0.58 -0.34 -0.51 -0.58 -0.42 -0.02 -0.70
H2ax Complex II 1227 0.44 0.60 0.40 -0.65 -0.20 0.52 0.33
Emerin Complex 24 5611 0.06 -0.38 0.14 0.40 0.49 -0.64 -0.03
Dystrophin Sarcoglyca…x Skeletal Muscle 337 -0.40 -0.55 -0.31 0.21 0.02 -0.77 -0.45
Dystrophin Glycoprote…C Skeletal Muscle 340 -0.30 -0.42 -0.13 0.23 0.05 -0.75 -0.25
Ddb2 Complex 726 -0.73 -0.79 -0.73 -0.10 -0.19 -0.25 -0.90
Cct Complex Chaperoni…t Testis Specific 150 -0.51 -0.32 -0.48 -0.35 -0.41 -0.39 -0.47
C Src Immune Complex 6938 -0.33 -0.50 -0.43 0.11 0.26 -0.26 -0.22
Bbs Chaperonin Complex 6247 -0.48 -0.40 -0.42 -0.27 -0.31 -0.51 -0.46
26s Proteasome 181 -0.51 -0.25 -0.55 -0.40 -0.37 0.43 -0.59
Er Localized Multipro…f Ig Heavy Chains 414 0.41 0.77 0.39 0.16 0.08 0.54 0.73
Er Localized Multipro…Chains Associated 413 0.48 0.74 0.47 0.18 0.17 0.57 0.74
CSA
Capillary density
Satellite cells / fibre
Soleus pyruvate
Plantaris pyruvate
Soleus octanoyl
PAS vs vehicle
r
1.0
0.5
0.0
-0.5
-1.0



Female: Complexes, treatment effect and phenotype associations
Pearson r across this sex's ten mice (treatment: PAS = 1); box: FDR ≤ 0.05 (limma, BH within level)
Phenotype associations
Treatment effect
V Atpase Ragulator Ax…kb1 Ampk Complex 6888
Tim23 Sort Complex Ro…50 Mitochondrial 8723
Tim22 Holo Complex Ag…29 Mitochondrial 8557
Thymoproteasome 8522
Snw1 Complex 1335
Selk Multiprotein Complex 6131
Respiratory Chain Com…te Mitochondrial 2905
Pyruvate Dehydrogenase Complex 1375
Profilin 1 Complex 2836
Parvulin Associated Pre Rrnp Complex 3047
Pa28 20s Proteasome 192
Micos Complex Mitochondrial 6255
Lebercilin Complex 5899
Immunoproteasome 39
Hrd1 Complex 6859
Emerin Complex 52 5615
742
Dynactin 6130
Cytochrome C Oxidase Liver Mitochondrial 546
Cytochrome C Oxidase Liver Mitochondrial 538
7305
Cytochrome C Oxidase Heart Mitochondrial 542
Cytochrome Bc1 Comple…2 2 Mitochondrial 403
Coa3 Hspa9 Timm44 Tim…ex Mitochondrial 8563
51
C Complex Spliceosome 1181
Aqp2 Binding Multipro…Generator Complex 437
Alix Actomyosin Tight Junction Complex 8172
55s Ribosome Mitochondrial 320
39s Ribosomal Subunit Mitochondrial 324
28s Ribosomal Subunit Mitochondrial 315
20s Proteasome 38
0.08 0.19 -0.11 -0.43 0.08 0.24 0.12
-0.29 -0.19 -0.19 -0.31 -0.21 0.31 -0.33
0.00 -0.23 0.12 -0.48 -0.07 0.13 -0.35
-0.41 -0.18 -0.49 -0.50 -0.22 0.54 -0.48
-0.10 -0.24 -0.01 0.05 0.38 -0.71 0.02
-0.03 0.17 0.02 0.12 0.01 0.48 0.25
-0.08 -0.25 -0.07 0.22 0.02 -0.55 -0.10
0.51 0.32 0.46 0.10 0.42 0.37 0.43
0.27 0.17 0.09 0.17 0.28 -0.43 0.33
0.11 0.13 0.13 -0.40 -0.14 -0.26 0.01
-0.49 -0.20 -0.55 -0.47 -0.27 0.48 -0.52
0.29 0.18 0.46 -0.16 0.05 0.20 0.20
-0.14 -0.22 -0.24 -0.10 0.13 -0.07 -0.31
-0.39 -0.14 -0.47 -0.50 -0.26 0.52 -0.46
0.14 0.17 0.06 -0.07 0.20 0.70 0.24
-0.02 -0.22 -0.01 0.12 0.19 -0.44 -0.17
-0.76 -0.42 -0.82 0.06 -0.47 -0.05 -0.63
0.17 -0.14 0.12 0.49 0.35 -0.40 0.05
0.31 0.19 0.56 0.17 0.26 -0.06 0.41
0.31 0.20 0.56 0.15 0.24 -0.04 0.41
0.31 0.21 0.41 0.08 0.18 0.20 0.30
0.10 -0.18 0.39 0.12 0.40 -0.15 0.15
0.22 0.36 0.20 -0.26 -0.20 0.29 0.19
-0.42 -0.25 -0.30 -0.34 -0.24 0.24 -0.44
-0.42 -0.15 -0.50 -0.26 -0.47 -0.52 -0.35
-0.20 -0.38 -0.43 0.08 0.03 -0.16 -0.43
0.19 0.19 0.13 0.04 0.20 -0.62 0.29
0.30 0.16 0.25 0.31 0.26 -0.60 0.30
-0.34 -0.47 -0.48 0.41 -0.09 0.03 -0.48
-0.33 -0.44 -0.46 0.31 -0.12 -0.01 -0.53
-0.25 -0.39 -0.38 0.48 -0.02 0.09 -0.27
-0.41 -0.18 -0.48 -0.50 -0.25 0.53 -0.49
CSA
Capillary density
Satellite cells / fibre
Soleus pyruvate
Plantaris pyruvate
Soleus octanoyl
PAS vs vehicle
r
1.0
0.5
0.0
-0.5
-1.0


Female: Complexes, treatment effect and phenotype associations
Pearson r across this sex's ten mice (treatment: PAS = 1); box: FDR ≤ 0.05 (limma, BH within level)
Phenotype associations
Treatment effect
Zfand2b Complex 8497 -0.00 -0.04 -0.04 0.42 0.15 0.58 0.07
Vacuolar Atpase 442 0.10 0.13 -0.11 -0.57 0.07 0.40 0.02
5380 0.14 0.09 0.19 -0.35 -0.17 -0.17 -0.01
Tnf Alpha Nf Kappa B Signaling Complex 6 5266 -0.23 -0.11 -0.13 -0.47 -0.31 -0.25 -0.31
Tle1 Corepressor Complex 1729 0.32 0.11 0.29 -0.07 0.00 -0.30 0.02
Sdh Mabc1 Pic Ant Atpase Complex 139 -0.00 -0.11 0.21 0.07 -0.20 -0.18 -0.05
Ribosome Cytoplasmic 306 0.08 0.08 0.12 -0.36 -0.16 -0.17 -0.03
Respiratory Chain Com…bda Mitochondrial 401 0.22 -0.08 0.17 0.18 0.24 -0.39 0.08
Respiratory Chain Com…pha Mitochondrial 400 0.22 0.08 0.30 0.03 0.09 -0.25 0.18
Respiratory Chain Complex I Mitochondrial 382 0.09 -0.06 0.16 0.09 0.05 -0.32 0.06
Respiratory Chain Complex I Mitochondrial 381 0.12 -0.05 0.18 0.12 0.09 -0.34 0.09
Respiratory Chain Com…it Mitochondrial 2928 0.14 -0.07 0.05 -0.03 0.15 -0.13 -0.05
Respiratory Chain Com…it Mitochondrial 2920 0.16 -0.12 0.14 0.19 0.24 -0.33 0.05
Respiratory Chain Com…kd Mitochondrial 2904 0.05 -0.16 0.05 0.10 0.07 -0.48 -0.02
Respiratory Chain Com…te Mitochondrial 2948 -0.04 -0.26 -0.01 0.20 0.17 -0.48 -0.07
Respiratory Chain Com…yme Mitochondrial 178 0.02 -0.15 0.11 0.16 0.08 -0.34 -0.00
Respiratory Chain Com…it Mitochondrial 2919 0.12 0.14 0.27 -0.08 -0.07 -0.09 0.17
Respiratory Chain Com…ly Mitochondrial 2884 0.04 -0.08 0.21 0.32 0.08 -0.30 0.13
P2x7 Receptor Signalling Complex 725 0.01 -0.03 0.04 -0.08 -0.20 -0.56 -0.21
Nop56p Associated Pre rRNA Complex 3055 0.06 0.03 0.07 -0.35 -0.16 -0.23 -0.08
Mitofilin Complex Mitochondrial 8587 0.21 0.00 0.39 0.10 0.11 -0.22 0.21
Mib Complex Mitochondrial 6249 0.21 0.08 0.40 -0.02 0.07 -0.05 0.21
Frataxin Complex 1094 -0.13 0.01 -0.26 -0.28 -0.07 0.20 -0.08
7961 -0.03 0.01 -0.02 0.28 -0.12 0.42 0.05
7960 0.08 0.17 0.04 0.21 -0.15 0.53 0.15
Emerin Complex 25 5613 -0.01 -0.12 0.02 0.05 -0.03 -0.49 -0.22
Cdc5l Complex 1183 -0.17 -0.04 -0.15 -0.44 -0.35 -0.35 -0.36
Cct Prefoldin Complex Cct6b Variant 3074 -0.42 -0.15 -0.42 -0.59 -0.50 -0.19 -0.43
Cct Prefoldin Complex Cct6a Variant 145 -0.35 -0.02 -0.45 -0.46 -0.55 -0.36 -0.31
Actb Anp32a C1qbp Psm…ma Psma1 Complex 7298 0.04 0.08 -0.02 0.23 0.08 0.12 -0.01
60s Ribosomal Subunit Cytoplasmic 308 0.09 0.08 0.13 -0.36 -0.15 -0.21 -0.03
40s Ribosomal Subunit Cytoplasmic 305 0.06 0.09 0.11 -0.36 -0.19 -0.09 -0.03
CSA
Capillary density
Satellite cells / fibre
Soleus pyruvate
Plantaris pyruvate
Soleus octanoyl
PAS vs vehicle
r
1.0
0.5
0.0
-0.5
-1.0

Female: Complexes, treatment effect and phenotype associations


Pearson r across this sex's ten mice (treatment: PAS = 1); box: FDR ≤ 0.05 (limma, BH within level)




Female: Clusters, treatment effect and phenotype associations
Pearson r across this sex's ten mice (treatment: PAS = 1); box: FDR ≤ 0.05 (limma, BH within level)
Phenotype associations
Treatment effect
String 07 Unannotated
String 06 Myogenesis
String 05 Extracellular Matrix Organization
String 04 Translation
String 03 Hemostasis
String 02 Protein Secretion
String 01 Oxidative Phosphorylation
0.01 0.06 -0.05 0.29 -0.12 0.52 0.06
0.33 -0.00 0.37 0.68 0.37 -0.38 0.32
0.48 0.12 0.56 0.23 0.40 -0.47 0.41
-0.29 -0.25 -0.34 -0.30 -0.38 0.02 -0.56
-0.08 0.18 -0.20 -0.41 -0.07 0.03 0.06
-0.06 0.22 -0.24 -0.50 -0.31 0.54 -0.09
0.10 0.01 0.32 0.22 0.13 0.31 0.16
CSA
Capillary density
Satellite cells / fibre
Soleus pyruvate
Plantaris pyruvate
Soleus octanoyl
PAS vs vehicle
r
1.0
0.5
0.0
-0.5
-1.0


Female: strongest robust protein per phenotype

Smallest association p that keeps its sign and p ≤ 0.05 with any one mouse left out




Female: Proteins associated with Capillary density, p ≤ 0.05 in either model (n = 325)

Phenotype effect of PAS in this sex: +138% (p 0.0052).
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Female: Proteins associated with Capillary density, p ≤ 0.05 in either model (n = 325)

Phenotype effect of PAS in this sex: +138% (p 0.0052).
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Female: Proteins associated with Capillary density, p ≤ 0.05 in either model (n = 325)

Phenotype effect of PAS in this sex: +138% (p 0.0052).
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Female: Proteins associated with Capillary density, p ≤ 0.05 in either model (n = 325)

Phenotype effect of PAS in this sex: +138% (p 0.0052).
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Female: Proteins associated with Capillary density, p ≤ 0.05 in either model (n = 325)


Phenotype effect of PAS in this sex: +138% (p 0.0052).
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model



Female: Proteins associated with Capillary density, p ≤ 0.05 in either model (n = 325)


Phenotype effect of PAS in this sex: +138% (p 0.0052).
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Female: Proteins associated with Capillary density, p ≤ 0.05 in either model (n = 325)
Phenotype effect of PAS in this sex: +138% (p 0.0052).
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model
Qrsl1
r -0.45 (VEH 0.82, PAS -0.58)
p 0.043 pooled, 0.79 within
Magt1
r 0.49 (VEH 0.33, PAS 0.36)
p 0.043 pooled, 0.2 within
Micu3 (mito)
r -0.71 (VEH -0.82, PAS -0.82)
p 0.044 pooled, 0.17 within
Tsfm
r 0.31 (VEH 0.39, PAS 0.49)
p 0.044 pooled, 0.0034 within
Cand1
r -0.36 (VEH 0.44, PAS 0.45)
p 0.045 pooled, 0.45 within
Timm13 (mito)
r 0.69 (VEH 0.15, PAS 0.28)
p 0.045 pooled, 0.93 within
Cdc37
r -0.69 (VEH -0.45, PAS -0.08)
p 0.045 pooled, 0.66 within
Dglucy
r 0.54 (VEH 0.86, PAS 0.60)
p 0.045 pooled, 0.17 within
Aurkaip1 (mito)
r -0.53 (VEH -0.64, PAS -0.35)
p 0.045 pooled, 0.16 within
Cul5
r -0.56 (VEH 0.40, PAS -0.02)
p 0.046 pooled, 0.97 within
Cast
r 0.61 (VEH 0.59, PAS 0.74)
p 0.046 pooled, 0.17 within
Tmem41b
r -0.26 (VEH -0.93, PAS -0.10)
p 0.046 pooled, 0.21 within
Slc25a24
r 0.48 (VEH 0.57, PAS 0.92)
p 0.047 pooled, 0.18 within
Hacd1
r 0.42 (VEH 0.41, PAS 0.04)
p 0.047 pooled, 0.53 within
Acss1 (mito)
r -0.43 (VEH -0.33, PAS -0.51)
p 0.048 pooled, 0.23 within
Map2k4
r -0.32 (VEH -0.78, PAS -0.19)
p 0.048 pooled, 0.15 within
Mustn1
r 0.44 (VEH -0.28, PAS 0.48)
p 0.049 pooled, 0.027 within
Map4
r 0.51 (VEH 0.06, PAS 0.16)
p 0.049 pooled, 0.96 within
Slc44a2
r 0.59 (VEH -0.51, PAS -0.00)
p 0.049 pooled, 0.21 within
Ppt1
r 0.36 (VEH 0.71, PAS 0.64)
p 0.05 pooled, 0.0018 within
Feature value
Phenotype
VEH_F
PAS_F


Female: Proteins associated with Capillary density, p ≤ 0.05 in either model (n = 325)

Phenotype effect of PAS in this sex: +138% (p 0.0052).
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Female: Proteins associated with Capillary density, p ≤ 0.05 in either model (n = 325)

Phenotype effect of PAS in this sex: +138% (p 0.0052).
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Female: Proteins associated with Capillary density, p ≤ 0.05 in either model (n = 325)

Phenotype effect of PAS in this sex: +138% (p 0.0052).
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Female: Proteins associated with Capillary density, p ≤ 0.05 in either model (n = 325)

Phenotype effect of PAS in this sex: +138% (p 0.0052).
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Female: Proteins associated with Capillary density, p ≤ 0.05 in either model (n = 325)

Phenotype effect of PAS in this sex: +138% (p 0.0052).
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Female: Proteins associated with Capillary density, p ≤ 0.05 in either model (n = 325)

Phenotype effect of PAS in this sex: +138% (p 0.0052).
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Female: Proteins associated with Capillary density, p ≤ 0.05 in either model (n = 325)

Phenotype effect of PAS in this sex: +138% (p 0.0052).
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Female: Proteins associated with Capillary density, p ≤ 0.05 in either model (n = 325)

Phenotype effect of PAS in this sex: +138% (p 0.0052).
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Female: Proteins associated with Capillary density, p ≤ 0.05 in either model (n = 325)

Phenotype effect of PAS in this sex: +138% (p 0.0052).
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Female: Proteins associated with Capillary density, p ≤ 0.05 in either model (n = 325)

Phenotype effect of PAS in this sex: +138% (p 0.0052).
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Female: Proteins associated with CSA, p ≤ 0.05 in either model (n = 242)

Phenotype effect of PAS in this sex: +22% (p 0.002).
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Female: Proteins associated with CSA, p ≤ 0.05 in either model (n = 242)

Phenotype effect of PAS in this sex: +22% (p 0.002).
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Female: Proteins associated with CSA, p ≤ 0.05 in either model (n = 242)

Phenotype effect of PAS in this sex: +22% (p 0.002).
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Female: Proteins associated with CSA, p ≤ 0.05 in either model (n = 242)

Phenotype effect of PAS in this sex: +22% (p 0.002).
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Female: Proteins associated with CSA, p ≤ 0.05 in either model (n = 242)


Phenotype effect of PAS in this sex: +22% (p 0.002).
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model



Female: Proteins associated with CSA, p ≤ 0.05 in either model (n = 242)


Phenotype effect of PAS in this sex: +22% (p 0.002).
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Female: Proteins associated with CSA, p ≤ 0.05 in either model (n = 242)
Phenotype effect of PAS in this sex: +22% (p 0.002).
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model
Goltt1b
r 0.43 (VEH 0.30, PAS 0.58)
p 0.039 pooled, 0.35 within
Gpx1 (mito)
r 0.58 (VEH 0.29, PAS 0.18)
p 0.04 pooled, 0.39 within
Casq1
r 0.73 (VEH 0.42, PAS 0.16)
p 0.04 pooled, 0.73 within
Cdc37
r -0.69 (VEH -0.29, PAS 0.31)
p 0.04 pooled, 0.87 within
Mapk12
r -0.56 (VEH 0.08, PAS 0.35)
p 0.04 pooled, 0.4 within
Sntb2
r -0.70 (VEH -0.77, PAS -0.82)
p 0.04 pooled, 0.035 within
Myh7
r 0.33 (VEH -0.27, PAS 0.56)
p 0.041 pooled, 0.11 within
Gpx1
r 0.60 (VEH 0.26, PAS 0.26)
p 0.041 pooled, 0.39 within
Cycs (mito)
r 0.70 (VEH 0.58, PAS -0.53)
p 0.041 pooled, 0.55 within
Tmod4
r 0.41 (VEH -0.14, PAS 0.32)
p 0.042 pooled, 0.22 within
Mrpl47 (mito)
r -0.64 (VEH -0.30, PAS -0.75)
p 0.042 pooled, 0.035 within
Tln1
r -0.66 (VEH -0.32, PAS 0.14)
p 0.043 pooled, 0.96 within
Prune1
r -0.74 (VEH -0.58, PAS 0.83)
p 0.043 pooled, 1 within
Ctnnb1
r -0.51 (VEH -0.77, PAS -0.01)
p 0.044 pooled, 0.33 within
CD99
r -0.53 (VEH -0.07, PAS -0.06)
p 0.045 pooled, 0.93 within
Mbp
r -0.49 (VEH 0.09, PAS -0.91)
p 0.045 pooled, 0.13 within
Hnrnpd
r 0.61 (VEH 0.41, PAS 0.41)
p 0.046 pooled, 0.24 within
Mrpl47
r -0.60 (VEH -0.32, PAS -0.68)
p 0.046 pooled, 0.035 within
Acaa1a
r 0.71 (VEH -0.39, PAS 0.47)
p 0.046 pooled, 0.59 within
Myoz1
r 0.42 (VEH 0.03, PAS 0.31)
p 0.046 pooled, 0.25 within
Feature value
Phenotype
VEH_F
PAS_F


Female: Proteins associated with CSA, p ≤ 0.05 in either model (n = 242)

Phenotype effect of PAS in this sex: +22% (p 0.002).
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Female: Proteins associated with CSA, p ≤ 0.05 in either model (n = 242)

Phenotype effect of PAS in this sex: +22% (p 0.002).
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Female: Proteins associated with CSA, p ≤ 0.05 in either model (n = 242)

Phenotype effect of PAS in this sex: +22% (p 0.002).
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Female: Proteins associated with CSA, p ≤ 0.05 in either model (n = 242)

Phenotype effect of PAS in this sex: +22% (p 0.002).
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Female: Proteins associated with CSA, p ≤ 0.05 in either model (n = 242)

Phenotype effect of PAS in this sex: +22% (p 0.002).
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Female: Proteins associated with CSA, p ≤ 0.05 in either model (n = 242)

Phenotype effect of PAS in this sex: +22% (p 0.002).
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Female: Proteins associated with Satellite cells / fibre, p ≤ 0.05 in either model (n = 484)

Phenotype effect of PAS in this sex: +47% (p 0.0093).
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Female: Proteins associated with Satellite cells / fibre, p ≤ 0.05 in either model (n = 484)

Phenotype effect of PAS in this sex: +47% (p 0.0093).
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Female: Proteins associated with Satellite cells / fibre, p ≤ 0.05 in either model (n = 484)

Phenotype effect of PAS in this sex: +47% (p 0.0093).
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Female: Proteins associated with Satellite cells / fibre, p ≤ 0.05 in either model (n = 484)

Phenotype effect of PAS in this sex: +47% (p 0.0093).
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Female: Proteins associated with Satellite cells / fibre, p ≤ 0.05 in either model (n = 484)

Phenotype effect of PAS in this sex: +47% (p 0.0093).
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Female: Proteins associated with Satellite cells / fibre, p ≤ 0.05 in either model (n = 484)

Phenotype effect of PAS in this sex: +47% (p 0.0093).
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Female: Proteins associated with Satellite cells / fibre, p ≤ 0.05 in either model (n = 484)

Phenotype effect of PAS in this sex: +47% (p 0.0093).
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Female: Proteins associated with Satellite cells / fibre, p ≤ 0.05 in either model (n = 484)

Phenotype effect of PAS in this sex: +47% (p 0.0093).
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Female: Proteins associated with Satellite cells / fibre, p ≤ 0.05 in either model (n = 484)

Phenotype effect of PAS in this sex: +47% (p 0.0093).
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Female: Proteins associated with Satellite cells / fibre, p ≤ 0.05 in either model (n = 484)

Phenotype effect of PAS in this sex: +47% (p 0.0093).
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Female: Proteins associated with Satellite cells / fibre, p ≤ 0.05 in either model (n = 484)

Phenotype effect of PAS in this sex: +47% (p 0.0093).
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Female: Proteins associated with Satellite cells / fibre, p ≤ 0.05 in either model (n = 484)

Phenotype effect of PAS in this sex: +47% (p 0.0093).
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Female: Proteins associated with Satellite cells / fibre, p ≤ 0.05 in either model (n = 484)

Phenotype effect of PAS in this sex: +47% (p 0.0093).
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Female: Proteins associated with Satellite cells / fibre, p ≤ 0.05 in either model (n = 484)

Phenotype effect of PAS in this sex: +47% (p 0.0093).
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Female: Proteins associated with Satellite cells / fibre, p ≤ 0.05 in either model (n = 484)

Phenotype effect of PAS in this sex: +47% (p 0.0093).
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Female: Proteins associated with Satellite cells / fibre, p ≤ 0.05 in either model (n = 484)

Phenotype effect of PAS in this sex: +47% (p 0.0093).
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Female: Proteins associated with Satellite cells / fibre, p ≤ 0.05 in either model (n = 484)

Phenotype effect of PAS in this sex: +47% (p 0.0093).
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Female: Proteins associated with Satellite cells / fibre, p ≤ 0.05 in either model (n = 484)

Phenotype effect of PAS in this sex: +47% (p 0.0093).
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Female: Proteins associated with Satellite cells / fibre, p ≤ 0.05 in either model (n = 484)

Phenotype effect of PAS in this sex: +47% (p 0.0093).
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Female: Proteins associated with Satellite cells / fibre, p ≤ 0.05 in either model (n = 484)

Phenotype effect of PAS in this sex: +47% (p 0.0093).
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Female: Proteins associated with Satellite cells / fibre, p ≤ 0.05 in either model (n = 484)

Phenotype effect of PAS in this sex: +47% (p 0.0093).
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Female: Proteins associated with Satellite cells / fibre, p ≤ 0.05 in either model (n = 484)

Phenotype effect of PAS in this sex: +47% (p 0.0093).
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Female: Proteins associated with Satellite cells / fibre, p ≤ 0.05 in either model (n = 484)

Phenotype effect of PAS in this sex: +47% (p 0.0093).
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Female: Proteins associated with Satellite cells / fibre, p ≤ 0.05 in either model (n = 484)

Phenotype effect of PAS in this sex: +47% (p 0.0093).
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Female: Proteins associated with Satellite cells / fibre, p ≤ 0.05 in either model (n = 484)

Phenotype effect of PAS in this sex: +47% (p 0.0093).
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Female: Proteins associated with Soleus pyruvate, p ≤ 0.05 in either model (n = 125)

Phenotype effect of PAS in this sex: +12% (p 0.45), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Female: Proteins associated with Soleus pyruvate, p ≤ 0.05 in either model (n = 125)

Phenotype effect of PAS in this sex: +12% (p 0.45), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Female: Proteins associated with Soleus pyruvate, p ≤ 0.05 in either model (n = 125)

Phenotype effect of PAS in this sex: +12% (p 0.45), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Female: Proteins associated with Soleus pyruvate, p ≤ 0.05 in either model (n = 125)

Phenotype effect of PAS in this sex: +12% (p 0.45), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Female: Proteins associated with Soleus pyruvate, p ≤ 0.05 in either model (n = 125)

Phenotype effect of PAS in this sex: +12% (p 0.45), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Female: Proteins associated with Soleus pyruvate, p ≤ 0.05 in either model (n = 125)

Phenotype effect of PAS in this sex: +12% (p 0.45), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Female: Proteins associated with Soleus pyruvate, p ≤ 0.05 in either model (n = 125)

Phenotype effect of PAS in this sex: +12% (p 0.45), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Female: Proteins associated with Plantaris pyruvate, p ≤ 0.05 in either model (n = 85)

Phenotype effect of PAS in this sex: +31% (p 0.3), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Female: Proteins associated with Plantaris pyruvate, p ≤ 0.05 in either model (n = 85)

Phenotype effect of PAS in this sex: +31% (p 0.3), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Female: Proteins associated with Plantaris pyruvate, p ≤ 0.05 in either model (n = 85)

Phenotype effect of PAS in this sex: +31% (p 0.3), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Female: Proteins associated with Plantaris pyruvate, p ≤ 0.05 in either model (n = 85)

Phenotype effect of PAS in this sex: +31% (p 0.3), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Female: Proteins associated with Plantaris pyruvate, p ≤ 0.05 in either model (n = 85)

Phenotype effect of PAS in this sex: +31% (p 0.3), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Female: Proteins associated with Soleus octanoyl, p ≤ 0.05 in either model (n = 332)

Phenotype effect of PAS in this sex: +73% (p 0.35), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Female: Proteins associated with Soleus octanoyl, p ≤ 0.05 in either model (n = 332)

Phenotype effect of PAS in this sex: +73% (p 0.35), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Female: Proteins associated with Soleus octanoyl, p ≤ 0.05 in either model (n = 332)

Phenotype effect of PAS in this sex: +73% (p 0.35), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Female: Proteins associated with Soleus octanoyl, p ≤ 0.05 in either model (n = 332)

Phenotype effect of PAS in this sex: +73% (p 0.35), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Female: Proteins associated with Soleus octanoyl, p ≤ 0.05 in either model (n = 332)

Phenotype effect of PAS in this sex: +73% (p 0.35), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Female: Proteins associated with Soleus octanoyl, p ≤ 0.05 in either model (n = 332)

Phenotype effect of PAS in this sex: +73% (p 0.35), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Female: Proteins associated with Soleus octanoyl, p ≤ 0.05 in either model (n = 332)

Phenotype effect of PAS in this sex: +73% (p 0.35), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Female: Proteins associated with Soleus octanoyl, p ≤ 0.05 in either model (n = 332)

Phenotype effect of PAS in this sex: +73% (p 0.35), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Female: Proteins associated with Soleus octanoyl, p ≤ 0.05 in either model (n = 332)

Phenotype effect of PAS in this sex: +73% (p 0.35), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Female: Proteins associated with Soleus octanoyl, p ≤ 0.05 in either model (n = 332)

Phenotype effect of PAS in this sex: +73% (p 0.35), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Female: Proteins associated with Soleus octanoyl, p ≤ 0.05 in either model (n = 332)

Phenotype effect of PAS in this sex: +73% (p 0.35), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Female: Proteins associated with Soleus octanoyl, p ≤ 0.05 in either model (n = 332)

Phenotype effect of PAS in this sex: +73% (p 0.35), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Female: Proteins associated with Soleus octanoyl, p ≤ 0.05 in either model (n = 332)

Phenotype effect of PAS in this sex: +73% (p 0.35), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Female: Proteins associated with Soleus octanoyl, p ≤ 0.05 in either model (n = 332)

Phenotype effect of PAS in this sex: +73% (p 0.35), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Female: Proteins associated with Soleus octanoyl, p ≤ 0.05 in either model (n = 332)

Phenotype effect of PAS in this sex: +73% (p 0.35), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Female: Proteins associated with Soleus octanoyl, p ≤ 0.05 in either model (n = 332)

Phenotype effect of PAS in this sex: +73% (p 0.35), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Female: Proteins associated with Soleus octanoyl, p ≤ 0.05 in either model (n = 332)

Phenotype effect of PAS in this sex: +73% (p 0.35), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Female: Pathways associated with Capillary density, p ≤ 0.05 in either model (n = 35)

Phenotype effect of PAS in this sex: +138% (p 0.0052).
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Female: Pathways associated with Capillary density, p ≤ 0.05 in either model (n = 35)

Phenotype effect of PAS in this sex: +138% (p 0.0052).
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Female: Pathways associated with CSA, p ≤ 0.05 in either model (n = 45)

Phenotype effect of PAS in this sex: +22% (p 0.002).
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Female: Pathways associated with CSA, p ≤ 0.05 in either model (n = 45)

Phenotype effect of PAS in this sex: +22% (p 0.002).
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Female: Pathways associated with CSA, p ≤ 0.05 in either model (n = 45)

Phenotype effect of PAS in this sex: +22% (p 0.002).
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Female: Pathways associated with Satellite cells / fibre, p ≤ 0.05 in either model (n = 38)

Phenotype effect of PAS in this sex: +47% (p 0.0093).
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Female: Pathways associated with Satellite cells / fibre, p ≤ 0.05 in either model (n = 38)

Phenotype effect of PAS in this sex: +47% (p 0.0093).
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Female: Pathways associated with Soleus pyruvate, p ≤ 0.05 in either model (n = 91)

Phenotype effect of PAS in this sex: +12% (p 0.45), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Female: Pathways associated with Soleus pyruvate, p ≤ 0.05 in either model (n = 91)

Phenotype effect of PAS in this sex: +12% (p 0.45), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Female: Pathways associated with Soleus pyruvate, p ≤ 0.05 in either model (n = 91)

Phenotype effect of PAS in this sex: +12% (p 0.45), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Female: Pathways associated with Soleus pyruvate, p ≤ 0.05 in either model (n = 91)

Phenotype effect of PAS in this sex: +12% (p 0.45), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Female: Pathways associated with Soleus pyruvate, p ≤ 0.05 in either model (n = 91)

Phenotype effect of PAS in this sex: +12% (p 0.45), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Female: Pathways associated with Plantaris pyruvate, p ≤ 0.05 in either model (n = 13)

Phenotype effect of PAS in this sex: +31% (p 0.3), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Female: Pathways associated with Soleus octanoyl, p ≤ 0.05 in either model (n = 132)

Phenotype effect of PAS in this sex: +73% (p 0.35), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Female: Pathways associated with Soleus octanoyl, p ≤ 0.05 in either model (n = 132)

Phenotype effect of PAS in this sex: +73% (p 0.35), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Female: Pathways associated with Soleus octanoyl, p ≤ 0.05 in either model (n = 132)

Phenotype effect of PAS in this sex: +73% (p 0.35), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Female: Pathways associated with Soleus octanoyl, p ≤ 0.05 in either model (n = 132)

Phenotype effect of PAS in this sex: +73% (p 0.35), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Female: Pathways associated with Soleus octanoyl, p ≤ 0.05 in either model (n = 132)

Phenotype effect of PAS in this sex: +73% (p 0.35), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Female: Pathways associated with Soleus octanoyl, p ≤ 0.05 in either model (n = 132)

Phenotype effect of PAS in this sex: +73% (p 0.35), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Female: Pathways associated with Soleus octanoyl, p ≤ 0.05 in either model (n = 132)

Phenotype effect of PAS in this sex: +73% (p 0.35), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model



Female: Modules associated with Capillary density, p ≤ 0.05 in either model (n = 1)

Phenotype effect of PAS in this sex: +138% (p 0.0052).
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Feature value

Phenotype

VEH_F PAS_F


Female: Modules associated with CSA, p ≤ 0.05 in either model (n = 3)

Phenotype effect of PAS in this sex: +22% (p 0.002).
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Female: Modules associated with Satellite cells / fibre, p ≤ 0.05 in either model (n = 2)

Phenotype effect of PAS in this sex: +47% (p 0.0093).
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Female: Modules associated with Soleus octanoyl, p ≤ 0.05 in either model (n = 1)

Phenotype effect of PAS in this sex: +73% (p 0.35), so these associations follow variation between mice, not a PAS response. Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Female: Complexes associated with Capillary density, p ≤ 0.05 in either model (n = 10)

Phenotype effect of PAS in this sex: +138% (p 0.0052).
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Female: Complexes associated with CSA, p ≤ 0.05 in either model (n = 10)

Phenotype effect of PAS in this sex: +22% (p 0.002).
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Female: Complexes associated with Satellite cells / fibre, p ≤ 0.05 in either model (n = 13)

Phenotype effect of PAS in this sex: +47% (p 0.0093).
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Female: Complexes associated with Soleus pyruvate, p ≤ 0.05 in either model (n = 7)

Phenotype effect of PAS in this sex: +12% (p 0.45), so these associations follow variation between mice, not a PAS response. Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Female: Complexes associated with Soleus octanoyl, p ≤ 0.05 in either model (n = 17)

Phenotype effect of PAS in this sex: +73% (p 0.35), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Female: Clusters associated with Satellite cells / fibre, p ≤ 0.05 in either model (n = 1)

Phenotype effect of PAS in this sex: +47% (p 0.0093).
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model



Female: Clusters associated with Soleus pyruvate, p ≤ 0.05 in either model (n = 1)

Phenotype effect of PAS in this sex: +12% (p 0.45), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model



Female: Clusters associated with Soleus octanoyl, p ≤ 0.05 in either model (n = 1)

Phenotype effect of PAS in this sex: +73% (p 0.35), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model

