


Male: phenotype effects of PAS
% change against vehicle, Welch t on log values, n = 5 v 5
CSA
Capillary density
Satellite cells / fibre
Soleus pyruvate
Plantaris pyruvate
Soleus octanoyl
-50%
+0%
+100%
PAS vs vehicle, % change (95% CI, log scale)
Male
p > 0.05


Male: phenotype associations by level


This sex's ten mice, pooled across PAS and vehicle or within treatment; p ≤ 0.05 of tests; BH within level, phenotype and model




Check: protein associations with and without pellet mitochondrial share

Share centred within group, added to the association model






Male: Modules, treatment effect and phenotype associations
Pearson r across this sex's ten mice (treatment: PAS = 1); box: FDR ≤ 0.05 (limma, BH within level)
Phenotype associations
Treatment effect
Yellow -0.69 0.14 -0.16 -0.09 0.55 0.01 -0.76
Turquoise -0.57 0.27 0.29 -0.54 -0.16 0.02 -0.15
Tan -0.44 -0.31 -0.35 0.12 0.54 0.28 -0.50
Salmon 0.25 -0.57 -0.26 0.37 0.18 0.08 -0.12
Red -0.02 0.27 0.55 -0.26 -0.27 -0.19 0.41
Purple -0.38 -0.28 0.00 -0.22 0.42 0.25 -0.35
Pink -0.44 -0.30 -0.37 0.00 0.58 0.12 -0.77
Magenta -0.48 -0.16 -0.10 -0.01 0.31 0.25 -0.46
Greenyellow 0.20 0.08 -0.29 0.08 0.38 0.30 -0.05
Green 0.22 -0.35 -0.33 0.34 0.45 0.16 -0.15
Cyan -0.55 -0.29 0.12 -0.00 0.60 -0.06 -0.29
Brown -0.23 -0.24 -0.09 -0.05 0.37 0.19 -0.18
Blue 0.68 -0.19 -0.32 0.43 0.14 0.10 0.10
Black -0.05 0.16 0.19 -0.33 0.11 0.06 0.09
CSA
Capillary density
Satellite cells / fibre
Soleus pyruvate
Plantaris pyruvate
Soleus octanoyl
PAS vs vehicle
r
1.0
0.5
0.0
-0.5
-1.0



Male: Complexes, treatment effect and phenotype associations
Pearson r across this sex's ten mice (treatment: PAS = 1); box: FDR ≤ 0.05 (limma, BH within level)
Phenotype associations
Treatment effect
Zfand2b Complex 8497 0.39 0.29 -0.51 0.47 -0.34 -0.05 -0.48
Tnfalpha Nfkappa B Signaling Complex 5232 -0.09 -0.63 -0.03 0.31 -0.38 -0.29 -0.06
Spliceosome E Complex 8391 0.14 -0.56 -0.17 0.44 0.18 -0.28 -0.17
Spliceosome A Complex 8372 0.09 -0.59 -0.14 0.38 0.18 -0.30 -0.17
Snw1 Complex 1335 -0.37 -0.61 -0.06 0.02 0.18 -0.02 -0.11
Selk Multiprotein Complex 6131 0.07 0.69 0.10 -0.32 -0.33 0.12 0.22
Sdh Mabc1 Pic Ant Atpase Complex 139 0.02 0.38 0.20 -0.44 -0.61 0.25 0.31
Sarcoglycan Sarcospan…trobrevin Complex 349 -0.33 -0.25 -0.53 0.27 0.58 0.35 -0.65
Respiratory Chain Com…kd Mitochondrial 2904 -0.44 0.30 0.42 -0.77 -0.05 0.02 0.11
Respiratory Chain Com…te Mitochondrial 2905 -0.42 0.33 0.41 -0.76 -0.02 0.01 0.10
Respiratory Chain Com…it Mitochondrial 2919 -0.37 0.33 0.40 -0.71 -0.08 0.03 0.17
Respiratory Chain Com…ly Mitochondrial 2884 -0.30 0.38 0.40 -0.69 -0.14 -0.06 0.23
Parvulin Associated Pre Rrnp Complex 3047 -0.86 -0.54 0.28 -0.34 0.04 -0.06 -0.21
Oligosaccharyltransferase Ostc I 1203 0.25 0.50 0.17 -0.24 -0.44 0.10 0.52
1200 0.21 0.48 0.38 -0.28 -0.36 -0.04 0.58
1199 0.08 0.48 0.39 -0.35 -0.36 -0.02 0.53
Nop56p Associated Pre rRNA Complex 3055 -0.63 -0.68 0.22 -0.07 -0.14 -0.28 -0.17
Mib Complex Mitochondrial 6249 -0.46 0.39 0.26 -0.62 -0.33 0.08 -0.03
5685 0.54 0.31 -0.51 0.56 -0.37 -0.09 -0.32
Large Drosha Complex 1332 0.15 -0.63 -0.07 0.41 0.22 -0.29 -0.00
Hrd1 Complex 6859 0.52 0.65 0.07 -0.31 0.01 0.29 0.38
Frataxin Complex 1094 0.03 0.53 -0.43 -0.13 -0.35 0.66 -0.19
7961 -0.05 0.25 0.46 -0.14 -0.62 -0.35 0.47
7960 -0.05 0.26 0.47 -0.15 -0.61 -0.35 0.47
Emerin Complex 52 5615 0.52 -0.36 -0.15 0.30 0.27 0.02 0.09
Emerin Complex 25 5613 0.18 -0.49 -0.38 0.36 0.43 -0.01 -0.23
Emerin Complex 24 5611 0.16 -0.51 -0.35 0.54 0.30 -0.01 -0.14
742 0.00 -0.44 0.14 -0.29 0.15 -0.15 -0.17
Cdc5l Complex 1183 -0.08 -0.75 0.07 0.22 0.30 -0.29 -0.11
C Complex Spliceosome 1181 0.27 -0.50 -0.16 0.42 0.10 -0.22 -0.09
Er Localized Multipro…f Ig Heavy Chains 414 -0.08 0.40 0.24 -0.14 -0.40 -0.13 0.16
Er Localized Multipro…Chains Associated 413 0.20 0.46 0.24 -0.05 -0.50 -0.16 0.30
CSA
Capillary density
Satellite cells / fibre
Soleus pyruvate
Plantaris pyruvate
Soleus octanoyl
PAS vs vehicle
r
1.0
0.5
0.0
-0.5
-1.0


Male: Complexes, treatment effect and phenotype associations

Pearson r across this sex's ten mice (treatment: PAS = 1); box: FDR ≤ 0.05 (limma, BH within level)





Male: Complexes, treatment effect and phenotype associations
Pearson r across this sex's ten mice (treatment: PAS = 1); box: FDR ≤ 0.05 (limma, BH within level)
Phenotype associations
Treatment effect
Vacuolar Atpase 442 0.11 0.10 -0.29 -0.02 0.26 0.20 0.01
5380 -0.55 -0.31 0.24 -0.05 -0.02 -0.32 -0.20
Tle1 Corepressor Complex 1729 0.43 -0.30 -0.13 0.30 0.45 -0.17 -0.05
Tim22 Holo Complex Ag…29 Mitochondrial 8557 0.41 -0.07 -0.20 0.07 -0.56 0.12 0.32
Thymoproteasome 8522 -0.09 -0.10 0.40 -0.25 0.00 -0.15 0.52
Spermatoproteasome 8532 -0.22 -0.10 0.42 -0.30 0.08 -0.14 0.38
Sarcoglycan Sarcospan Complex Sg Spn 343 -0.33 -0.08 -0.45 0.15 0.47 0.45 -0.62
Respiratory Chain Com…it Mitochondrial 2928 -0.37 0.16 0.48 -0.85 -0.03 0.06 0.27
Profilin 1 Complex 2836 0.81 0.10 -0.18 0.30 -0.26 -0.05 0.45
Pa700 Complex 32 0.47 0.17 -0.41 0.57 -0.30 -0.08 -0.05
Pa28 20s Proteasome 192 0.06 -0.35 0.16 -0.09 0.13 0.06 0.56
P2x7 Receptor Signalling Complex 725 0.22 -0.04 -0.05 -0.01 0.17 0.14 0.17
8918 -0.67 0.13 0.43 -0.35 0.17 -0.07 0.03
Lebercilin Complex 5899 -0.42 -0.19 -0.01 -0.43 0.62 -0.04 -0.51
Immunoproteasome 39 0.21 -0.22 -0.01 0.20 0.11 0.07 0.50
Dystrophin Sarcoglyca…x Skeletal Muscle 337 -0.32 -0.15 -0.58 0.20 0.51 0.41 -0.71
Dynactin 6130 -0.43 0.16 -0.13 -0.47 0.59 0.05 -0.55
Ddb2 Complex 726 0.58 -0.26 -0.32 0.53 0.06 0.28 0.34
Cytochrome C Oxidase Liver Mitochondrial 546 -0.81 0.08 0.14 -0.46 -0.05 0.01 -0.44
Cytochrome C Oxidase Liver Mitochondrial 538 -0.81 0.08 0.14 -0.46 -0.05 0.01 -0.44
7305 -0.15 -0.14 -0.21 0.15 -0.18 -0.14 -0.36
Cytochrome C Oxidase Heart Mitochondrial 542 -0.61 0.23 -0.05 -0.60 -0.07 0.31 -0.44
Cct Prefoldin Complex Cct6b Variant 3074 -0.09 -0.11 -0.24 0.04 0.14 0.23 -0.38
Bbs Chaperonin Complex 6247 -0.66 -0.09 -0.03 -0.21 0.29 0.30 -0.42
Aqp2 Binding Multipro…Generator Complex 437 -0.26 0.14 -0.06 -0.29 0.35 0.17 -0.07
Alix Actomyosin Tight Junction Complex 8172 0.30 -0.12 -0.02 -0.01 0.31 -0.26 0.16
Actb Anp32a C1qbp Psm…ma Psma1 Complex 7298 0.74 -0.04 -0.36 0.33 0.08 0.23 0.36
55s Ribosome Mitochondrial 320 -0.20 0.09 0.22 -0.11 -0.27 -0.13 -0.02
39s Ribosomal Subunit Mitochondrial 324 -0.22 0.11 0.23 -0.09 -0.28 -0.20 -0.02
28s Ribosomal Subunit Mitochondrial 315 -0.18 0.05 0.20 -0.14 -0.25 -0.06 -0.01
26s Proteasome 181 0.10 -0.35 0.03 -0.20 -0.01 0.17 0.51
20s Proteasome 38 0.04 -0.30 0.19 -0.12 0.09 -0.00 0.55
CSA
Capillary density
Satellite cells / fibre
Soleus pyruvate
Plantaris pyruvate
Soleus octanoyl
PAS vs vehicle
r
1.0
0.5
0.0
-0.5
-1.0


Male: Complexes, treatment effect and phenotype associations
Pearson r across this sex's ten mice (treatment: PAS = 1); box: FDR ≤ 0.05 (limma, BH within level)
Phenotype associations
Treatment effect
Pa700 20s Pa28 Complex 193
Cct Complex Chaperoni…t Testis Specific 150
0.55
-0.03
-0.34
0.57
-0.24
-0.05
0.30
-0.45
-0.03
-0.21
-0.27
0.14
0.37
-0.46
r
1.0
0.5
0.0
-0.5
-1.0
CSA
Capillary density
Satellite cells / fibre
Soleus pyruvate
Plantaris pyruvate
Soleus octanoyl
PAS vs vehicle


Male: Clusters, treatment effect and phenotype associations
Pearson r across this sex's ten mice (treatment: PAS = 1); box: FDR ≤ 0.05 (limma, BH within level)
Phenotype associations
Treatment effect
String 07 Unannotated
String 06 Myogenesis
String 05 Extracellular Matrix Organization
String 04 Translation
String 03 Hemostasis
String 02 Protein Secretion
String 01 Oxidative Phosphorylation
0.05 0.26 0.41 -0.07 -0.61 -0.34 0.51
-0.52 0.41 -0.02 -0.16 0.15 -0.09 -0.45
-0.40 -0.17 -0.29 0.38 0.63 -0.13 -0.65
0.19 -0.39 0.00 0.30 -0.21 -0.24 0.06
0.12 -0.30 -0.22 0.24 0.56 0.15 -0.09
0.17 0.18 -0.28 0.00 -0.06 0.41 0.04
-0.16 0.54 0.32 -0.60 -0.49 0.03 0.21
CSA
Capillary density
Satellite cells / fibre
Soleus pyruvate
Plantaris pyruvate
Soleus octanoyl
PAS vs vehicle
r
1.0
0.5
0.0
-0.5
-1.0


Male: strongest robust protein per phenotype

Smallest association p that keeps its sign and p ≤ 0.05 with any one mouse left out




Male: Proteins associated with Capillary density, p ≤ 0.05 in either model (n = 605)

Phenotype effect of PAS in this sex: -18% (p 0.54), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Capillary density, p ≤ 0.05 in either model (n = 605)

Phenotype effect of PAS in this sex: -18% (p 0.54), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Capillary density, p ≤ 0.05 in either model (n = 605)

Phenotype effect of PAS in this sex: -18% (p 0.54), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Capillary density, p ≤ 0.05 in either model (n = 605)

Phenotype effect of PAS in this sex: -18% (p 0.54), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Capillary density, p ≤ 0.05 in either model (n = 605)

Phenotype effect of PAS in this sex: -18% (p 0.54), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Capillary density, p ≤ 0.05 in either model (n = 605)

Phenotype effect of PAS in this sex: -18% (p 0.54), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Capillary density, p ≤ 0.05 in either model (n = 605)

Phenotype effect of PAS in this sex: -18% (p 0.54), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Capillary density, p ≤ 0.05 in either model (n = 605)

Phenotype effect of PAS in this sex: -18% (p 0.54), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Capillary density, p ≤ 0.05 in either model (n = 605)

Phenotype effect of PAS in this sex: -18% (p 0.54), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Capillary density, p ≤ 0.05 in either model (n = 605)

Phenotype effect of PAS in this sex: -18% (p 0.54), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Capillary density, p ≤ 0.05 in either model (n = 605)

Phenotype effect of PAS in this sex: -18% (p 0.54), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Capillary density, p ≤ 0.05 in either model (n = 605)

Phenotype effect of PAS in this sex: -18% (p 0.54), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Capillary density, p ≤ 0.05 in either model (n = 605)

Phenotype effect of PAS in this sex: -18% (p 0.54), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Capillary density, p ≤ 0.05 in either model (n = 605)

Phenotype effect of PAS in this sex: -18% (p 0.54), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Capillary density, p ≤ 0.05 in either model (n = 605)

Phenotype effect of PAS in this sex: -18% (p 0.54), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Capillary density, p ≤ 0.05 in either model (n = 605)

Phenotype effect of PAS in this sex: -18% (p 0.54), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Capillary density, p ≤ 0.05 in either model (n = 605)

Phenotype effect of PAS in this sex: -18% (p 0.54), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Capillary density, p ≤ 0.05 in either model (n = 605)

Phenotype effect of PAS in this sex: -18% (p 0.54), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Capillary density, p ≤ 0.05 in either model (n = 605)

Phenotype effect of PAS in this sex: -18% (p 0.54), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Capillary density, p ≤ 0.05 in either model (n = 605)

Phenotype effect of PAS in this sex: -18% (p 0.54), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Capillary density, p ≤ 0.05 in either model (n = 605)

Phenotype effect of PAS in this sex: -18% (p 0.54), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Capillary density, p ≤ 0.05 in either model (n = 605)

Phenotype effect of PAS in this sex: -18% (p 0.54), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Capillary density, p ≤ 0.05 in either model (n = 605)

Phenotype effect of PAS in this sex: -18% (p 0.54), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Capillary density, p ≤ 0.05 in either model (n = 605)

Phenotype effect of PAS in this sex: -18% (p 0.54), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Capillary density, p ≤ 0.05 in either model (n = 605)

Phenotype effect of PAS in this sex: -18% (p 0.54), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Capillary density, p ≤ 0.05 in either model (n = 605)

Phenotype effect of PAS in this sex: -18% (p 0.54), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Capillary density, p ≤ 0.05 in either model (n = 605)

Phenotype effect of PAS in this sex: -18% (p 0.54), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Capillary density, p ≤ 0.05 in either model (n = 605)

Phenotype effect of PAS in this sex: -18% (p 0.54), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Capillary density, p ≤ 0.05 in either model (n = 605)

Phenotype effect of PAS in this sex: -18% (p 0.54), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Capillary density, p ≤ 0.05 in either model (n = 605)

Phenotype effect of PAS in this sex: -18% (p 0.54), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Capillary density, p ≤ 0.05 in either model (n = 605)

Phenotype effect of PAS in this sex: -18% (p 0.54), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with CSA, p ≤ 0.05 in either model (n = 818)

Phenotype effect of PAS in this sex: +13% (p 0.25), so these associations follow variation between mice, not a PAS response. Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with CSA, p ≤ 0.05 in either model (n = 818)

Phenotype effect of PAS in this sex: +13% (p 0.25), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with CSA, p ≤ 0.05 in either model (n = 818)

Phenotype effect of PAS in this sex: +13% (p 0.25), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with CSA, p ≤ 0.05 in either model (n = 818)

Phenotype effect of PAS in this sex: +13% (p 0.25), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with CSA, p ≤ 0.05 in either model (n = 818)

Phenotype effect of PAS in this sex: +13% (p 0.25), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with CSA, p ≤ 0.05 in either model (n = 818)

Phenotype effect of PAS in this sex: +13% (p 0.25), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with CSA, p ≤ 0.05 in either model (n = 818)

Phenotype effect of PAS in this sex: +13% (p 0.25), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with CSA, p ≤ 0.05 in either model (n = 818)

Phenotype effect of PAS in this sex: +13% (p 0.25), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with CSA, p ≤ 0.05 in either model (n = 818)

Phenotype effect of PAS in this sex: +13% (p 0.25), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with CSA, p ≤ 0.05 in either model (n = 818)

Phenotype effect of PAS in this sex: +13% (p 0.25), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with CSA, p ≤ 0.05 in either model (n = 818)

Phenotype effect of PAS in this sex: +13% (p 0.25), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with CSA, p ≤ 0.05 in either model (n = 818)

Phenotype effect of PAS in this sex: +13% (p 0.25), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with CSA, p ≤ 0.05 in either model (n = 818)

Phenotype effect of PAS in this sex: +13% (p 0.25), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with CSA, p ≤ 0.05 in either model (n = 818)

Phenotype effect of PAS in this sex: +13% (p 0.25), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with CSA, p ≤ 0.05 in either model (n = 818)

Phenotype effect of PAS in this sex: +13% (p 0.25), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with CSA, p ≤ 0.05 in either model (n = 818)

Phenotype effect of PAS in this sex: +13% (p 0.25), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with CSA, p ≤ 0.05 in either model (n = 818)

Phenotype effect of PAS in this sex: +13% (p 0.25), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with CSA, p ≤ 0.05 in either model (n = 818)

Phenotype effect of PAS in this sex: +13% (p 0.25), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with CSA, p ≤ 0.05 in either model (n = 818)

Phenotype effect of PAS in this sex: +13% (p 0.25), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with CSA, p ≤ 0.05 in either model (n = 818)

Phenotype effect of PAS in this sex: +13% (p 0.25), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with CSA, p ≤ 0.05 in either model (n = 818)


Phenotype effect of PAS in this sex: +13% (p 0.25), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model



Male: Proteins associated with CSA, p ≤ 0.05 in either model (n = 818)


Phenotype effect of PAS in this sex: +13% (p 0.25), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with CSA, p ≤ 0.05 in either model (n = 818)
Phenotype effect of PAS in this sex: +13% (p 0.25), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model
Nomo1
r -0.28 (VEH -0.25, PAS -0.66)
p 0.013 pooled, 0.0054 within
Atic
r 0.23 (VEH -0.60, PAS 0.46)
p 0.013 pooled, 0.024 within
CD55
r -0.79 (VEH -0.65, PAS -0.90)
p 0.013 pooled, 0.017 within
Ufc1
r -0.60 (VEH -0.47, PAS -0.90)
p 0.013 pooled, 0.0046 within
Etfdh
r -0.38 (VEH -0.52, PAS -0.40)
p 0.013 pooled, 0.011 within
Hspa9
r 0.74 (VEH 0.31, PAS 0.86)
p 0.013 pooled, 0.02 within
Samm50
r -0.52 (VEH -0.32, PAS -0.73)
p 0.013 pooled, 0.009 within
Ywhag
r -0.59 (VEH -0.49, PAS -0.90)
p 0.014 pooled, 0.0063 within
Nploc4
r 0.57 (VEH 0.83, PAS 0.71)
p 0.014 pooled, 0.0014 within
Nlrx1
r -0.54 (VEH -0.37, PAS -0.78)
p 0.014 pooled, 0.013 within
Dhrs1
r -0.33 (VEH -0.31, PAS -0.35)
p 0.014 pooled, 0.019 within
Pdia4
r -0.25 (VEH -0.17, PAS -0.25)
p 0.014 pooled, 0.04 within
Calu
r -0.75 (VEH -0.76, PAS -0.81)
p 0.014 pooled, 0.014 within
Atxn2l
r 0.48 (VEH 0.54, PAS 0.44)
p 0.015 pooled, 0.016 within
Eno1
r -0.77 (VEH -0.59, PAS -0.93)
p 0.015 pooled, 0.018 within
Eno3
r 0.55 (VEH 0.32, PAS 0.63)
p 0.015 pooled, 0.049 within
Aimp1
r 0.63 (VEH 0.64, PAS 0.69)
p 0.015 pooled, 0.013 within
Mrps34 (mito)
r -0.57 (VEH 0.16, PAS -0.84)
p 0.015 pooled, 0.017 within
Vtn
r 0.71 (VEH 0.57, PAS 0.77)
p 0.015 pooled, 0.0095 within
Lamp1
r -0.56 (VEH -0.74, PAS -0.49)
p 0.015 pooled, 0.014 within
Feature value
Phenotype
VEH_M
PAS_M


Male: Proteins associated with CSA, p ≤ 0.05 in either model (n = 818)

Phenotype effect of PAS in this sex: +13% (p 0.25), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with CSA, p ≤ 0.05 in either model (n = 818)

Phenotype effect of PAS in this sex: +13% (p 0.25), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with CSA, p ≤ 0.05 in either model (n = 818)

Phenotype effect of PAS in this sex: +13% (p 0.25), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with CSA, p ≤ 0.05 in either model (n = 818)

Phenotype effect of PAS in this sex: +13% (p 0.25), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with CSA, p ≤ 0.05 in either model (n = 818)


Phenotype effect of PAS in this sex: +13% (p 0.25), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with CSA, p ≤ 0.05 in either model (n = 818)

Phenotype effect of PAS in this sex: +13% (p 0.25), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model





Male: Proteins associated with CSA, p ≤ 0.05 in either model (n = 818)
Phenotype effect of PAS in this sex: +13% (p 0.25), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model
Opa1
r -0.49 (VEH -0.30, PAS -0.88)
p 0.029 pooled, 0.0067 within
Cdc42
r -0.50 (VEH -0.63, PAS -0.53)
p 0.029 pooled, 0.033 within
Coxta4 (mito)
r 0.73 (VEH 0.45, PAS 0.89)
p 0.029 pooled, 0.038 within
Rps21
r 0.59 (VEH 0.24, PAS 0.81)
p 0.029 pooled, 0.011 within
Nptn
r -0.50 (VEH 0.43, PAS -0.85)
p 0.029 pooled, 0.023 within
Cct5
r -0.66 (VEH -0.37, PAS -0.86)
p 0.029 pooled, 0.018 within
Lamc1
r -0.78 (VEH -0.87, PAS -0.71)
p 0.029 pooled, 0.04 within
Thnsl1 (mito)
r 0.57 (VEH 0.48, PAS 0.50)
p 0.029 pooled, 0.17 within
Cpt2 (mito)
r -0.52 (VEH -0.35, PAS -0.69)
p 0.03 pooled, 0.035 within
Got1
r -0.44 (VEH -0.58, PAS -0.47)
p 0.03 pooled, 0.014 within
Immt (mito)
r -0.71 (VEH -0.59, PAS -0.80)
p 0.03 pooled, 0.079 within
Prdx6
r -0.50 (VEH 0.42, PAS -0.71)
p 0.03 pooled, 0.085 within
Chid1
r 0.22 (VEH 0.31, PAS -0.05)
p 0.03 pooled, 0.11 within
Prdx6 (mito)
r -0.42 (VEH 0.60, PAS -0.75)
p 0.03 pooled, 0.11 within
Mpp1
r 0.29 (VEH -0.06, PAS 0.91)
p 0.03 pooled, 0.0016 within
Emc2
r -0.60 (VEH -0.26, PAS -0.90)
p 0.03 pooled, 0.0063 within
Myom1
r -0.72 (VEH -0.39, PAS -0.78)
p 0.03 pooled, 0.075 within
Vdac1 (mito)
r 0.71 (VEH 0.51, PAS 0.89)
p 0.03 pooled, 0.043 within
Mrpl17 (mito)
r -0.51 (VEH -0.36, PAS -0.82)
p 0.03 pooled, 0.032 within
Cat (mito)
r 0.39 (VEH 0.17, PAS 0.70)
p 0.03 pooled, 0.047 within
Feature value
Phenotype
VEH_M
PAS_M


Male: Proteins associated with CSA, p ≤ 0.05 in either model (n = 818)
Phenotype effect of PAS in this sex: +13% (p 0.25), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model
Mrps7
r -0.65 (VEH -0.61, PAS -0.69)
p 0.03 pooled, 0.011 within
Ndufa2 (mito)
r 0.75 (VEH 0.92, PAS 0.67)
p 0.03 pooled, 0.053 within
Tars2 (mito)
r -0.58 (VEH -0.16, PAS -0.91)
p 0.031 pooled, 0.035 within
Higd1a (mito)
r 0.63 (VEH 0.24, PAS 0.81)
p 0.031 pooled, 0.052 within
Dtd1
r 0.42 (VEH 0.65, PAS 0.04)
p 0.031 pooled, 0.072 within
Ppox
r -0.62 (VEH -0.54, PAS -0.71)
p 0.031 pooled, 0.0097 within
Mrps11 (mito)
r 0.24 (VEH 0.40, PAS 0.12)
p 0.031 pooled, 0.066 within
Pcbp1
r -0.50 (VEH -0.55, PAS -0.28)
p 0.031 pooled, 0.057 within
Ank1
r 0.42 (VEH NA, PAS 0.33)
p 0.031 pooled, 0.038 within
Crat (mito)
r -0.65 (VEH -0.24, PAS -0.90)
p 0.031 pooled, 0.04 within
Fundc2 (mito)
r -0.36 (VEH -0.21, PAS -0.73)
p 0.032 pooled, 0.019 within
Rheb
r -0.56 (VEH -0.22, PAS -0.71)
p 0.032 pooled, 0.044 within
Cox7c
r -0.68 (VEH -0.65, PAS -0.72)
p 0.032 pooled, 0.039 within
Ssr3
r -0.08 (VEH -0.40, PAS -0.08)
p 0.032 pooled, 0.021 within
Ndufa5 (mito)
r 0.31 (VEH 0.27, PAS 0.13)
p 0.032 pooled, 0.12 within
Stim1
r 0.44 (VEH 0.55, PAS 0.27)
p 0.032 pooled, 0.07 within
Dcakd
r -0.31 (VEH -0.17, PAS -0.45)
p 0.032 pooled, 0.039 within
Snx3
r 0.34 (VEH 0.21, PAS 0.64)
p 0.032 pooled, 0.023 within
Tpm4
r 0.29 (VEH -0.30, PAS 0.82)
p 0.033 pooled, 0.058 within
Capns1
r -0.46 (VEH -0.34, PAS -0.51)
p 0.033 pooled, 0.048 within
Feature value
Phenotype
VEH_M
PAS_M


Male: Proteins associated with CSA, p ≤ 0.05 in either model (n = 818)

Phenotype effect of PAS in this sex: +13% (p 0.25), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with CSA, p ≤ 0.05 in either model (n = 818)

Phenotype effect of PAS in this sex: +13% (p 0.25), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with CSA, p ≤ 0.05 in either model (n = 818)


Phenotype effect of PAS in this sex: +13% (p 0.25), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model



Male: Proteins associated with CSA, p ≤ 0.05 in either model (n = 818)


Phenotype effect of PAS in this sex: +13% (p 0.25), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with CSA, p ≤ 0.05 in either model (n = 818)
Phenotype effect of PAS in this sex: +13% (p 0.25), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model
Gpc1
r -0.52 (VEH 0.52, PAS -0.77)
p 0.044 pooled, 0.077 within
Aldoart2
r 0.48 (VEH 0.20, PAS 0.60)
p 0.045 pooled, 0.045 within
Tmem65
r -0.56 (VEH -0.69, PAS -0.61)
p 0.045 pooled, 0.023 within
Lrrc38
r -0.52 (VEH -0.27, PAS -0.81)
p 0.045 pooled, 0.012 within
Atp5pb
r -0.44 (VEH -0.39, PAS -0.48)
p 0.045 pooled, 0.043 within
Mrpl57
r 0.11 (VEH 0.30, PAS 0.15)
p 0.045 pooled, 0.017 within
Tnnc2
r -0.74 (VEH -0.75, PAS -0.75)
p 0.045 pooled, 0.16 within
Timm22
r -0.26 (VEH -0.35, PAS -0.27)
p 0.046 pooled, 0.044 within
Sec61b
r 0.85 (VEH 0.77, PAS 0.89)
p 0.046 pooled, 0.085 within
Bola1
r 0.37 (VEH 0.04, PAS 0.72)
p 0.046 pooled, 0.027 within
Serpina1b
r -0.62 (VEH -0.38, PAS -0.71)
p 0.046 pooled, 0.055 within
Ndufb8
r -0.32 (VEH -0.51, PAS -0.22)
p 0.046 pooled, 0.039 within
Cyc1
r -0.43 (VEH -0.46, PAS -0.46)
p 0.047 pooled, 0.056 within
Letm1 (mito)
r -0.26 (VEH -0.30, PAS -0.27)
p 0.047 pooled, 0.089 within
Chmp6
r -0.61 (VEH -0.37, PAS -0.71)
p 0.047 pooled, 0.047 within
Svip
r 0.24 (VEH 0.45, PAS 0.37)
p 0.047 pooled, 0.016 within
Atp5po
r -0.39 (VEH -0.43, PAS -0.54)
p 0.047 pooled, 0.035 within
Ndufa10
r -0.35 (VEH -0.49, PAS -0.41)
p 0.047 pooled, 0.03 within
Bak1 (mito)
r 0.72 (VEH 0.97, PAS 0.53)
p 0.047 pooled, 0.11 within
Aurkaip1 (mito)
r 0.40 (VEH 0.71, PAS 0.03)
p 0.047 pooled, 0.081 within
Feature value
Phenotype
VEH_M
PAS_M


Male: Proteins associated with CSA, p ≤ 0.05 in either model (n = 818)

Phenotype effect of PAS in this sex: +13% (p 0.25), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with CSA, p ≤ 0.05 in either model (n = 818)

Phenotype effect of PAS in this sex: +13% (p 0.25), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with CSA, p ≤ 0.05 in either model (n = 818)

Phenotype effect of PAS in this sex: +13% (p 0.25), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with CSA, p ≤ 0.05 in either model (n = 818)

Phenotype effect of PAS in this sex: +13% (p 0.25), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with CSA, p ≤ 0.05 in either model (n = 818)

Phenotype effect of PAS in this sex: +13% (p 0.25), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Satellite cells / fibre, p ≤ 0.05 in either model (n = 895)

Phenotype effect of PAS in this sex: +43% (p 0.086), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Satellite cells / fibre, p ≤ 0.05 in either model (n = 895)

Phenotype effect of PAS in this sex: +43% (p 0.086), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Satellite cells / fibre, p ≤ 0.05 in either model (n = 895)

Phenotype effect of PAS in this sex: +43% (p 0.086), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Satellite cells / fibre, p ≤ 0.05 in either model (n = 895)

Phenotype effect of PAS in this sex: +43% (p 0.086), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Satellite cells / fibre, p ≤ 0.05 in either model (n = 895)

Phenotype effect of PAS in this sex: +43% (p 0.086), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Satellite cells / fibre, p ≤ 0.05 in either model (n = 895)

Phenotype effect of PAS in this sex: +43% (p 0.086), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Satellite cells / fibre, p ≤ 0.05 in either model (n = 895)

Phenotype effect of PAS in this sex: +43% (p 0.086), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Satellite cells / fibre, p ≤ 0.05 in either model (n = 895)

Phenotype effect of PAS in this sex: +43% (p 0.086), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Satellite cells / fibre, p ≤ 0.05 in either model (n = 895)

Phenotype effect of PAS in this sex: +43% (p 0.086), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Satellite cells / fibre, p ≤ 0.05 in either model (n = 895)

Phenotype effect of PAS in this sex: +43% (p 0.086), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Satellite cells / fibre, p ≤ 0.05 in either model (n = 895)

Phenotype effect of PAS in this sex: +43% (p 0.086), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Satellite cells / fibre, p ≤ 0.05 in either model (n = 895)

Phenotype effect of PAS in this sex: +43% (p 0.086), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Satellite cells / fibre, p ≤ 0.05 in either model (n = 895)

Phenotype effect of PAS in this sex: +43% (p 0.086), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Satellite cells / fibre, p ≤ 0.05 in either model (n = 895)

Phenotype effect of PAS in this sex: +43% (p 0.086), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Satellite cells / fibre, p ≤ 0.05 in either model (n = 895)

Phenotype effect of PAS in this sex: +43% (p 0.086), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Satellite cells / fibre, p ≤ 0.05 in either model (n = 895)

Phenotype effect of PAS in this sex: +43% (p 0.086), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Satellite cells / fibre, p ≤ 0.05 in either model (n = 895)

Phenotype effect of PAS in this sex: +43% (p 0.086), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Satellite cells / fibre, p ≤ 0.05 in either model (n = 895)

Phenotype effect of PAS in this sex: +43% (p 0.086), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Satellite cells / fibre, p ≤ 0.05 in either model (n = 895)


Phenotype effect of PAS in this sex: +43% (p 0.086), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model



Male: Proteins associated with Satellite cells / fibre, p ≤ 0.05 in either model (n = 895)


Phenotype effect of PAS in this sex: +43% (p 0.086), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model





Male: Proteins associated with Satellite cells / fibre, p ≤ 0.05 in either model (n = 895)
Phenotype effect of PAS in this sex: +43% (p 0.086), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model
Pdhb (mito)
r 0.31 (VEH 0.88, PAS -0.23)
p 0.02 pooled, 0.013 within
Naca
r -0.41 (VEH -0.86, PAS 0.27)
p 0.02 pooled, 0.042 within
Anxa11
r 0.41 (VEH 0.91, PAS -0.58)
p 0.02 pooled, 0.03 within
Hnrnpa1
r -0.11 (VEH -0.86, PAS 0.45)
p 0.02 pooled, 0.017 within
Supv3l1 (mito)
r -0.42 (VEH -0.95, PAS -0.09)
p 0.02 pooled, 0.01 within
Mrpl51
r 0.15 (VEH 0.21, PAS 0.13)
p 0.02 pooled, 0.089 within
Diablo (mito)
r -0.30 (VEH -0.85, PAS -0.33)
p 0.02 pooled, 0.00086 within
Rplp0
r 0.23 (VEH 0.75, PAS -0.05)
p 0.02 pooled, 0.007 within
Sucla2
r 0.24 (VEH 0.82, PAS 0.29)
p 0.02 pooled, 0.00096 within
Atp5mk
r 0.20 (VEH 0.93, PAS -0.37)
p 0.02 pooled, 0.011 within
Dpysl2
r -0.36 (VEH -0.27, PAS -0.30)
p 0.02 pooled, 0.058 within
Dld (mito)
r -0.28 (VEH -0.94, PAS 0.37)
p 0.02 pooled, 0.0044 within
Slc25a12 (mito)
r 0.14 (VEH 0.76, PAS -0.48)
p 0.02 pooled, 0.023 within
Slc25a3 (mito)
r 0.20 (VEH 0.90, PAS -0.22)
p 0.02 pooled, 0.0032 within
Ywhah
r 0.32 (VEH 0.87, PAS -0.87)
p 0.02 pooled, 0.72 within
Agl
r 0.56 (VEH 0.97, PAS 0.55)
p 0.02 pooled, 0.0033 within
Acsl1
r 0.22 (VEH 0.93, PAS -0.61)
p 0.021 pooled, 0.014 within
Mtfr1l (mito)
r -0.29 (VEH -0.80, PAS -0.19)
p 0.021 pooled, 0.006 within
Uqcrh (mito)
r -0.30 (VEH -0.83, PAS -0.33)
p 0.021 pooled, 0.0012 within
Apool (mito)
r 0.22 (VEH 0.84, PAS -0.33)
p 0.021 pooled, 0.013 within
Feature value
Phenotype
VEH_M
PAS_M

Male: Proteins associated with Satellite cells / fibre, p ≤ 0.05 in either model (n = 895)


Phenotype effect of PAS in this sex: +43% (p 0.086), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Satellite cells / fibre, p ≤ 0.05 in either model (n = 895)
Phenotype effect of PAS in this sex: +43% (p 0.086), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model
Lias (mito)
r -0.31 (VEH -0.85, PAS -0.17)
p 0.023 pooled, 0.0055 within
Mrpl1
r 0.40 (VEH 0.85, PAS 0.05)
p 0.023 pooled, 0.22 within
Prdx3 (mito)
r -0.36 (VEH -0.82, PAS -0.49)
p 0.023 pooled, 0.00094 within
Mrpl46 (mito)
r -0.21 (VEH -0.86, PAS 0.14)
p 0.023 pooled, 0.0021 within
Chchd7
r -0.27 (VEH -0.70, PAS -0.08)
p 0.023 pooled, 0.043 within
Gfm1 (mito)
r 0.56 (VEH 1.00, PAS -0.07)
p 0.023 pooled, 0.048 within
Pcyt1a
r -0.39 (VEH -0.51, PAS -0.22)
p 0.023 pooled, 0.12 within
Timm23 (mito)
r 0.44 (VEH 0.70, PAS 0.38)
p 0.023 pooled, 0.0091 within
Dmac1
r 0.11 (VEH 0.72, PAS -0.67)
p 0.023 pooled, 0.098 within
Ndufa6
r 0.23 (VEH 0.91, PAS -0.27)
p 0.024 pooled, 0.0041 within
Rpl23a
r -0.18 (VEH -0.85, PAS 0.63)
p 0.024 pooled, 0.11 within
Isca2 (mito)
r -0.15 (VEH -0.57, PAS 0.44)
p 0.024 pooled, 0.052 within
Adhfe1
r 0.35 (VEH 0.87, PAS -0.01)
p 0.024 pooled, 0.035 within
Cavin4
r -0.34 (VEH -0.64, PAS 0.27)
p 0.024 pooled, 0.056 within
Chchd2
r -0.26 (VEH -0.78, PAS -0.27)
p 0.024 pooled, 0.0028 within
Tefm (mito)
r 0.43 (VEH 0.60, PAS 0.17)
p 0.024 pooled, 0.084 within
Rab12
r 0.19 (VEH 0.87, PAS -0.81)
p 0.024 pooled, 0.13 within
Lonp1 (mito)
r 0.38 (VEH 0.83, PAS 0.05)
p 0.024 pooled, 0.017 within
Tbc1d17
r 0.40 (VEH 0.82, PAS -0.08)
p 0.025 pooled, 0.038 within
Camk2g
r 0.47 (VEH 0.74, PAS 0.23)
p 0.025 pooled, 0.027 within
Feature value
Phenotype
VEH_M
PAS_M


Male: Proteins associated with Satellite cells / fibre, p ≤ 0.05 in either model (n = 895)

Phenotype effect of PAS in this sex: +43% (p 0.086), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Satellite cells / fibre, p ≤ 0.05 in either model (n = 895)

Phenotype effect of PAS in this sex: +43% (p 0.086), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Satellite cells / fibre, p ≤ 0.05 in either model (n = 895)

Phenotype effect of PAS in this sex: +43% (p 0.086), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Satellite cells / fibre, p ≤ 0.05 in either model (n = 895)

Phenotype effect of PAS in this sex: +43% (p 0.086), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Satellite cells / fibre, p ≤ 0.05 in either model (n = 895)


Phenotype effect of PAS in this sex: +43% (p 0.086), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model



Male: Proteins associated with Satellite cells / fibre, p ≤ 0.05 in either model (n = 895)


Phenotype effect of PAS in this sex: +43% (p 0.086), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Satellite cells / fibre, p ≤ 0.05 in either model (n = 895)
Phenotype effect of PAS in this sex: +43% (p 0.086), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model
Ldhb (mito)
r 0.42 (VEH 0.72, PAS -0.04)
p 0.042 pooled, 0.06 within
Ndufb8
r 0.17 (VEH 0.95, PAS -0.40)
p 0.042 pooled, 0.024 within
Trap1 (mito)
r 0.17 (VEH 0.80, PAS -0.07)
p 0.042 pooled, 0.0097 within
Pa2g4
r -0.06 (VEH -0.72, PAS 0.41)
p 0.042 pooled, 0.051 within
Mrpl13 (mito)
r -0.41 (VEH -0.73, PAS -0.02)
p 0.043 pooled, 0.051 within
Rps3
r 0.25 (VEH 0.81, PAS 0.13)
p 0.043 pooled, 0.0063 within
Aspscr1
r -0.06 (VEH -0.51, PAS 0.54)
p 0.043 pooled, 0.27 within
Atp5f1d
r -0.28 (VEH -0.85, PAS -0.02)
p 0.043 pooled, 0.036 within
Ech1 (mito)
r -0.28 (VEH -0.82, PAS 0.15)
p 0.043 pooled, 0.099 within
Tmed10
r 0.27 (VEH 0.85, PAS -0.21)
p 0.043 pooled, 0.037 within
Psma5
r 0.16 (VEH 0.78, PAS -0.40)
p 0.043 pooled, 0.067 within
Maob
r 0.11 (VEH 0.86, PAS -0.41)
p 0.043 pooled, 0.032 within
Rpl26
r -0.26 (VEH -0.79, PAS 0.39)
p 0.044 pooled, 0.024 within
Slc25a46
r -0.29 (VEH -0.98, PAS -0.23)
p 0.044 pooled, 0.0077 within
Aqp1
r 0.15 (VEH 0.91, PAS 0.09)
p 0.044 pooled, 0.0011 within
Tpd52l2
r -0.52 (VEH -0.51, PAS -0.25)
p 0.044 pooled, 0.1 within
Prdx3
r -0.34 (VEH -0.69, PAS -0.58)
p 0.044 pooled, 0.0051 within
Mrpl51 (mito)
r 0.12 (VEH 0.06, PAS 0.15)
p 0.045 pooled, 0.13 within
Letm1
r 0.26 (VEH 0.83, PAS -0.18)
p 0.045 pooled, 0.016 within
r 0.35 (VEH 0.78, PAS 0.10)
p 0.045 pooled, 0.044 within
Feature value
Phenotype
VEH_M
PAS_M


Male: Proteins associated with Satellite cells / fibre, p ≤ 0.05 in either model (n = 895)

Phenotype effect of PAS in this sex: +43% (p 0.086), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Satellite cells / fibre, p ≤ 0.05 in either model (n = 895)

Phenotype effect of PAS in this sex: +43% (p 0.086), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Satellite cells / fibre, p ≤ 0.05 in either model (n = 895)

Phenotype effect of PAS in this sex: +43% (p 0.086), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Satellite cells / fibre, p ≤ 0.05 in either model (n = 895)

Phenotype effect of PAS in this sex: +43% (p 0.086), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Satellite cells / fibre, p ≤ 0.05 in either model (n = 895)

Phenotype effect of PAS in this sex: +43% (p 0.086), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Satellite cells / fibre, p ≤ 0.05 in either model (n = 895)

Phenotype effect of PAS in this sex: +43% (p 0.086), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Satellite cells / fibre, p ≤ 0.05 in either model (n = 895)

Phenotype effect of PAS in this sex: +43% (p 0.086), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Satellite cells / fibre, p ≤ 0.05 in either model (n = 895)

Phenotype effect of PAS in this sex: +43% (p 0.086), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Satellite cells / fibre, p ≤ 0.05 in either model (n = 895)

Phenotype effect of PAS in this sex: +43% (p 0.086), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Satellite cells / fibre, p ≤ 0.05 in either model (n = 895)

Phenotype effect of PAS in this sex: +43% (p 0.086), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Satellite cells / fibre, p ≤ 0.05 in either model (n = 895)

Phenotype effect of PAS in this sex: +43% (p 0.086), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Satellite cells / fibre, p ≤ 0.05 in either model (n = 895)

Phenotype effect of PAS in this sex: +43% (p 0.086), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Satellite cells / fibre, p ≤ 0.05 in either model (n = 895)

Phenotype effect of PAS in this sex: +43% (p 0.086), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Satellite cells / fibre, p ≤ 0.05 in either model (n = 895)

Phenotype effect of PAS in this sex: +43% (p 0.086), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Satellite cells / fibre, p ≤ 0.05 in either model (n = 895)

Phenotype effect of PAS in this sex: +43% (p 0.086), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Soleus pyruvate, p ≤ 0.05 in either model (n = 658)

Phenotype effect of PAS in this sex: +9% (p 0.77), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Soleus pyruvate, p ≤ 0.05 in either model (n = 658)

Phenotype effect of PAS in this sex: +9% (p 0.77), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Soleus pyruvate, p ≤ 0.05 in either model (n = 658)

Phenotype effect of PAS in this sex: +9% (p 0.77), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Soleus pyruvate, p ≤ 0.05 in either model (n = 658)

Phenotype effect of PAS in this sex: +9% (p 0.77), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Soleus pyruvate, p ≤ 0.05 in either model (n = 658)

Phenotype effect of PAS in this sex: +9% (p 0.77), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Soleus pyruvate, p ≤ 0.05 in either model (n = 658)

Phenotype effect of PAS in this sex: +9% (p 0.77), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Soleus pyruvate, p ≤ 0.05 in either model (n = 658)

Phenotype effect of PAS in this sex: +9% (p 0.77), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Soleus pyruvate, p ≤ 0.05 in either model (n = 658)

Phenotype effect of PAS in this sex: +9% (p 0.77), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Soleus pyruvate, p ≤ 0.05 in either model (n = 658)

Phenotype effect of PAS in this sex: +9% (p 0.77), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Soleus pyruvate, p ≤ 0.05 in either model (n = 658)

Phenotype effect of PAS in this sex: +9% (p 0.77), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Soleus pyruvate, p ≤ 0.05 in either model (n = 658)

Phenotype effect of PAS in this sex: +9% (p 0.77), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Soleus pyruvate, p ≤ 0.05 in either model (n = 658)

Phenotype effect of PAS in this sex: +9% (p 0.77), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Soleus pyruvate, p ≤ 0.05 in either model (n = 658)

Phenotype effect of PAS in this sex: +9% (p 0.77), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Soleus pyruvate, p ≤ 0.05 in either model (n = 658)

Phenotype effect of PAS in this sex: +9% (p 0.77), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Soleus pyruvate, p ≤ 0.05 in either model (n = 658)

Phenotype effect of PAS in this sex: +9% (p 0.77), so these associations follow variation between mice, not a PAS response. Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Soleus pyruvate, p ≤ 0.05 in either model (n = 658)

Phenotype effect of PAS in this sex: +9% (p 0.77), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Soleus pyruvate, p ≤ 0.05 in either model (n = 658)

Phenotype effect of PAS in this sex: +9% (p 0.77), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Soleus pyruvate, p ≤ 0.05 in either model (n = 658)

Phenotype effect of PAS in this sex: +9% (p 0.77), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Soleus pyruvate, p ≤ 0.05 in either model (n = 658)

Phenotype effect of PAS in this sex: +9% (p 0.77), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Soleus pyruvate, p ≤ 0.05 in either model (n = 658)

Phenotype effect of PAS in this sex: +9% (p 0.77), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Soleus pyruvate, p ≤ 0.05 in either model (n = 658)

Phenotype effect of PAS in this sex: +9% (p 0.77), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Soleus pyruvate, p ≤ 0.05 in either model (n = 658)

Phenotype effect of PAS in this sex: +9% (p 0.77), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Soleus pyruvate, p ≤ 0.05 in either model (n = 658)

Phenotype effect of PAS in this sex: +9% (p 0.77), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Soleus pyruvate, p ≤ 0.05 in either model (n = 658)

Phenotype effect of PAS in this sex: +9% (p 0.77), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Soleus pyruvate, p ≤ 0.05 in either model (n = 658)

Phenotype effect of PAS in this sex: +9% (p 0.77), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Soleus pyruvate, p ≤ 0.05 in either model (n = 658)

Phenotype effect of PAS in this sex: +9% (p 0.77), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Soleus pyruvate, p ≤ 0.05 in either model (n = 658)

Phenotype effect of PAS in this sex: +9% (p 0.77), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Soleus pyruvate, p ≤ 0.05 in either model (n = 658)

Phenotype effect of PAS in this sex: +9% (p 0.77), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Soleus pyruvate, p ≤ 0.05 in either model (n = 658)


Phenotype effect of PAS in this sex: +9% (p 0.77), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model



Male: Proteins associated with Soleus pyruvate, p ≤ 0.05 in either model (n = 658)


Phenotype effect of PAS in this sex: +9% (p 0.77), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Soleus pyruvate, p ≤ 0.05 in either model (n = 658)
Phenotype effect of PAS in this sex: +9% (p 0.77), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model
Cdv3
r 0.45 (VEH 0.75, PAS -0.33)
p 0.041 pooled, 0.086 within
Mcu (mito)
r -0.42 (VEH -0.93, PAS 0.16)
p 0.041 pooled, 0.18 within
Tmed9
r -0.32 (VEH -0.62, PAS 0.44)
p 0.041 pooled, 0.13 within
Crat (mito)
r -0.48 (VEH -0.89, PAS 0.03)
p 0.041 pooled, 0.093 within
Eif4g1
r 0.36 (VEH 0.84, PAS -0.38)
p 0.042 pooled, 0.12 within
Rab10
r -0.47 (VEH -0.70, PAS 0.16)
p 0.042 pooled, 0.13 within
Perm1
r 0.43 (VEH 0.84, PAS -0.11)
p 0.042 pooled, 0.096 within
Dld (mito)
r 0.47 (VEH 0.76, PAS -0.57)
p 0.042 pooled, 0.2 within
Pnpla7
r -0.36 (VEH -0.57, PAS 0.32)
p 0.042 pooled, 0.13 within
Rhot2
r -0.51 (VEH -0.67, PAS -0.08)
p 0.042 pooled, 0.1 within
Gnai2
r -0.43 (VEH -0.65, PAS -0.18)
p 0.042 pooled, 0.11 within
Gpd1
r -0.35 (VEH -0.82, PAS 0.39)
p 0.042 pooled, 0.13 within
Bdh1 (mito)
r -0.49 (VEH -0.65, PAS -0.20)
p 0.043 pooled, 0.12 within
Ssbp1 (mito)
r 0.48 (VEH 0.84, PAS 0.11)
p 0.043 pooled, 0.12 within
Ces1d
r -0.58 (VEH -0.81, PAS -0.40)
p 0.043 pooled, 0.1 within
Ethe1
r 0.39 (VEH 0.83, PAS -0.25)
p 0.043 pooled, 0.12 within
Plin3
r 0.45 (VEH 0.57, PAS 0.29)
p 0.043 pooled, 0.1 within
Mrpl14
r 0.44 (VEH 0.59, PAS 0.12)
p 0.043 pooled, 0.12 within
Gadd45gip1
r 0.50 (VEH 0.79, PAS -0.35)
p 0.043 pooled, 0.12 within
Hsd17b4
r -0.63 (VEH -0.68, PAS -0.83)
p 0.044 pooled, 0.071 within
Feature value
Phenotype
VEH_M
PAS_M


Male: Proteins associated with Soleus pyruvate, p ≤ 0.05 in either model (n = 658)

Phenotype effect of PAS in this sex: +9% (p 0.77), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Soleus pyruvate, p ≤ 0.05 in either model (n = 658)

Phenotype effect of PAS in this sex: +9% (p 0.77), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Plantaris pyruvate, p ≤ 0.05 in either model (n = 678)

Phenotype effect of PAS in this sex: -23% (p 0.46), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Plantaris pyruvate, p ≤ 0.05 in either model (n = 678)

Phenotype effect of PAS in this sex: -23% (p 0.46), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Plantaris pyruvate, p ≤ 0.05 in either model (n = 678)

Phenotype effect of PAS in this sex: -23% (p 0.46), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Plantaris pyruvate, p ≤ 0.05 in either model (n = 678)

Phenotype effect of PAS in this sex: -23% (p 0.46), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Plantaris pyruvate, p ≤ 0.05 in either model (n = 678)

Phenotype effect of PAS in this sex: -23% (p 0.46), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Plantaris pyruvate, p ≤ 0.05 in either model (n = 678)

Phenotype effect of PAS in this sex: -23% (p 0.46), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Plantaris pyruvate, p ≤ 0.05 in either model (n = 678)


Phenotype effect of PAS in this sex: -23% (p 0.46), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model



Male: Proteins associated with Plantaris pyruvate, p ≤ 0.05 in either model (n = 678)


Phenotype effect of PAS in this sex: -23% (p 0.46), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Plantaris pyruvate, p ≤ 0.05 in either model (n = 678)
Phenotype effect of PAS in this sex: -23% (p 0.46), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model
B3galnt2
r -0.49 (VEH -0.65, PAS -0.30)
p 0.0056 pooled, 0.037 within
Flnc
r 0.76 (VEH 0.85, PAS 0.32)
p 0.0057 pooled, 0.011 within
Mrpl51
r -0.31 (VEH -0.19, PAS -0.52)
p 0.0058 pooled, 0.0033 within
Mfn2
r -0.30 (VEH -0.12, PAS -0.87)
p 0.0058 pooled, 0.0057 within
Bzw2
r -0.27 (VEH -0.07, PAS -0.75)
p 0.0058 pooled, 0.018 within
Myo1c
r 0.63 (VEH 0.72, PAS 0.15)
p 0.0059 pooled, 0.053 within
Acp6
r -0.34 (VEH -0.21, PAS -0.79)
p 0.006 pooled, 0.00054 within
Mrps27
r -0.44 (VEH -0.77, PAS -0.28)
p 0.006 pooled, 0.0078 within
Crip1
r 0.58 (VEH 0.60, PAS 0.47)
p 0.0061 pooled, 0.0059 within
Epb41l2
r 0.66 (VEH 0.95, PAS 0.33)
p 0.0062 pooled, 0.025 within
Sar1b
r -0.26 (VEH -0.21, PAS -0.58)
p 0.0062 pooled, 0.0016 within
Golga2
r 0.62 (VEH 0.88, PAS 0.55)
p 0.0063 pooled, 0.085 within
Lrpprc
r -0.35 (VEH -0.27, PAS -0.73)
p 0.0063 pooled, 0.0024 within
Prdx2 (mito)
r 0.39 (VEH 0.48, PAS 0.57)
p 0.0064 pooled, 0.003 within
Nefh
r 0.58 (VEH 0.57, PAS 0.33)
p 0.0064 pooled, 0.12 within
Cox5a (mito)
r 0.23 (VEH 0.04, PAS 0.59)
p 0.0065 pooled, 0.028 within
Mbp
r 0.66 (VEH 0.68, PAS 0.58)
p 0.0065 pooled, 0.071 within
Mlip
r 0.46 (VEH 0.53, PAS 0.02)
p 0.0065 pooled, 0.031 within
Flna
r 0.54 (VEH 0.64, PAS 0.32)
p 0.0066 pooled, 0.015 within
Acp6 (mito)
r -0.34 (VEH -0.23, PAS -0.73)
p 0.0066 pooled, 0.0072 within
Feature value
Phenotype
VEH_M
PAS_M


Male: Proteins associated with Plantaris pyruvate, p ≤ 0.05 in either model (n = 678)

Phenotype effect of PAS in this sex: -23% (p 0.46), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Plantaris pyruvate, p ≤ 0.05 in either model (n = 678)

Phenotype effect of PAS in this sex: -23% (p 0.46), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Plantaris pyruvate, p ≤ 0.05 in either model (n = 678)

Phenotype effect of PAS in this sex: -23% (p 0.46), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Plantaris pyruvate, p ≤ 0.05 in either model (n = 678)

Phenotype effect of PAS in this sex: -23% (p 0.46), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Plantaris pyruvate, p ≤ 0.05 in either model (n = 678)

Phenotype effect of PAS in this sex: -23% (p 0.46), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Plantaris pyruvate, p ≤ 0.05 in either model (n = 678)

Phenotype effect of PAS in this sex: -23% (p 0.46), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Plantaris pyruvate, p ≤ 0.05 in either model (n = 678)

Phenotype effect of PAS in this sex: -23% (p 0.46), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Plantaris pyruvate, p ≤ 0.05 in either model (n = 678)

Phenotype effect of PAS in this sex: -23% (p 0.46), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Plantaris pyruvate, p ≤ 0.05 in either model (n = 678)


Phenotype effect of PAS in this sex: -23% (p 0.46), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model



Male: Proteins associated with Plantaris pyruvate, p ≤ 0.05 in either model (n = 678)


Phenotype effect of PAS in this sex: -23% (p 0.46), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Plantaris pyruvate, p ≤ 0.05 in either model (n = 678)
Phenotype effect of PAS in this sex: -23% (p 0.46), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model
Nfu1 (mito)
r 0.41 (VEH 0.38, PAS 0.45)
p 0.026 pooled, 0.029 within
Slc25a20
r -0.27 (VEH -0.17, PAS -0.75)
p 0.026 pooled, 0.0081 within
Ryr1
r -0.33 (VEH -0.07, PAS -0.66)
p 0.026 pooled, 0.098 within
Clptm1
r -0.42 (VEH -0.53, PAS 0.19)
p 0.026 pooled, 0.12 within
Capzb
r 0.60 (VEH 0.66, PAS 0.43)
p 0.026 pooled, 0.23 within
Vdac2
r -0.21 (VEH -0.05, PAS -0.78)
p 0.026 pooled, 0.011 within
S100a10
r 0.57 (VEH 0.54, PAS 0.40)
p 0.027 pooled, 0.2 within
Msra (mito)
r 0.35 (VEH 0.03, PAS 0.94)
p 0.027 pooled, 0.071 within
Hspa5
r -0.24 (VEH -0.15, PAS -0.20)
p 0.027 pooled, 0.18 within
Htra2 (mito)
r -0.33 (VEH -0.39, PAS -0.94)
p 0.027 pooled, 0.0055 within
Tom1
r 0.32 (VEH 0.29, PAS 0.40)
p 0.027 pooled, 0.012 within
Mrpl27
r -0.26 (VEH -0.17, PAS -0.57)
p 0.027 pooled, 0.0048 within
Rbm3
r 0.32 (VEH 0.40, PAS 0.39)
p 0.027 pooled, 0.015 within
Ssr4
r -0.20 (VEH 0.09, PAS -0.64)
p 0.028 pooled, 0.076 within
Nt5dc3
r -0.44 (VEH -0.55, PAS -0.97)
p 0.028 pooled, 0.00075 within
Myadm
r 0.54 (VEH 0.65, PAS 0.17)
p 0.028 pooled, 0.13 within
Myorg
r -0.47 (VEH -0.25, PAS -0.63)
p 0.028 pooled, 0.39 within
Slc25a12
r -0.38 (VEH -0.26, PAS -0.62)
p 0.028 pooled, 0.01 within
Cs (mito)
r -0.58 (VEH -0.97, PAS -0.07)
p 0.028 pooled, 0.029 within
Des
r 0.56 (VEH 0.90, PAS -0.09)
p 0.028 pooled, 0.073 within
Feature value
Phenotype
VEH_M
PAS_M


Male: Proteins associated with Plantaris pyruvate, p ≤ 0.05 in either model (n = 678)

Phenotype effect of PAS in this sex: -23% (p 0.46), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Plantaris pyruvate, p ≤ 0.05 in either model (n = 678)

Phenotype effect of PAS in this sex: -23% (p 0.46), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Plantaris pyruvate, p ≤ 0.05 in either model (n = 678)


Phenotype effect of PAS in this sex: -23% (p 0.46), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model



Male: Proteins associated with Plantaris pyruvate, p ≤ 0.05 in either model (n = 678)


Phenotype effect of PAS in this sex: -23% (p 0.46), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Plantaris pyruvate, p ≤ 0.05 in either model (n = 678)
Phenotype effect of PAS in this sex: -23% (p 0.46), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model
Sudg2
r -0.19 (VEH -0.22, PAS -0.48)
p 0.04 pooled, 0.0041 within
Rilp
r -0.31 (VEH -0.29, PAS -0.97)
p 0.04 pooled, 0.0009 within
Cdh13
r 0.37 (VEH 0.42, PAS 0.10)
p 0.041 pooled, 0.1 within
Mrpl46 (mito)
r 0.10 (VEH -0.02, PAS 0.50)
p 0.041 pooled, 0.031 within
Plpp7
r -0.43 (VEH -0.34, PAS -0.28)
p 0.041 pooled, 0.56 within
Rack1
r -0.11 (VEH 0.07, PAS -0.56)
p 0.041 pooled, 0.038 within
Atp5mg (mito)
r -0.20 (VEH -0.08, PAS -0.56)
p 0.042 pooled, 0.032 within
B2m
r 0.47 (VEH 0.62, PAS 0.13)
p 0.042 pooled, 0.05 within
Mrps18a
r -0.32 (VEH -0.37, PAS -0.47)
p 0.042 pooled, 0.0049 within
Cpox (mito)
r 0.33 (VEH 0.01, PAS 0.80)
p 0.043 pooled, 0.21 within
Hspa8
r -0.13 (VEH 0.13, PAS -0.83)
p 0.043 pooled, 0.048 within
Slc25a12 (mito)
r -0.38 (VEH -0.32, PAS -0.44)
p 0.043 pooled, 0.037 within
Eloc
r -0.49 (VEH NA, PAS -0.75)
p 0.043 pooled, 0.37 within
Mrpl11 (mito)
r 0.26 (VEH 0.06, PAS 0.76)
p 0.043 pooled, 0.047 within
Rhot2
r -0.33 (VEH -0.14, PAS -0.79)
p 0.043 pooled, 0.029 within
Grsf1
r 0.31 (VEH 0.85, PAS -0.11)
p 0.043 pooled, 0.0018 within
Prph
r 0.51 (VEH 0.50, PAS 0.33)
p 0.043 pooled, 0.44 within
March5 (mito)
r -0.59 (VEH -0.79, PAS -0.41)
p 0.043 pooled, 0.053 within
Pin4
r 0.20 (VEH 0.25, PAS 0.44)
p 0.043 pooled, 0.0029 within
Vwa8 (mito)
r -0.22 (VEH -0.12, PAS -0.33)
p 0.043 pooled, 0.09 within
Feature value
Phenotype
VEH_M
PAS_M


Male: Proteins associated with Plantaris pyruvate, p ≤ 0.05 in either model (n = 678)

Phenotype effect of PAS in this sex: -23% (p 0.46), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Plantaris pyruvate, p ≤ 0.05 in either model (n = 678)

Phenotype effect of PAS in this sex: -23% (p 0.46), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Plantaris pyruvate, p ≤ 0.05 in either model (n = 678)

Phenotype effect of PAS in this sex: -23% (p 0.46), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Plantaris pyruvate, p ≤ 0.05 in either model (n = 678)

Phenotype effect of PAS in this sex: -23% (p 0.46), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Plantaris pyruvate, p ≤ 0.05 in either model (n = 678)

Phenotype effect of PAS in this sex: -23% (p 0.46), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Plantaris pyruvate, p ≤ 0.05 in either model (n = 678)

Phenotype effect of PAS in this sex: -23% (p 0.46), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Plantaris pyruvate, p ≤ 0.05 in either model (n = 678)

Phenotype effect of PAS in this sex: -23% (p 0.46), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Plantaris pyruvate, p ≤ 0.05 in either model (n = 678)

Phenotype effect of PAS in this sex: -23% (p 0.46), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Plantaris pyruvate, p ≤ 0.05 in either model (n = 678)

Phenotype effect of PAS in this sex: -23% (p 0.46), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Soleus octanoyl, p ≤ 0.05 in either model (n = 47)

Phenotype effect of PAS in this sex: -16% (p 0.57), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Soleus octanoyl, p ≤ 0.05 in either model (n = 47)

Phenotype effect of PAS in this sex: -16% (p 0.57), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Proteins associated with Soleus octanoyl, p ≤ 0.05 in either model (n = 47)

Phenotype effect of PAS in this sex: -16% (p 0.57), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Pathways associated with Capillary density, p ≤ 0.05 in either model (n = 20)

Phenotype effect of PAS in this sex: -18% (p 0.54), so these associations follow variation between mice, not a PAS response. Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Pathways associated with CSA, p ≤ 0.05 in either model (n = 45)

Phenotype effect of PAS in this sex: +13% (p 0.25), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Pathways associated with CSA, p ≤ 0.05 in either model (n = 45)

Phenotype effect of PAS in this sex: +13% (p 0.25), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Pathways associated with CSA, p ≤ 0.05 in either model (n = 45)

Phenotype effect of PAS in this sex: +13% (p 0.25), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Pathways associated with Satellite cells / fibre, p ≤ 0.05 in either model (n = 34)

Phenotype effect of PAS in this sex: +43% (p 0.086), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Pathways associated with Satellite cells / fibre, p ≤ 0.05 in either model (n = 34)

Phenotype effect of PAS in this sex: +43% (p 0.086), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Pathways associated with Soleus pyruvate, p ≤ 0.05 in either model (n = 24)

Phenotype effect of PAS in this sex: +9% (p 0.77), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Pathways associated with Soleus pyruvate, p ≤ 0.05 in either model (n = 24)

Phenotype effect of PAS in this sex: +9% (p 0.77), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Pathways associated with Plantaris pyruvate, p ≤ 0.05 in either model (n = 113)

Phenotype effect of PAS in this sex: -23% (p 0.46), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Pathways associated with Plantaris pyruvate, p ≤ 0.05 in either model (n = 113)

Phenotype effect of PAS in this sex: -23% (p 0.46), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Pathways associated with Plantaris pyruvate, p ≤ 0.05 in either model (n = 113)

Phenotype effect of PAS in this sex: -23% (p 0.46), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Pathways associated with Plantaris pyruvate, p ≤ 0.05 in either model (n = 113)

Phenotype effect of PAS in this sex: -23% (p 0.46), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Pathways associated with Plantaris pyruvate, p ≤ 0.05 in either model (n = 113)

Phenotype effect of PAS in this sex: -23% (p 0.46), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Pathways associated with Plantaris pyruvate, p ≤ 0.05 in either model (n = 113)

Phenotype effect of PAS in this sex: -23% (p 0.46), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Pathways associated with Soleus octanoyl, p ≤ 0.05 in either model (n = 1)

Phenotype effect of PAS in this sex: -16% (p 0.57), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model



Male: Modules associated with Capillary density, p ≤ 0.05 in either model (n = 1)

Phenotype effect of PAS in this sex: -18% (p 0.54), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Feature value

Phenotype

VEH_M PAS_M


Male: Modules associated with CSA, p ≤ 0.05 in either model (n = 2)

Phenotype effect of PAS in this sex: +13% (p 0.25), so these associations follow variation between mice, not a PAS response. Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model



Male: Modules associated with Satellite cells / fibre, p ≤ 0.05 in either model (n = 1)

Phenotype effect of PAS in this sex: +43% (p 0.086), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Feature value

Phenotype

VEH_M PAS_M


Male: Modules associated with Soleus pyruvate, p ≤ 0.05 in either model (n = 1)

Phenotype effect of PAS in this sex: +9% (p 0.77), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Modules associated with Plantaris pyruvate, p ≤ 0.05 in either model (n = 1)

Phenotype effect of PAS in this sex: -23% (p 0.46), so these associations follow variation between mice, not a PAS response. Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Complexes associated with Capillary density, p ≤ 0.05 in either model (n = 10)

Phenotype effect of PAS in this sex: -18% (p 0.54), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Complexes associated with CSA, p ≤ 0.05 in either model (n = 9)

Phenotype effect of PAS in this sex: +13% (p 0.25), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Complexes associated with Satellite cells / fibre, p ≤ 0.05 in either model (n = 2)

Phenotype effect of PAS in this sex: +43% (p 0.086), so these associations follow variation between mice, not a PAS response. Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model




Male: Complexes associated with Soleus pyruvate, p ≤ 0.05 in either model (n = 4)

Phenotype effect of PAS in this sex: +9% (p 0.77), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model



Male: Complexes associated with Plantaris pyruvate, p ≤ 0.05 in either model (n = 3)

Phenotype effect of PAS in this sex: -23% (p 0.46), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model



Male: Clusters associated with Plantaris pyruvate, p ≤ 0.05 in either model (n = 1)

Phenotype effect of PAS in this sex: -23% (p 0.46), so these associations follow variation between mice, not a PAS response.
Lines within vehicle and within PAS (5 mice each), the within-treatment model; dashed: all 10 mice, the pooled model

