


Sex difference: Treatment × sex, every protein

limpa dpcDE, BH within universe; labelled: FDR ≤ 0.05 and the 10 smallest p





Proteins per tier
Dashed: proteins at p ≤ 0.05 expected by chance in each direction; Π can include p > 0.05
Full proteome
Mitochondrial universe
FDR ≤ 0.05
Π ≤ 0.05
p ≤ 0.05
0
0
43
133
36
36
40
109
46
46
50
0
50
100
150
60
30
0
30
60
Proteins (down | up)
down
up


All 149 proteins at p ≤ 0.05, per mouse
Rows z-scored across these mice, clustered on 1 - Pearson r (display only)
VEH_F
PAS_F
VEH_M
PAS_M
z
2
1
0
-1
-2
VEH_F1
VEH_F2
VEH_F3
VEH_F4
VEH_F5
PAS_F1
PAS_F2
PAS_F3
PAS_F4
PAS_F5
VEH_M1
VEH_M2
VEH_M3
VEH_M4
VEH_M5
PAS_M1
PAS_M2
PAS_M3
PAS_M4
PAS_M5


Sex difference: every protein at p ≤ 0.05, in order of p

log2 abundance per mouse, bar at the group median; log2FC: female PAS response minus male PAS response




Sex difference: every protein at p ≤ 0.05, in order of p

log2 abundance per mouse, bar at the group median; log2FC: female PAS response minus male PAS response




Sex difference: every protein at p ≤ 0.05, in order of p

log2 abundance per mouse, bar at the group median; log2FC: female PAS response minus male PAS response




Sex difference: every protein at p ≤ 0.05, in order of p

log2 abundance per mouse, bar at the group median; log2FC: female PAS response minus male PAS response




Sex difference: every protein at p ≤ 0.05, in order of p

log2 abundance per mouse, bar at the group median; log2FC: female PAS response minus male PAS response




Sex difference: every protein at p ≤ 0.05, in order of p

log2 abundance per mouse, bar at the group median; log2FC: female PAS response minus male PAS response



Sex difference: every protein at p ≤ 0.05, in order of p

log2 abundance per mouse, bar at the group median; log2FC: female PAS response minus male PAS response



Sex difference: every protein at p ≤ 0.05, in order of p

log2 abundance per mouse, bar at the group median; log2FC: female PAS response minus male PAS response

