

Female: GO:BP

*12 strongest terms the EnrichmentMap flag keeps; * FDR≤0.05*




Female: Reactome


*12 strongest terms the EnrichmentMap flag keeps; * FDR≤0.05*





Female: MitoCarta

*12 strongest terms the EnrichmentMap flag keeps; * FDR≤0.05*





Female: signal by GO Slim family
fgsea; a term counts in every family it belongs to; every family with a term at p ≤ 0.05
Anatomical structure development
Muscle system process
Signaling
Cytoskeleton organization
Cell differentiation
Nervous system process
Mitotic cell cycle
Carbohydrate metabolic process
Telomere organization
Protein catabolic process
Programmed cell death
Detoxification
Transmembrane transport
Protein-containing complex assembly
Mitochondrion organization
Lipid metabolic process
Immune system process
Generation of precursor metabolites and energy
Carbohydrate derivative metabolic process
Amino acid metabolic process
0
5
GO:BP terms at p ≤ 0.05 (down | up)
down
up



20 strongest GO:BP and Reactome terms, ordered by shared genes
EnrichmentMap similarity, average linkage; black ring FDR ≤ 0.05
Regulation Of Growth
Regulation Of Chemotaxis
Regulation Of Cell Cycle
Cell Cycle
Microtubule Cytoskeleton Organization
Cell Cycle Mitotic
Protein Modification Process
Proteasome Mediated Ubiquitin Dependent Protein Catabolic Process
Protein Targeting
Multicellular Organismal Movement
Skeletal Muscle Contraction
Neuromuscular Process
Striated Muscle Contraction
Muscle Contraction
Striated Muscle Contraction
Sarcomere Organization
Myofibril Assembly
Cellular Response To Toxic Substance
Cellular Homeostasis
Small Molecule Catabolic Process
-2
-1
0
1
2
NES
NES
3
2
1
0
-1
-2
-3
-log10(p)
3.0
3.5
4.0
4.5


GO Slim cellular compartments
fgsea, full proteome
Endoplasmic Reticulum
Extracellular Region
Extracellular Matrix
External Encapsulating Structure
Vacuole
Peroxisome
Lipid Droplet
Organelle
Lysosome
Cytoskeleton
Golgi Apparatus
Nuclear Envelope
Mitochondrion
Nucleolus
Chromosome
Plasma Membrane
Cytoplasmic Vesicle
Cytosol
Nucleus
Ribosome
Cilium
Endosome
Microtubule Organizing Center
Nucleoplasm
p 2.3e-08
p 4.9e-05
p 0.019
p 0.019
p 0.24
p 0.39
p 0.42
p 0.42
p 0.54
p 0.66
p 0.64
p 0.8
p 1
p 0.86
p 0.77
p 0.85
p 0.14
p 0.00036
p 0.00038
p 0.013
p 0.011
p 0.0041
p 0.0087
p 9.7e-05
-2
-1
0
1
2
NES
FDR > 0.05
FDR ≤ 0.05


Female: every term at p ≤ 0.05, in order of p (fgsea running score)




Female: every term at p ≤ 0.05, in order of p (fgsea running score)




Female: every term at p ≤ 0.05, in order of p (fgsea running score)




Female: every term at p ≤ 0.05, in order of p (fgsea running score)




Female: every term at p ≤ 0.05, in order of p (fgsea running score)




Female: every term at p ≤ 0.05, in order of p (fgsea running score)




Female: every term at p ≤ 0.05, in order of p (fgsea running score)




Female: every term at p ≤ 0.05, in order of p (fgsea running score)



Female: every term at p ≤ 0.05, in order of p (fgsea running score)



Female: every term at p ≤ 0.05, in order of p (fgsea running score)



Female: every term at p ≤ 0.05, in order of p (fgsea running score)

