

Sex difference: Treatment × sex, every protein

limpa dpcDE, BH within universe; labelled: FDR ≤ 0.05 and the 10 smallest p




Proteins per tier

Dashed: proteins at p ≤ 0.05 expected by chance in each direction; Π can include p > 0.05




All 162 proteins at p ≤ 0.05, per mouse

Rows z-scored across these mice, clustered on 1 - Pearson r (display only)




Sex difference: every protein at p ≤ 0.05, in order of p

log2 abundance per mouse, bar at the group median; log2FC: female PAS response minus male PAS response




Sex difference: every protein at p ≤ 0.05, in order of p

log2 abundance per mouse, bar at the group median; log2FC: female PAS response minus male PAS response




Sex difference: every protein at p ≤ 0.05, in order of p

log2 abundance per mouse, bar at the group median; log2FC: female PAS response minus male PAS response




Sex difference: every protein at p ≤ 0.05, in order of p

log2 abundance per mouse, bar at the group median; log2FC: female PAS response minus male PAS response




Sex difference: every protein at p ≤ 0.05, in order of p

log2 abundance per mouse, bar at the group median; log2FC: female PAS response minus male PAS response




Sex difference: every protein at p ≤ 0.05, in order of p


log2 abundance per mouse, bar at the group median; log2FC: female PAS response minus male PAS response



Sex difference: every protein at p ≤ 0.05, in order of p

log2 abundance per mouse, bar at the group median; log2FC: female PAS response minus male PAS response




Sex difference: every protein at p ≤ 0.05, in order of p
log2 abundance per mouse, bar at the group median; log2FC: female PAS response minus male PAS response
log2 abundance
Cmbl
log2FC 0.83
p 0.043, FDR 0.59, Π 0.074
Apobec2
log2FC 1.44
p 0.044, FDR 0.59, Π 0.011
Ppib
log2FC 0.71
p 0.044, FDR 0.59, Π 0.11
Acyp2
log2FC 0.85
p 0.044, FDR 0.59, Π 0.069
Ufc1
log2FC -1.40
p 0.045, FDR 0.59, Π 0.013
Thbs4
log2FC 0.87
p 0.045, FDR 0.59, Π 0.067
Pptc7
log2FC -0.77
p 0.046, FDR 0.59, Π 0.093
Col1a1
log2FC 1.59
p 0.046, FDR 0.59, Π 0.0073
Alg9
log2FC 0.60
p 0.046, FDR 0.59, Π 0.16
Uqcc5
log2FC -0.75
p 0.047, FDR 0.59, Π 0.1
Myom1
log2FC 1.82
p 0.047, FDR 0.59, Π 0.0039
Gaa
log2FC 0.95
p 0.047, FDR 0.59, Π 0.055
Srsf5
log2FC 1.01
p 0.048, FDR 0.59, Π 0.046
Sod2
log2FC 0.86
p 0.048, FDR 0.59, Π 0.074
Gls
log2FC 1.25
p 0.048, FDR 0.59, Π 0.022
Dhodh
log2FC -0.84
p 0.048, FDR 0.59, Π 0.077
Park7
log2FC 0.62
p 0.048, FDR 0.59, Π 0.15
Ptdss2
log2FC 0.68
p 0.048, FDR 0.59, Π 0.13
Itga7
log2FC -0.56
p 0.049, FDR 0.59, Π 0.18
Vps29
log2FC 0.75
p 0.049, FDR 0.59, Π 0.11
VEH_F
PAS_F
VEH_M
PAS_M

Sex difference: every protein at p ≤ 0.05, in order of p

log2 abundance per mouse, bar at the group median; log2FC: female PAS response minus male PAS response

