




Sex difference: GO:BP
12 strongest terms the EnrichmentMap flag keeps; * FDR≤0.05
up: PAS raised the set more in females than in males
Muscle Contraction
Response To Toxic Substance
Adaptive Thermogenesis
External Encapsulating Structure Organization
Organic Anion Transport
Response To Xenobiotic Stimulus
Establishment Of Protein Localization To Mitochondrion
Animal Organ Morphogenesis
Response To Abiotic Stimulus
Phosphorylation
Actin Filament Based Process
Muscle Cell Development
-log10 p
down
up
log2 FC
NES
3
2
1
0
-1
-2
-3


Sex difference: Reactome


*12 strongest terms the EnrichmentMap flag keeps; * FDR≤0.05*
up: PAS raised the set more in females than in males





Sex difference: MitoCarta
12 strongest terms the EnrichmentMap flag keeps; * FDR≤0.05
up: PAS raised the set more in females than in males
Carbohydrate Metabolism *
Lipid Metabolism *
Lysine Metabolism
Detoxification
Branched Chain Amino Acid Metabolism
SLC25A Family
Proteases
Apoptosis
Civ Assembly Factors
Nucleotide Metabolism
Complex I
Mitochondrial Central Dogma *
-log10 p
down
up
log2 FC
NES
3
2
1
0
-1
-2
-3


Sex difference: signal by GO Slim family
fgsea; a term counts in every family it belongs to; every family with a term at p ≤ 0.05
Anatomical structure development
Cytoskeleton organization
Muscle system process
Cell differentiation
Signaling
Nervous system process
Nucleobase-containing small molecule metabolic process
Detoxification
Vesicle-mediated transport
Telomere organization
Protein maturation
Protein catabolic process
Programmed cell death
Mitotic cell cycle
Membrane organization
Generation of precursor metabolites and energy
Circulatory system process
Cell adhesion
Carbohydrate metabolic process
Carbohydrate derivative metabolic process
0
5
10
GO:BP terms at p ≤ 0.05 (down | up)
down
up



20 strongest GO:BP and Reactome terms, ordered by shared genes
EnrichmentMap similarity, average linkage; black ring FDR ≤ 0.05
Regulation Of Phosphorylation
Phosphorylation
Regulation Of Transferase Activity
Protein Modification Process
Cell Cycle
Adaptive Thermogenesis
Cellular Response To Toxic Substance
Response To Toxic Substance
Non Integrin Membrane ECM Interactions
External Encapsulating Structure Organization
Striated Muscle Cell Differentiation
Muscle Cell Development
Myofibril Assembly
Sarcomere Organization
Cellular Component Assembly Involved In Morphogenesis
Actomyosin Structure Organization
Actin Filament Based Process
Muscle Contraction
Striated Muscle Contraction
Muscle Contraction
-2
-1
0
1
2
NES
NES
3
2
1
0
-1
-2
-3
-log10(p)
2.7
3.0
3.3
3.6
3.9


GO Slim cellular compartments
fgsea, full proteome
Extracellular Matrix
External Encapsulating Structure
Extracellular Region
Vacuole
Lysosome
Cytoskeleton
Organelle
Lipid Droplet
Endoplasmic Reticulum
Mitochondrion
Plasma Membrane
Nuclear Envelope
Chromosome
Golgi Apparatus
Peroxisome
Nucleolus
Cytoplasmic Vesicle
Cytosol
Ribosome
Nucleus
Cilium
Endosome
Microtubule Organizing Center
Nucleoplasm
p 0.0008
p 0.001
p 8.1e-05
p 0.029
p 0.051
p 0.044
p 0.13
p 0.29
p 0.17
p 0.52
p 0.72
p 0.64
p 0.96
p 0.96
p 0.82
p 0.81
p 0.85
p 0.19
p 0.1
p 0.017
p 0.052
p 0.019
p 0.0059
p 5.6e-05
-2
-1
0
1
2
NES
FDR > 0.05
FDR ≤ 0.05


Sex difference: every term at p ≤ 0.05, in order of p (fgsea running score)




Sex difference: every term at p ≤ 0.05, in order of p (fgsea running score)




Sex difference: every term at p ≤ 0.05, in order of p (fgsea running score)




Sex difference: every term at p ≤ 0.05, in order of p (fgsea running score)




Sex difference: every term at p ≤ 0.05, in order of p (fgsea running score)




Sex difference: every term at p ≤ 0.05, in order of p (fgsea running score)



Sex difference: every term at p ≤ 0.05, in order of p (fgsea running score)




Sex difference: every term at p ≤ 0.05, in order of p (fgsea running score)




Sex difference: every term at p ≤ 0.05, in order of p (fgsea running score)


Collagen Formation
NES 1.80, p 0.0026, FDR 0.042, n 16
singscore p 0.012




Cellular Response To Chemical Stress
NES 1.77, p 0.0027, FDR 0.042, n 40
singscore p 0.076




Extracellular Matrix Organization
NES 1.65, p 0.0028, FDR 0.042, n 56
singscore p 0.085




ECM Proteoglycans
NES 1.84, p 0.003, FDR 0.042, n 18
singscore p 0.04




Binding And Uptake Of Ligands By Scav...
NES 1.80, p 0.0033, FDR 0.042, n 15
singscore p 0.061




Cell Cycle Mitotic
NES -1.73, p 0.0035, FDR 0.042, n 33
singscore p 0.011




Degradation Of The Extracellular Matrix
NES 1.77, p 0.0042, FDR 0.046, n 21
singscore p 0.021



Cellular Responses To Stimuli
NES 1.41, p 0.013, FDR 0.13, n 120
singscore p 0.29




Mitotic G2 G2 M Phases
NES -1.72, p 0.015, FDR 0.14, n 18
singscore p 0.014



Mitochondrial Protein Import
NES 1.50, p 0.019, FDR 0.16, n 22
singscore p 0.058



M Phase
NES -1.54, p 0.022, FDR 0.18, n 30
singscore p 0.019



Integrin Cell Surface Interactions
NES 1.51, p 0.031, FDR 0.22, n 21
singscore p 0.072



Sex difference: every term at p ≤ 0.05, in order of p (fgsea running score)

