




Sex difference: GO:BP
12 strongest terms after the EnrichmentMap collapse; * FDR≤0.05
Actin Filament Based Process
Muscle Cell Development
Muscle Contraction
Response To Toxic Substance
Striated Muscle Cell Differentiation
Cellular Component Assembly Involved In Morphogenesis
Myofibril Assembly
Sarcomere Organization
Regulation Of Phosphorylation
Regulation Of Transferase Activity
Protein Modification Process
Phosphorylation
-log10 p
down
up
log2 FC
NES
3
2
1
0
-1
-2
-3


Sex difference: MitoCarta

*12 strongest terms after the EnrichmentMap collapse; * FDR≤0.05*





Sex difference: signal by GO Slim family
fgsea; a term counts in every family it belongs to; every family with a term at p ≤ 0.05
Anatomical structure development
Cytoskeleton organization
Muscle system process
Cell differentiation
Signaling
Nervous system process
Nucleobase-containing small molecule metabolic process
Detoxification
Vesicle-mediated transport
Telomere organization
Protein maturation
Protein catabolic process
Programmed cell death
Mitotic cell cycle
Membrane organization
Generation of precursor metabolites and energy
Circulatory system process
Cell adhesion
Carbohydrate metabolic process
Carbohydrate derivative metabolic process
0
5
10
GO:BP terms at p ≤ 0.05 (down | up)
down
up



20 strongest GO:BP terms, ordered by shared genes
EnrichmentMap similarity, average linkage; black ring FDR ≤ 0.05
Temperature Homeostasis
Adaptive Thermogenesis
Homeostatic Process
Organic Anion Transport
Cellular Response To Toxic Substance
Response To Toxic Substance
Regulation Of Phosphorylation
Phosphorylation
Regulation Of Transferase Activity
Protein Modification Process
Cell Cycle
External Encapsulating Structure Organization
Striated Muscle Cell Differentiation
Muscle Cell Development
Myofibril Assembly
Sarcomere Organization
Cellular Component Assembly Involved In Morphogenesis
Actomyosin Structure Organization
Actin Filament Based Process
Muscle Contraction
-2
-1
0
1
2
NES
NES
3
2
1
0
-1
-2
-3
-log10(p)
2.4
2.7
3.0
3.3


GO Slim cellular compartments
fgsea, full proteome
Extracellular Matrix
External Encapsulating Structure
Extracellular Region
Vacuole
Lysosome
Cytoskeleton
Organelle
Lipid Droplet
Endoplasmic Reticulum
Mitochondrion
Plasma Membrane
Nuclear Envelope
Chromosome
Golgi Apparatus
Peroxisome
Nucleolus
Cytoplasmic Vesicle
Cytosol
Ribosome
Nucleus
Cilium
Endosome
Microtubule Organizing Center
Nucleoplasm
p 0.0008
p 0.001
p 8.1e-05
p 0.029
p 0.051
p 0.044
p 0.13
p 0.29
p 0.17
p 0.52
p 0.72
p 0.64
p 0.96
p 0.96
p 0.82
p 0.81
p 0.85
p 0.19
p 0.1
p 0.017
p 0.052
p 0.019
p 0.0059
p 5.6e-05
-2
-1
0
1
2
NES
FDR > 0.05
FDR ≤ 0.05


Sex difference: every term at p ≤ 0.05, in order of p (fgsea running score)




Sex difference: every term at p ≤ 0.05, in order of p (fgsea running score)




Sex difference: every term at p ≤ 0.05, in order of p (fgsea running score)




Sex difference: every term at p ≤ 0.05, in order of p (fgsea running score)




Sex difference: every term at p ≤ 0.05, in order of p (fgsea running score)




Sex difference: every term at p ≤ 0.05, in order of p (fgsea running score)



Sex difference: every term at p ≤ 0.05, in order of p (fgsea running score)




Sex difference: every term at p ≤ 0.05, in order of p (fgsea running score)




Sex difference: every term at p ≤ 0.05, in order of p (fgsea running score)


Translation
NES -1.77, p 0.00087, FDR 0.0075, n 81




Lipid Metabolism
NES 1.60, p 0.0055, FDR 0.039, n 68



Lysine Metabolism
NES 1.59, p 0.022, FDR 0.14, n 12



Detoxification
NES 1.50, p 0.031, FDR 0.17, n 33



Complex I
NES -1.36, p 0.049, FDR 0.22, n 45

