




Male: perfect separations against chance
Dot: observed; grey: median and 95th percentile over all 252 relabellings of the same mice
p 1.000
Clusters
p 1.000
Complexes
p 0.103
Modules
p 0.341
GO:BP / MitoCarta
p 0.540
Proteins
0
10
20
30
40
50
Features separating PAS from vehicle perfectly



Male: AUC of every feature
Proteins
GO:BP / MitoCarta
Modules
Complexes
Clusters
Features
AUC (1: higher in PAS)
0.00
0.25
0.50
0.75
1.00


Male: Complexes, PAS vs vehicle, every feature at p ≤ 0.05 (n = 1)

Exact Wilcoxon p, BH within level; higher or lower in PAS





Male: Modules, PAS vs vehicle, every feature at p ≤ 0.05 (n = 2)
Exact Wilcoxon p, BH within level; higher or lower in PAS
Pink
0.00 (lower), p 0.008, FDR(
Red
0.08 (lower), p 0.032, FDR(
Sensitivity
1 - specificity
0.00
0.25
0.50
0.75
1.00



Male: GO:BP / MitoCarta, PAS vs vehicle, every feature at p ≤ 0.05 (n = 19)
Exact Wilcoxon p, BH within level; higher or lower in PAS
ellular Response ToOxygen
0.00 (lower), p 0.008, FDR(
sma Membrane Organizati
0.00 (lower), p 0.008, FDR(
rotein Complex Oligomeriza
0.00 (lower), p 0.008, FDR(
ctomyosin Structure Organi
0.04 (lower), p 0.016, FDR(
mbraneless Organelle Ass
0.04 (lower), p 0.016, FDR(
Nervous System Process
0.04 (lower), p 0.016, FDR(
Positive Regulation Of Prot...
0.96 (higher), p 0.016, FDR
egulation Of Muscle Contra
0.04 (lower), p 0.016, FDR(
esponse To Transforming Gi
0.04 (lower), p 0.016, FDR(
Cell Cell Junction Organiza...
0.08 (lower), p 0.032, FDR(
Cell Substrate Junction Org...
0.08 (lower), p 0.032, FDR(
Multicellular Organismal Mo.
0.08 (lower), p 0.032, FDR(
egative Regulation Of Prot.
0.08 (lower), p 0.032, FDR(
Organelle Assembly
0.08 (lower), p 0.032, FDR(
Phosphorylation
0.92 (higher), p 0.032, FDR
rotein Homooligomerizatio
0.08 (lower), p 0.032, FDR(
Renal System Development
0.08 (lower), p 0.032, FDR(
ensory System Developmen
0.08 (lower), p 0.032, FDR(
upramolecular Fiber Organ
0.08 (lower), p 0.032, FDR(
Sensitivity
1 - specificity
0.00
0.25
0.50
0.75
1.00


Male: Proteins, PAS vs vehicle, every feature at p ≤ 0.05 (n = 52)
Exact Wilcoxon p, BH within level; higher or lower in PAS
Epb42
C0.00 (lower), p 0.008, FDI
Tpm1
C0.00 (lower), p 0.008, FDI
Tpm2
C0.00 (lower), p 0.008, FDI
Ywhah
C1.00 (higher), p 0.008, FD
Atp5pf
C0.00 (lower), p 0.008, FDI
Saraf
C1.00 (higher), p 0.008, FD
Tnnt3
C0.00 (lower), p 0.008, FDI
Plin4
C0.04 (lower), p 0.016, FDI
Cilp2
C0.04 (lower), p 0.016, FDI
Prx
C0.04 (lower), p 0.016, FDI
Nefl
C0.04 (lower), p 0.016, FDI
Alpl
C0.04 (lower), p 0.016, FDI
Dmd
C0.04 (lower), p 0.016, FDI
Prph
C0.04 (lower), p 0.016, FDI
Dpep1
C0.04 (lower), p 0.016, FDI
Capzb
C0.04 (lower), p 0.016, FDI
Cav1
C0.04 (lower), p 0.016, FDI
Mybph
C0.04 (lower), p 0.016, FDI
Fahd2a
C0.04 (lower), p 0.016, FDI
Jagn1
C0.96 (higher), p 0.016, FD
Sensitivity
1 - specificity
0.00
0.25
0.50
0.75
1.00

Male: Proteins, PAS vs vehicle, every feature at p ≤ 0.05 (n = 52)

Exact Wilcoxon p, BH within level; higher or lower in PAS




Male: Proteins, PAS vs vehicle, every feature at p ≤ 0.05 (n = 52)
Exact Wilcoxon p, BH within level; higher or lower in PAS
Trim72
C 0.08 (lower), p 0.032, FDI
Pptc7
C 0.92 (higher), p 0.032, FD
Lnpk
C 0.92 (higher), p 0.032, FD
Aldh5a1
C 0.92 (higher), p 0.032, FD
Plpp7
C 0.92 (higher), p 0.032, FD
Gde1
C 0.92 (higher), p 0.032, FD
Myo1c
IC 0.08 (lower), p 0.032, FDI
Kcnj11
C 0.92 (higher), p 0.032, FD
Atl2
C 0.88 (higher), p 0.048, FD
Myl1
IC 0.12 (lower), p 0.048, FDI
Cmbl
IC 0.12 (lower), p 0.048, FDI
Ccdc47
C 0.88 (higher), p 0.048, FD
Sensitivity
1 - specificity
0.00
0.25
0.50
0.75
1.00