



Proteins, every contrast
Full proteome; dashed line p = 0.05
PAS vs VEH, female
PAS vs VEH, male
Treatment × sex
Sex in VEH
Sex in PAS
-log10 p
10.0
7.5
5.0
2.5
0.0
-2.5
0.0
2.5
5.0
log2 fold change
FDR ≤ 0.05
p > 0.05
p ≤ 0.05
Π ≤ 0.05


Proteins moving with each phenotype
Across PAS and vehicle mice of each sex; 92 at p ≤ 0.05 by chance
CSA
Capillary density
Satellite cells / fibre
Soleus pyruvate
Plantaris pyruvate
Soleus octanoyl
120
FDR 0
133
FDR 0
466
FDR 193
430
FDR 94
136
FDR 4
157
FDR 6
384
FDR 7
464
FDR 86
274
FDR 14
320
FDR 42
457
FDR 129
406
FDR 83
56
FDR 0
96
FDR 0
439
FDR 92
522
FDR 197
40
FDR 0
43
FDR 0
411
FDR 67
463
FDR 106
177
FDR 2
187
FDR 2
28
FDR 0
25
FDR 0
Female
Female + mito share
Male
Male + mito share
p ≤ 0.05 / chance
5
4
3
2
1


PAS effect on each phenotype
Welch t on log values within sex, n = 5 v 5
CSA
Capillary density
Satellite cells / fibre
Soleus pyruvate
Plantaris pyruvate
Soleus octanoyl
p > 0.05
p ≤ 0.05
Female
Male
-50%
+0%
+100%
+300%
+700%
PAS vs vehicle, % change (95% CI, log scale)