



Proteins, every contrast
Full proteome; dashed line p = 0.05
PAS vs VEH, female
PAS vs VEH, male
Treatment × sex
Sex in VEH
Sex in PAS
-log10 p
10.0
7.5
5.0
2.5
0.0
-2.5
0.0
2.5
5.0
log2 fold change
FDR ≤ 0.05
p > 0.05
p ≤ 0.05
p and Π ≤ 0.05


Proteins tracking each phenotype
Per-sex slopes, adjusted for pellet mitochondrial share; 92 at p ≤ 0.05 by chance
CSA
Capillary density
Satellite cells / fibre
Soleus pyruvate
Plantaris pyruvate
Soleus octanoyl
87
FDR 0
443
FDR 135
112
FDR 0
191
FDR 5
392
FDR 11
281
FDR 0
257
FDR 0
480
FDR 110
264
FDR 0
43
FDR 0
416
FDR 34
83
FDR 0
59
FDR 0
518
FDR 166
163
FDR 1
205
FDR 4
24
FDR 0
136
FDR 2
Female slope
Male slope
Male − female
p ≤ 0.05 / chance
5
4
3
2
1


PAS effect on each phenotype
Welch t on log values within sex, n = 5 v 5
CSA
Capillary density
Satellite cells / fibre
Soleus pyruvate
Plantaris pyruvate
Soleus octanoyl
p > 0.05
p ≤ 0.05
Female
Male
-50%
+0%
+100%
+300%
+700%
PAS vs vehicle, % change (95% CI, log scale)