
a What the proteome PCs track
R² of each factor on each component
PC1 (43.1%)
PC2 (11.1%)
PC3 (8.7%)
PC4 (6.8%)
Sex
0.89
0.08
0.00
0.00
Treatment
0.00
0.00
0.02
0.21
Treatment × sex
0.01
0.06
0.01
0.07
MitoCarta share
0.88
0.03
0.00
0.00
Run depth (log2 precursors)
0.69
0.23
0.00
0.01
R²
0.0
0.5
1.0
c Perfect separators against relabelling
Dot: observed; bar: 0 to the null's 95th centile; tick: null median
Pathways
Female
Male
p = 0.13
p = 0.33
0
10
20
30
Modules
p = 0.063
p = 1
0
1
Complexes
Female
Male
p = 0.35
p = 1
0
1
2
3
4
Clusters
p = 1
p = 1
0
1
features separating perfectly
b Modules against contrasts and phenotypes
r-equivalent of each t, bold if absolute r ≥ 0.4; AUC cells show the AUC, filled by 2·AUC − 1; J: subsample stability
Contrast
Female traits
Male traits
AUC (PAS vs vehicle)
Female
Male
Treatment × sex
CSA
Capillaries
Sat. cells
Sol. pyr
Pla. pyr
Sol. oct
CSA
Capillaries
Sat. cells
Sol. pyr
Pla. pyr
Sol. oct
Female
Male
84%
turquoise (J 0.58) · OXPHOS
-0.28 0.44 -0.48 -0.31 -0.22 -0.12 -0.05 -0.04 0.20 0.63 0.37 -0.09 -0.13 -0.41 0.29 0.36 0.68
3%
blue (J 0.49)
0.48 0.28 0.18 0.56 0.43 0.57 0.07 0.16 0.23 -0.08 0.15 0.53 -0.25 -0.13 -0.17 1.00 0.64
7%
brown (J 0.46)
0.19 0.50 -0.26 0.15 0.19 0.18 0.03 -0.02 0.38 0.41 0.31 0.26 -0.04 -0.37 -0.04 0.60 0.84
22%
yellow (J 0.46) · Selenoamino Acid Metabolism
-0.10 -0.20 0.07 -0.00 -0.01 -0.06 -0.45 -0.12 0.01 -0.43 -0.24 0.07 -0.15 0.04 0.04 0.40 0.44
4%
green (J 0.53) · Platelet Activation Signaling And Aggregation
-0.00 -0.17 0.12 -0.05 0.11 -0.13 -0.30 -0.11 -0.05 0.06 -0.20 -0.25 0.17 0.45 0.17 0.40 0.32
13%
red (J 0.67) · Myogenesis
0.18 -0.33 0.35 0.22 -0.12 0.31 0.47 0.39 -0.41 -0.28 0.09 -0.17 0.04 0.25 0.05 0.52 0.08
37%
black (J 0.50) · Selenoamino Acid Metabolism
-0.30 0.09 -0.27 -0.24 -0.09 -0.29 -0.49 -0.31 0.27 0.26 -0.17 -0.24 0.12 -0.12 0.25 0.20 0.52
10%
pink (J 0.29)
0.00 -0.17 0.12 0.14 0.08 0.06 -0.22 -0.00 -0.14 0.04 -0.31 -0.19 0.06 0.17 0.08 0.56 0.40
100
50
0
proteins (% MitoCarta)
p ≤ 0.05
r-equivalent or 2·AUC − 1
-1
0
1