

Betweenness in Enriched Categories

Ruben Mud

November 16, 2024


Main Result

Theorem

There are functors $\mathbf{BetSp} \begin{matrix} \xleftarrow{L} \\ \xrightarrow{R} \end{matrix} \mathbf{EnCat}$


Examples of Betweenness

- ▶ Pasch (1882):



Examples of Betweenness

- ▶ Pasch (1882):




- ▶ Menger (1928): In a metric space (X, d)

$$d(a, x) + d(x, b) = d(a, b)$$

Examples of Betweenness

- ▶ Pasch (1882):



- ▶ Menger (1928): In a metric space (X, d)

$$d(a, x) + d(x, b) = d(a, b)$$


- ▶ Glivenko (1937): In a lattice (L, \wedge, \vee, \leq) ,

$$(a \wedge x) \vee (x \wedge b) = x = (a \vee x) \wedge (x \vee b)$$

Definition

A relation $[\cdot, \cdot, \cdot] \subseteq X^3$ is a betweenness if

- (B1) Symmetry: $[a, x, b] \longleftrightarrow [b, x, a]$,
- (B2) Reflexivity: $[a, b, b]$ holds for all $a, b \in X$,
- (B3) Minimality: $[a, b, a]$ and $[b, a, b]$, then $a = b$
- (B4) Transitivity: $[a, x, b]$ and $[a, y, x]$, then $[a, y, b]$.



Definition (Bankston, 2013)

A road system on X is a family $\mathcal{R} \subseteq 2^X$ such that:

1. $\{a\} \in \mathcal{R}$ for all $a \in X$,
2. for all $a, b \in X$ there exists $R \in \mathcal{R}$ such that $a, b \in R$.

Definition

x is between a and b whenever

$$x \in \bigcap_{a, b \in R \in \mathcal{R}} R$$

“All roads from a to b go via x ”

New Example

“All functions $f : A \longrightarrow B$ go via X ”

New Example

“All functions $f : A \longrightarrow B$ go via X ”

Composition is surjective

$$\mathrm{Hom}(A, X) \times \mathrm{Hom}(X, B) \longrightarrow \mathrm{Hom}(A, B)$$

New Example

“All functions $f : A \longrightarrow B$ go via X ”

Composition is surjective

$$\mathrm{Hom}(A, X) \times \mathrm{Hom}(X, B) \longrightarrow \mathrm{Hom}(A, B)$$

$$d(a, x) + d(x, b) \geq d(a, b)$$

$R : \mathbf{EnCat} \longrightarrow \mathbf{BetSp}$

On a \mathcal{V} -category \mathcal{A} define $[-, -, -]_V \subseteq \text{ob}(\mathcal{A})^3$

$[A, B, C]_V$ if and only if M_{ABC} and M_{CBA} are split epi

Where $M_{ABC} : \mathcal{A}(A, B) \otimes \mathcal{A}(B, C) \longrightarrow \mathcal{A}(A, C)$

$R : \mathbf{EnCat} \longrightarrow \mathbf{BetSp}$

On a \mathcal{V} -category \mathcal{A} define $[-, -, -]_V \subseteq \text{ob}(\mathcal{A})^3$

$[A, B, C]_V$ if and only if M_{ABC} and M_{CBA} are split epi

Where $M_{ABC} : \mathcal{A}(A, B) \otimes \mathcal{A}(B, C) \longrightarrow \mathcal{A}(A, C)$

This definition is symmetric (B1)

$$[A, B, C] \longleftrightarrow [C, B, A]$$

$[A, B, B]$ holds $\forall A, B \in \text{ob}(\mathcal{A})$.

$$\begin{array}{ccccc}
 \mathcal{A}(A, B) \otimes \mathcal{A}(B, B) & \xrightarrow{M} & \mathcal{A}(A, B) & \xleftarrow{M} & \mathcal{A}(A, A) \otimes \mathcal{A}(A, B) \\
 \uparrow 1 \otimes j_B & \nearrow r & & \nwarrow l & \uparrow j_A \otimes 1 \\
 \mathcal{A}(A, B) \otimes I & & & & I \otimes \mathcal{A}(A, B)
 \end{array}$$

$[A, B, B]$ holds $\forall A, B \in \text{ob}(\mathcal{A})$.

$$\begin{array}{ccccc}
 \mathcal{A}(A, B) \otimes \mathcal{A}(B, B) & \xrightarrow{M} & \mathcal{A}(A, B) & \xleftarrow{M} & \mathcal{A}(A, A) \otimes \mathcal{A}(A, B) \\
 \uparrow 1 \otimes j_B & \nearrow r & & \nwarrow l & \uparrow j_A \otimes 1 \\
 \mathcal{A}(A, B) \otimes I & & & & I \otimes \mathcal{A}(A, B)
 \end{array}$$

$$M_{ABB}(1 \otimes j_B)r^{-1} = \text{Id}$$

if $[A, X, B]$ and $[A, Y, X]$ then $[A, Y, B]$

$$\begin{array}{ccc}
 (\mathcal{A}(A, Y) \otimes \mathcal{A}(Y, X)) \otimes \mathcal{A}(X, B) & \xrightarrow{a} & \mathcal{A}(A, Y) \otimes (\mathcal{A}(Y, X) \otimes \mathcal{A}(X, B)), \\
 \downarrow M_{AYX} \otimes 1 \quad \uparrow \varphi_{AYX} \otimes 1 & & \downarrow 1 \otimes M_{YXB} \\
 \mathcal{A}(A, X) \otimes \mathcal{A}(X, B) & \xleftarrow{\varphi_{AXB}} & \mathcal{A}(A, Y) \otimes \mathcal{A}(Y, B) \\
 \searrow M_{AXB} & \swarrow M_{AYB} & \\
 & \mathcal{A}(A, B) &
 \end{array}$$

if $[A, X, B]$ and $[A, Y, X]$ then $[A, Y, B]$

$$\begin{array}{ccc}
 (\mathcal{A}(A, Y) \otimes \mathcal{A}(Y, X)) \otimes \mathcal{A}(X, B) & \xrightarrow{a} & \mathcal{A}(A, Y) \otimes (\mathcal{A}(Y, X) \otimes \mathcal{A}(X, B)), \\
 \downarrow M_{AYX} \otimes 1 \quad \uparrow \varphi_{AYX} \otimes 1 & & \downarrow 1 \otimes M_{YXB} \\
 \mathcal{A}(A, X) \otimes \mathcal{A}(X, B) & \begin{array}{c} \xleftarrow{\varphi_{AXB}} \\ \xrightarrow{M_{AXB}} \end{array} & \mathcal{A}(A, Y) \otimes \mathcal{A}(Y, B) \\
 & \searrow M_{AYB} & \swarrow \\
 & \mathcal{A}(A, B) &
 \end{array}$$


$$M_{AYB}(1 \otimes M_{YXB})a(\varphi_{AYX} \otimes 1)\varphi_{AXB} = \text{Id}$$

Since $[A, B, A]_{\mathcal{V}}, [B, A, B]_{\mathcal{V}}$ is an equivalence relation,

$$(\mathcal{A}, \mathcal{V}) \longmapsto (\text{ob}(\mathcal{A}) / \sim, [-, -, -]_{\mathcal{V}})$$

defines the object part of our functor $L : \mathbf{EnCat} \longrightarrow \mathbf{BetSp}$.

\mathbb{R}^2 is a Category



We can think of points $f \in \overrightarrow{AB}$ as morphisms $f : A \longrightarrow B$

the functor $\mathbf{BetSp} \longrightarrow \mathbf{EnCat}$

Given $(X, [\cdot, \cdot, \cdot]) \in \mathbf{BetSp}$, define a 2^X -category X .

the functor $\mathbf{BetSp} \longrightarrow \mathbf{EnCat}$

Given $(X, [\cdot, \cdot, \cdot]) \in \mathbf{BetSp}$, define a 2^X -category X .

- Objects: points in X ,

the functor $\mathbf{BetSp} \longrightarrow \mathbf{EnCat}$

Given $(X, [\cdot, \cdot, \cdot]) \in \mathbf{BetSp}$, define a 2^X -category X .

- Objects: points in X ,
- Hom-objects: $X(a, b) := \{x \in X : [x, b, a]\}$

the functor $\mathbf{BetSp} \longrightarrow \mathbf{EnCat}$

Given $(X, [\cdot, \cdot, \cdot]) \in \mathbf{BetSp}$, define a 2^X -category X .

- ▶ Objects: points in X ,
- ▶ Hom-objects: $X(a, b) := \{x \in X : [x, b, a]\}$
- ▶ Composition: $X(a, b) \cap X(b, c) \subseteq X(a, c)$

the functor $\mathbf{BetSp} \longrightarrow \mathbf{EnCat}$

Given $(X, [\cdot, \cdot, \cdot]) \in \mathbf{BetSp}$, define a 2^X -category X .

- Objects: points in X ,
- Hom-objects: $X(a, b) := \{x \in X : [x, b, a]\}$
- Composition: $X(a, b) \cap X(b, c) \subseteq X(a, c)$

$$([x, b, a] \text{ and } [x, c, b]) \xrightarrow{B^4} [x, c, a]$$

the functor $\mathbf{BetSp} \longrightarrow \mathbf{EnCat}$

Given $(X, [\cdot, \cdot, \cdot]) \in \mathbf{BetSp}$, define a 2^X -category X .

- Objects: points in X ,
- Hom-objects: $X(a, b) := \{x \in X : [x, b, a]\}$
- Composition: $X(a, b) \cap X(b, c) \subseteq X(a, c)$

$$([x, b, a] \text{ and } [x, c, b]) \xrightarrow{B^4} [x, c, a]$$

- Unit element: $X \subseteq X(a, a)$

$[x, a, a]$ holds for all $x \in X$ by B2

Given $f : X \longrightarrow Y \in \mathbf{BetSp}$.

$$[a, b, c]_X \implies [f(a), f(b), f(c)]_Y$$

Given $f : X \longrightarrow Y \in \mathbf{BetSp}$.

$$[a, b, c]_X \implies [f(a), f(b), f(c)]_Y$$

$$fX(a, b) \subseteq Y(f(a), f(b))$$

Given $f : X \longrightarrow Y \in \mathbf{BetSp}$.

$$[a, b, c]_X \implies [f(a), f(b), f(c)]_Y$$

$$fX(a, b) \subseteq Y(f(a), f(b))$$

Thus we can view Y as enriched over 2^X via $f^{-1}(Y(f(a), f(b)))$

$$\begin{array}{ccc} X(a, x) \cap X(x, b) & \longrightarrow & X(a, b) \\ \downarrow & & \downarrow \\ f^{-1}(Y(fa, fx)) \cap f^{-1}(Y(fx, fb)) & \longrightarrow & f^{-1}(Y(fa, fb)) \end{array}$$

and

$$\begin{array}{ccc} X & \longrightarrow & X(x, x) \\ & \searrow & \downarrow \\ & & f^{-1}(Y(fx, fx)) \end{array}$$

commute.

Functoriality

$$\begin{array}{ccc} X(a, x) \cap X(x, b) & \longrightarrow & X(a, b) \\ \downarrow & & \downarrow \\ f^{-1}(Y(fa, fx)) \cap f^{-1}(Y(fx, fb)) & \longrightarrow & f^{-1}(Y(fa, fb)) \end{array}$$

and

$$\begin{array}{ccc} X & \longrightarrow & X(x, x) \\ & \searrow & \downarrow \\ & & f^{-1}(Y(fx, fx)) \end{array}$$

commute.

Thus $f : X \longrightarrow Y \in \mathbf{EnCat}$



F.William Lawvere (1973) *Metric Spaces, Generalized Logic, and Closed Categories*, Rendiconti del Seminario Matematico e Fisico di Milano, 43: 135 – 166



V. Pambuccian (2011) *The Axiomatics of Ordered Geometry I. Ordered Incidence Spaces*, Expositiones Mathematicae 29: 24 – 66