



Published in final edited form as:

Nano Today. 2023 June ; 50: . doi:10.1016/j.nantod.2023.101822.

Two decades of vaccine development against atherosclerosis

Miguel A. Moreno-Gonzalez^{a,b,1}, Oscar A. Ortega-Rivera^{a,b,1}, Nicole F. Steinmetz^{a,b,c,d,e,f,*}

^aDepartment of NanoEngineering, University of California-San Diego, La Jolla, CA 92039, USA

^bCenter for Nano-ImmunoEngineering, University of California-San Diego, La Jolla, CA 92039, USA

^cInstitute for Materials Discovery and Design, University of California-San Diego, La Jolla, CA 92039, USA

^dDepartment of Bioengineering, University of California-San Diego, La Jolla, CA 92039, USA

^eDepartment of Radiology, University of California-San Diego, La Jolla, CA 92039, USA

^fMoore's Cancer Center, University of California-San Diego, La Jolla, CA 92039, USA

Abstract

Atherosclerosis is an immune-mediated chronic inflammatory disease that leads to the development of fatty plaques in the arterial walls, ultimately increasing the risk of thrombosis, stroke, and myocardial infarction. The immune response in this complex disease is both atheroprotective and pro-atherogenic, involving both innate and adaptive immunity. Current treatments include the adjustment of lifestyle factors, cholesterol-lowering drugs such as statins, and immunotherapy, whereas vaccine development has received comparatively little attention. In this review, we discuss the potential of antigen-specific vaccination as a preventative approach based on more than 20 years of research and innovation. Vaccination targets include proteins that are more abundant in atherosclerotic patients, such as oxidized low-density lipoprotein (LDL), apolipoprotein B-100, proprotein convertase subtilisin/kexin type-9 serine protease (PCSK9), cholesteryl ester transfer protein (CETP), and heat shock proteins HSP60 and HSP65. Immunization with such proteins or their peptide epitopes has been shown to induce T-cell activation, produce antigen-specific antibodies, reduce the size of atherosclerotic lesions, and/or reduce serum cholesterol levels. Vaccination against atherosclerosis therefore offers a new strategy to address the burden on healthcare systems caused by cardiovascular disease, the leading cause of death worldwide.

*Corresponding author at: Department of NanoEngineering, University of California-San Diego, La Jolla, CA 92039, USA. nsteinmetz@ucsd.edu (N.F. Steinmetz).

¹Co-first author.

Declaration of Competing Interest

The authors declare the following financial interests/personal relationships which may be considered as potential competing interests: Dr. Steinmetz is a co-founder of, has equity in, and has a financial interest with Mosaic ImmunoEngineering Inc. Dr. Steinmetz serves as Director, Board Member, and Acting Chief Scientific Officer, and paid consultant to Mosaic. The other authors declare no potential conflicts of interest.

Keywords

Atherosclerosis; Vaccines; Nanotechnology

Introduction

Cardiovascular disease (CVD) is the leading cause of mortality worldwide, with ~18 million CVD-related fatalities in 2019 representing 31% of all deaths. [1] The major underlying cause of cardiovascular events is atherosclerosis, a chronic inflammatory disease with an autoimmune component, which involves the disruption of normal cholesterol homeostasis. [2,3] In healthy individuals, cholesterol is delivered to cells by low-density lipoprotein (LDL) and is removed for excretion by high-density lipoprotein (HDL). During atherosclerosis, LDL particles accumulate in the arterial intima to form fibro-fatty plaques, which promote local inflammation. This leads to the recruitment of macrophages and T-cells that exacerbate the condition by releasing pro-inflammatory cytokines. Ultimately, the inflammation and hemodynamic anomalies cause local failures of vascular integrity that can trigger thrombosis leading to myocardial infarction and ischemic stroke [2,4,5].

In the clinic, the first-line approach for the prevention and treatment of atherosclerosis is the adjustment of lifestyle factors that contribute to high serum LDL levels, including poor diet, smoking, and lack of exercise. Therapeutic intervention typically involves a group of drugs known as statins, which reduce levels of LDL-cholesterol (LDL-C) in the blood by inhibiting the hepatic enzyme hydroxymethylglutaryl-coenzyme A (HMG-CoA) reductase, a rate-limiting step in the cholesterol biosynthesis pathway. [3] The long-term use of statins, however, can increase the expression of proprotein convertase subtilisin/kexin type-9 serine protease (PCSK9), a hepatic enzyme that inhibits the LDL receptor (LDL-R), thus preventing cholesterol metabolism. Two monoclonal antibody (mAb) therapies against PCSK9 (alirocumab and evolocumab) were therefore approved in 2015 for patients in which statins are ineffective or poorly tolerated. The FDA also recently approved siRNA-based gene therapy against PCSK9 to reduce levels of LDL-C. However, many atherosclerosis patients remain at risk due to chronic inflammation. [6,7].

The ability to therapeutically target components of the cholesterol homeostatic network and/or pro-inflammatory cascade has led to proposals for atherosclerosis vaccines based on two different principles: reducing serum cholesterol levels and blocking the pro-inflammatory response to LDL-C accumulation. In this review, we consider the molecular basis of atherosclerosis and the associated inflammation and then focus on key targets for vaccine development. These include LDL (primarily responsible for the formation of atherosclerotic plaques) and its major component apolipoprotein B (ApoB); cholesteryl ester transfer protein (CETP), which transfers cholesteryl ester between HDL and LDL; PCSK9, which regulates the LDL-R; and heat shock proteins (HSP60 and HSP65), which form part of the stress-response pathway in the arterial walls and are involved in the autoimmune response. We carried out a systematic analysis of patents, academic publications, and clinical trials to provide a comprehensive analysis of the state of the art in atherosclerosis vaccines and to evaluate their feasibility as a strategy to address this chronic disease.

The immunology of atherosclerosis

The formation of atherosclerotic plaques in humans and animal models triggers both innate and adaptive immune responses (Fig. 1). The initiating event is the retention of infiltrating LDL particles on proteoglycans in the subendothelial space. [8] The trapped lipoproteins are modified by proteases and lipases, promoting their ability to bind proteoglycans and thus causing their aggregation. [9] They are also modified by myeloperoxidase, lipoxygenase, and reactive oxygen and nitrogen species to form oxidized LDL (oxLDL), which elicits an inflammatory response from the innate immune system. [10].

Endothelial cells respond to the accumulation of modified lipoproteins by expressing adhesion molecules such as vascular cell adhesion protein 1 (VCAM-1). [11] Circulating monocytes and other leukocytes are recruited to these sites (Fig. 1). Infiltrating monocytes differentiate into macrophages in response to macrophage colony-stimulating factor (M-CSF) and granulocyte-macrophage colony stimulating factor (GM-CSF), which are produced by endothelial cells and several other cell types. Macrophages thus become a major cell population in atherosclerotic plaques. [12] The engulfment of lipoproteins leads to transformation of macrophages into foam cells. Macrophages express scavenger receptors, among which scavenger receptor class A and CD36 are the most important for the uptake of modified LDL. Foam cells lose their ability to migrate and eventually die within the arterial intima, thus creating a core area in the plaque that consists of apoptotic and necrotic cells, cholesterol crystals, and other extracellular material. [13,14].

Atherosclerotic plaques also accumulate T cells and cells that express major histocompatibility complex class II (MHC-II), forming the basis of the adaptive immune response in atherosclerosis. HLA-DR is expressed by several cell types in the plaque and presents antigens to CD4⁺ T cells. Dendritic cells in atherosclerotic plaques and in the adjacent adventitia take up plaque-derived antigens and migrate to lymph nodes, where they display these antigens to naive T cells. [15,16] Autoantigens derived from LDL particles are important for atherosclerosis and the frequent presence of anti-oxLDL antibodies indicates that B cells react to oxLDL. These antibodies tend to be more prevalent in patients with coronary artery disease than in healthy controls. [17] Anti-oxLDL antibodies can be either IgM or IgG, indicating that an isotype class-switch occurs, which requires assistance from T cells. Indeed, T cells from patients with atherosclerosis recognize LDL components displayed by antigen-presenting cells (APCs). [2,18].

The main adaptive effector cells in atherosclerotic plaques are CD4⁺ helper T (Th) cells, some of which react to ApoB-100 peptides presented by APCs on MHC-II. [19] Both oxLDL and native LDL contain ApoB-derived T-cell epitopes. However, APCs express scavenger receptors that recognize oxLDL, so oxidation increases the uptake of LDL for antigen presentation. Th1 cells are the most abundant Th cells in the plaques, and they secrete interferon γ (IFN γ), which promotes monocyte infiltration, macrophage activation, and foam-cell formation, as well as inducing Th1 cell differentiation in synergy with IL-12, which is also produced in the plaque. [2] Th2 cells are much less abundant than Th1 cells [12]. The Th2-related cytokine IL-4 appears to promote atherosclerosis, whereas IL-5, IL-13 and IL-33 limit disease development. [2,5] Pro-atherogenic conditions and oxLDL

can also induce a small population of Th17 cells in plaques, although there are conflicting reports concerning the role of their signature cytokine (IL-17 A), which is also produced by other cells. [2,5] Small populations of regulatory T (T_{reg}) cells expressing forkhead box protein P3 (FOXP3) are also found in atherosclerotic plaques, where they control auto-reactive T-cell clones and disease progression and thus play an important protective role during atherogenesis. [20] Whereas $CD4^{+}$ T cells are present in plaques during all disease stages, $CD8^{+}$ T cells are more abundant during the very early stages of lesion development but seem to have a lesser impact than $CD4^{+}$ T cells in hyperlipidemic mice. The proportion of $CD8^{+}$ T cells gradually decreases as the lesion matures. $CD8^{+} T_{reg}$ cells may limit atherosclerosis by controlling pro-atherosclerotic pathways. [2,5].

B cells are also found in atherosclerotic plaques. [5] B1 cells represent a first-line innate defense against common pathogens and secrete germline-encoded IgM antibodies in a T-cell-independent manner. In cardiovascular disease, the titers of IgMs recognizing LDL or ApoB epitopes are inversely correlated with atherosclerosis, complications, and outcome. [21] IgMs that recognize oxLDL inhibit its uptake by macrophages and prevent myeloid cell inflammation, suggesting an atheroprotective role for B1 cells. [22] In contrast, B2 cells are activated by Tfh to differentiate into plasma cells that secrete IgG antibodies. IgG titers against native and oxidized LDL or ApoB are positively correlated with atherosclerotic disease in mice and humans, suggesting B2 cells are pro-atherogenic. Accordingly, the inhibition of B2 cells could be atheroprotective, whereas specifically interfering with plasma cell functions is largely pro-atherogenic. [5] However, the GLACIER clinical phase II study using a monoclonal IgG1 antibody against a human ApoB peptide failed to show a reduction in atherosclerotic inflammation, so the role of IgG remains controversial. [23] It is worth mentioning that this clinical trial only tested one mAb, so more trials must be conducted to make general conclusions on the atheroprotective potential of oxidized LDL or ApoB IgG.

Analysis of atherosclerosis treatment: from passive immunotherapy to vaccines

Vaccination is a form of active immunotherapy intended to elicit the production of antibodies by the patient. Before considering the use of vaccination as a strategy to prevent and treat atherosclerosis, it is therefore important to consider passive immunotherapies that have been developed for this disease. This approach involves the therapeutic administration of recombinant antibody products that block the function of target proteins by inhibiting their activity and/or leading to their degradation.

The most advanced antibody therapies against atherosclerosis are those that target PCSK9. Two products have been approved (alirocumab and evolocumab), both of which target and neutralize PCSK9 in the blood, preventing it from binding to the LDL receptor. The FOURIER clinical trial ([ClinicalTrials.gov](https://clinicaltrials.gov/ct2/show/study/NCT01764633) ID: [NCT01764633](https://clinicaltrials.gov/ct2/show/study/NCT01764633)) showed that evolocumab can reduce circulating LDL-C levels by 59% in patients with atherosclerotic cardiovascular disease who are concurrently treated with statins. This was associated with a 15% reduction in the risk of composite of cardiovascular death, myocardial infarction, stroke, hospitalization for unstable angina, or coronary revascularization. [24] The ODYSSEY

OUTCOMES clinical trial ([ClinicalTrials.gov](https://clinicaltrials.gov/ct2/show/study/NCT01663402) ID: [NCT01663402](https://clinicaltrials.gov/ct2/show/study/NCT01663402)) showed that alirocumab can reduce circulating LDL-C levels by 62.7%, 61.0% and 54.7% after 4, 12 and 48 months of treatment, respectively. [25] There have been 60 clinical trials of evolocumab and alirocumab that have tested various demographics such as age, race, ethnicity and genetics, demonstrating positive results for mortality, safety and efficacy. [26–85] However, the drawbacks of passive immunotherapy include the cost and the need for repeated dosing and lifelong treatment, which make the drugs inaccessible for most patients. The cost in the USA is \$4,500-\$8,000 per year if used to treat higher-risk patients with LDL cholesterol 100 mg/dL despite intensive statin therapy, which is prohibitive in developing countries. [86].

Active immunization (vaccination) is a promising alternative for the treatment of atherosclerosis, following the remarkable effects of vaccination campaigns against various infectious diseases, [87] as well as more recent advances against cancer and chronic diseases, including CVD. [88] However, given that the therapeutic targets are self-antigens, there are a few risks and challenges that must be addressed. The first obstacle when vaccinating against self-antigens is to overcome immune tolerance. This can be achieved through potent carriers and adjuvants capable of mounting a strong immune response against weakly immunogenic targets. Several nanotechnology-based adjuvants, such as nanoliposomes and virus-like particles (VLPs), have proven effective (this will be elaborated in more detail in the next sections). Based on their size regimen, these nanoparticle formulations are effective at interacting with immune cells, and they have highly immunostimulatory properties. [89,90] For immunotherapies targeting self, the second challenge is avoiding target-specific immunotoxicity. Mounting humoral responses against a self-target is preferred and safer, because cytotoxic CD8⁺ T cell responses against a self-antigen may lead to autoimmune diseases. [91] However, there are preclinical studies of vaccines that decrease atherosclerotic lesions by increasing CD8⁺ T cells. [92,93] Overall, if efficacious and safe vaccination can be achieved, this would be superior over other therapeutic approaches such as mAb, siRNA, and gene editing. mAb therapies are effective, but they are expensive, they require repeated dosing, and they lead to poor patient compliance compared to vaccination. In the case of siRNA, there is one FDA approved therapy, Inclisiran, but again requires lifelong treatment. [58] Finally, gene editing is a new and promising alternative, but the approaches are at early stages. [94] Verve Therapeutics recently started the first phase 1b clinical trial to evaluate the safety of PCSK9 gene editing medicine, and they expect results in 2023 ([ClinicalTrials.gov](https://clinicaltrials.gov/ct2/show/study/NCT05398029) Identifier: [NCT05398029](https://clinicaltrials.gov/ct2/show/study/NCT05398029)).

We therefore carried out a detailed quantitative analysis of vaccination approaches for atherosclerosis, aiming to identify all the targets that have been considered in preclinical studies and clinical trials. We searched the peer-reviewed literature in PubMed Central using the keywords “atherosclerosis vaccine” and only included results from the year 2000 onwards since the conception of the idea to develop a vaccine started during the late 90’s. [17,18] We then evaluated the specific targets that have been considered as antigens, using keywords such as oxidized LDL (oxLDL), apolipoprotein B-100 (ApoB-100), heat shock proteins (HSPs), cholesteryl ester transfer protein (CETP), and proprotein convertase subtilisin/kexin type 9 (PCSK9). The relevant results were refined to a list of 113 original research articles and reviews. We also screened the [ClinicalTrials.gov](https://clinicaltrials.gov/) database for vaccines

and antibody therapies against atherosclerosis, revealing 7 vaccine products targeting PCSK9, CETP and angiotensin II, as well as 60 clinical trials for monoclonal antibody therapies against PCSK9 and 1 against LDL.

We also conducted a patent search using the website [Lens.com](https://lens.com) and the keywords “atherosclerosis” and “vaccine” in the title, abstract and claims. This resulted in a list of 973 patents (granted patents and applications) which we refined to 248 by restricting the date of registration (starting from 2000) and manual screening for common target antigens. The list included patents covering monoclonal antibodies against PCSK9 (67), CD20 (1) and the phosphorylcholine head group (70).

Ultimately, this combined screen of the academic literature, clinical trials, and patents over the last 21 years (Fig. 2) enabled us to define three broad categories of targets: (1) lipid-related antigens, (2) non-lipid-related antigens, and (3) pathogen-related antigens, because infections can also cause CVD (Table 1). The most prominent lipid-related antigens were LDL, ApoB, CETP and PCSK9, whereas the most prominent non-lipid antigens were HSP60 and HSP65. The intended effect of vaccination was either to reduce serum cholesterol levels or block the inflammatory response to the formation of atherosclerotic plaques.

Next, we compiled Table 2, which lists the 77 preclinical studies on atherosclerosis vaccines targeting LDL, ApoB, CETP, PCSK9, HSP60/HSP65 and angiotensin. The table also shows the year of publication, specific antigen, carrier used (if any), adjuvant used (if any), animal model, and summary of major results that can be used as a point of comparison such as percent reduction in atherosclerotic lesion area (sometimes reported as plaque size, lesion size or lesion area), antibody titers, T cell and T_{reg} activation, protein target inhibition, effects on cholesterol (total cholesterol, LDL-C, and HDL-C) and LDL-R, effects on macrophages and dendritic cells, and effects on certain cytokines, such as IFN γ , tumor necrosis factor α (TNF α), IL-10, IL-4 and transforming growth factor β (TGF β). Although we could find only one paper describing angiotensin-based atherosclerosis vaccines, most of the clinical trials on atherosclerosis have targeted this protein. The other preclinical studies involving angiotensin mainly focused on hypertension, which is a major risk factor for atherosclerosis, so we decided to include them in this study. The specific amino acid sequence of each peptide vaccine is also included, based on UniProt data. If the organism for each peptide was not specified, we assumed that the authors referred to human proteins. If the protein was described as murine, we assumed the species was the mouse rather than any other rodent.

Major target proteins for vaccine development

Low-density lipoprotein

The predominant reason for the onset of atherosclerosis is the accumulation of LDL on the arterial walls. LDL consists of cholesterol, triglycerides, lipophilic antioxidants, phospholipids and ApoB-100, the major lipoprotein component.[97,166] ApoB-100 in the blood is usually described as “bad cholesterol” because it can cause the buildup of cholesterol in the arteries. Native LDL does not elicit an immune response, but oxLDL

is a major factor in the development of atherosclerosis. [166] One pathway for the oxidation of LDL is through exposure to the enzyme myeloperoxidase (MPO), hydrogen peroxide (H_2O_2) and reactive nitrogen species (NO_2^-). [167].

The pro-atherogenic protein oxLDL is detected by APCs such as macrophages and dendritic cells, which process the protein and present epitopes to T cells. Macrophages engulf oxLDL as a defense mechanism, and the epitopes are presented on the cell surface via MHC-I and MHC-II molecules. [98] CD8^+ cytotoxic T cells and CD4^+ helper T cells then recognize and bind these MHC molecules, initiating the production of oxLDL-specific T cells. As discussed above, these are primarily Th1 cells that produce pro-inflammatory cytokines such as $\text{IFN}\gamma$, IL-2, IL-12 and $\text{TNF}\alpha$. The uptake of oxLDL by macrophages also promotes the formation of cholesterol-rich foam cells that accumulate on the arterial wall, creating fatty streaks and exacerbating the inflammation. [100].

The oxLDL epitopes also give rise to oxLDL-specific autoantibodies in humans and animal models. [99,96,95] Most such autoantibodies are the Th1-specific IgG2a isotype. [98] A Th1 response is a pro-inflammatory reaction and confers an undesirable prognosis, with higher IgG2a titers correlating with more severe manifestations. [95,166] Based on the recruitment of T cells, the activation of macrophages, and the production of antibodies, it is evident that oxLDL can initiate both cellular and humoral immune responses. [95,100].

Several studies have explored the effects of vaccination with oxLDL epitopes. [99,100,102,95,97] Homologous malondialdehyde (MDA)-LDL antigens were tested for the treatment of apolipoprotein E knockout mice (*ApoE^{-/-}*), revealing no change in the IgM titer but a substantial increase in the titer of T-cell dependent IgG. [95] Treatment with MDA-LDL reduced the mean lesion size by 46%. Similarly, oxLDL-pulsed dendritic cells in LDL-R knockout mice (*Ldlr^{-/-}*) induced high titers of T-cell-dependent IgG against ox-LDL, resulting in an 87% reduction in plaque size in the carotid arteries and a 75% reduction in the abundance of $\text{IFN}\gamma$. [100] The fact that the upregulation of IgG confers an atheroprotective effect, while high levels of circulating IgG are indicators of a bad prognosis shows how beautiful and complex this disease is. Perhaps this could explain why it has been challenging to develop treatments that can succeed in the clinic.

Another treatment option that has been explored in preclinical models is the induction of mucosal tolerance to oxLDL. [98,101] This is based on the hypothesis that tolerance to the autoantigen should ameliorate the inflammatory effects, as reported for other Th1-mediated autoimmune diseases such as multiple sclerosis, rheumatoid arthritis, and type 1 diabetes. [98] One possible tolerance mechanism mediated by autoantigens is the upregulation of immunosuppressive T_{reg} cells, thus maintaining self-tolerance. It is noted that there are contradictions in targeting oxLDL and distinct approaches are being pursued: While immunization against oxLDL aims to upregulate the IgG humoral immune response, tolerance induction aims to upregulate the T_{reg} cellular immune response to prevent autoimmune response associated with the disease. These are distinct mechanism approaches targeting the same molecule. Oral tolerance to oxLDL was shown to reduce mean lesion areas by up to 71.2% in carotid arteries by the upregulation of $\text{CD4}^+\text{CD25}^+\text{Foxp3}^+$ T_{reg} cells

and TGF β , [98] whereas nasal tolerance induced a modest 47.6% reduction in lesion size by promoting the accumulation of CD4⁺LAP⁺ and CD4⁺CD25⁺Foxp3⁺ T_{reg} cells. [101].

Apolipoprotein B-100

ApoB-100 is a major component of the LDL protein and it binds oxLDL products such as MDA via its histidine and lysine residues. [103] Once MDA binds to ApoB-100, the ApoB-100 protein and its peptide sequences are targeted by the immune system. Plasma antibodies from CVD patients recognize immunogenic ApoB-100 epitopes, confirming this protein is an important target for atherosclerosis vaccines. [103,105,168].

The 302-amino-acid ApoB-100 protein contains several highly immunogenic peptides. [166,104,93,169,103,105,106,170,168,171] Scanning the entire human ApoB-100 protein in 20-amino-acid windows revealed 102 peptides that elicit a humoral immune response. [166,169] The MDA-modified human ApoB-100 peptides p143 (ApoB₂₁₃₁₋₂₁₅₀, IALDDAKINFNEKLSQLQTY) and p210 (ApoB₃₁₃₆₋₃₁₅₅, KTTKQSFDSLVSVAQYKKNKH) led to a 60% reduction in atherosclerotic lesion area when administered with an alum adjuvant. [103] Similarly, MDA-modified p45 (human ApoB₆₈₈₋₇₀₇, IEIGLEGKGFEPTLEALFGK) led to a 48% reduction in atherosclerotic lesion area, whereas the lesser-known p74 (ApoB₁₁₂₃₋₁₁₄₂, VISIPRLQAEARSEILAHWS) led to a 31% reduction in aortic lesion area. [105] More importantly, both peptides converted a pro-inflammatory Th1 response to an anti-inflammatory Th2 response, as evident from the isotype switch from IgG2a to IgG1. The same group also tested the p45 (human ApoB₆₆₁₋₆₈₀, IEIGLEGKGFEPTLEALFGK) and p210 peptides, which reduced atherosclerotic lesion area by 66% and 59%, respectively despite no increase in the titer of peptide-specific IgG, which was initially considered an important correlate of efficacy. [106] This suggested that the protective effect of the vaccines was independent of the antibodies, and may reflect a cellular rather than a humoral immune response. It is important to note that despite both studies being done by the same researchers, they identified P45 with two different sequences, but with the same amino acids.

To test this hypothesis, other groups investigated the MHC-II-restricted murine ApoB epitopes ApoB₃₅₀₁₋₃₅₁₆ (SQEYSGSVANEANVYL) and ApoB₉₇₈₋₉₉₃ (TGAYSNASSTESASY), [110] p101 (ApoB₇₀₅₋₇₂₀), p102 (ApoB₄₄₁₋₄₅₆, TLYALSHAVNSYFDVD) and p103 (ApoB₃₉₅₃₋₃₉₆₈, LYYKEDKTSLASAAS), [112] p18 (ApoB₃₀₃₀₋₃₀₄₄, SLFFSAQPFEITAST), [114] and p6 (ApoB₉₇₈₋₉₉₃, TGAYSNASSTESASY) [113] rather than B-cell epitopes, achieving lesion size reductions of 35–52% accompanied by significant increase in IL-10 production.

This newfound knowledge inspired other researchers to look for different pathways that could explain the atheroprotective effects of ApoB-100 epitope vaccines. The depletion of T_{reg} cells is known to worsen atherosclerotic plaques, whereas the addition of these cells ameliorates the condition. [107] One promising alternative was to investigate the potential upregulation of T_{reg} cells by ApoB-100 vaccines as already shown in oxLDL tolerance experiments. Researchers at Lund University immunized mice with the so-called aBp210 vaccine based on p210, resulting in a 37% reduction in atherosclerotic lesion area in the aorta accompanied by a fivefold expansion of splenic CD4⁺CD25⁺Foxp3⁺ T_{reg} cells. [107]

To confirm that the atheroprotective effect was mediated by the expansion of T_{reg} cells, they administered anti-CD25 antibodies that block the effect of circulating T_{reg} cells and found that this neutralized the effects of the aBp210 vaccine. Other researchers working with p210 obtained similar results accompanied by an expansion of Foxp3⁺ T_{reg} cells, a decrease in the levels of pro-inflammatory cytokines such as IFN- γ and IL-17, and an increase in the level of anti-inflammatory cytokine IL-10. [109,20,111].

To determine whether the atheroprotective effect could also be mediated by CD8⁺ cytotoxic T cells, the CD4⁺ vs CD8⁺ immune response was evaluated following the administration of p210. [92] This achieved a significant 57% reduction in aortic lesion area, but the CD4⁺ T cell levels did not increase following immunization. Surprisingly, the CD8⁺ T-cell population expanded whereas dendritic cells were depleted at the immunization sites, suggesting that the adaptive immune response was interfering with the innate response. CD8⁺CD25⁺ T cells were responsible for the effectiveness of the p210 vaccine, and that the adoptive transfer of CD8⁺ T cells from the p210-immunized mice to recipient mice could also protect against atherosclerosis.

Cholesteryl ester transfer protein

High levels of HDL-C correlate with a lower risk of CVD. [172] Perhaps the most promising target to increase HDL-C levels is CETP, a hydrophobic glycoprotein that circulates in the plasma and is mainly secreted by the liver. [166,173] This protein binds to HDL and promotes the removal of cholesterol from tissues and its transport to the liver for reverse cholesterol transfer, which leads to cholesterol excretion. CETP maintains equilibrium lipid concentrations by mediating the transport of cholesteryl esters, triglycerides and phospholipids between lipoprotein particles, but it also promotes atherogenesis by transferring cholesteryl ester from the antiatherogenic HDL to the pro-atherogenic LDL and very-low-density lipoprotein (VLDL). CETP therefore has both pro-atherogenic and anti-atherogenic functions (Fig. 3).

The pro-atherogenic properties of CETP were originally investigated in transgenic mice. Expression of the human *CETP* gene in mice reduced HDL levels and led to an increase in the abundance of VLDL and LDL. [173] Furthermore, expression of the human *CETP* gene in *ApoE*^{-/-} and *Ldlr*^{-/-} mice resulted in the development and progression of atherosclerosis. [166] Based on these results, the pro-atherogenic activity of CETP has been inhibited using anti-CETP antibodies, which were shown to lower the plasma concentration of CETP and in turn increase the amount of HDL-C in circulation. [116,117,123,119,115,120] CETP-specific antibodies have been induced by administering a DNA vaccine consisting of the plasmid pCR-X8-HBc-CETP, which encodes the B-cell epitope of the CETP C-terminal fragment (CETPC) displayed by hepatitis B virus core (HBc) particles. [119] Intramuscular immunization of female New Zealand white rabbits resulted in the production of anti-CETP antibodies, leading to an 80.6% reduction in aortic lesion area, lower total cholesterol and LDL-C levels, and higher HDL-C levels compared to control rabbits on a high cholesterol diet. The same pCR-X8-HBc-CETP plasmid combined with the linear polysaccharide chitosan to form chitosan/pCETP nanoparticles was used for the intranasal immunization

of male New Zealand white rabbits, leading to a 59.2% reduction in aortic lesion size accompanied high titers of anti-CETP IgG and lower levels of LDL-C. [120].

Researchers at the Universidad Nacional Autonoma de Mexico recently administered their therapeutic intranasal vaccine (HB-ATV-8) to pigs, which more closely resemble human anatomy, physiology and biochemistry than rodents, thus providing a better prediction of performance in clinical trials. [126] The HB-ATV-8 vaccine is based on micellar nanoparticles consisting of the C-terminal residues H486–S496 of CETP, which are necessary for lipid binding and transfer. As expected, the immunized animals produced anti-CETP IgG and the aortal atherosclerotic lesions were less severe. Indeed, 50% of the vaccinated pigs lacked any atherosclerotic lesions and only type I lesions (with isolated macrophage foam cells) were found in the other 50%. These are excellent results compared to the pigs that consumed a high fat diet but received no vaccination, which developed type II and type III lesions characterized by intracellular lipid accumulation and small extracellular lipid pools, respectively.

Researchers have also developed vaccines that combine B-cell epitopes of CETP with tetanus toxoid (TT) residues that act as helper T cell epitopes. [124,125,127,118,122,115,121] The first CETP/TT vaccine combined TT₈₃₀₋₈₄₃ (QYIKANSKFIGITE) with human CETP₄₆₁₋₄₇₆ (INPEITRDGFLLLQM, the residues needed for lipid transfer). [115] The intramuscular immunization of New Zealand white rabbits led to a 39.6% reduction in aortic lesion area, higher HDL-C levels, and lower LDL-C levels. Similarly, the recombinant chimeric enzyme vaccine AnsB-TTP-CETPC combined the enzyme asparaginase (AnsB), TT₈₃₁₋₈₅₄ (YIKANSKFIGITELKKLESKINKV) and human CETP₄₄₈₋₄₇₆ (ALMNSKGVSLFDIINPEITRDGFLLLQM). [118] This vaccine, administered subcutaneously, induced anti-CETP antibodies that inhibited CETP activity, increasing HDL-C levels and reducing LDL-C levels in the immunized rabbits, and reducing the size of aortic atherosclerotic plaques by 42.3% compared to the ovalbumin (OVA) neutral control and by 47.6% compared to the rHSP65 negative control. Subcutaneous CETP/TT vaccines have also been shown to reduce atherosclerotic lesion area and the levels of IL-4, but surprisingly increased the level of IFN γ . [125,127] Other CETP/TT vaccines have been combined with human intestinal trefoil factor (TFF3) to withstand acidity and proteases in the gastrointestinal tract. [122] Oral administration reduced CETP activity and atherosclerotic lesion area, increased the abundance of anti-atherogenic IL-10 and TGF β , but in contrast to earlier reports also depleted pro-atherogenic TNF α and IFN γ . Only one CETP vaccine has been tested in a clinical trial ([ClinicalTrials.gov](https://clinicaltrials.gov/ct2/show/study?term=NCT01284582) ID: NCT01284582). [174] This phase I clinical trial tested the safety, immunogenicity and dose response of ATH03, a vaccine against CETP, in healthy male subjects with HDL-C blood concentrations equal or below 80 mg/dl from May 2011 to July 2012, but no results have been made publicly available.

Proprotein convertase subtilisin/kexin type 9

LDL-C is removed from circulation by its receptor (LDL-R) located on the surface of hepatocytes, which is followed by endocytosis and lysosomal fusion. [128] Within the lysosome, LDL-C is removed and the receptor is recycled back to the cell surface. The

liver also produces the protease PCSK9, which regulates the hepatic availability of LDL-R. [128,175] PCSK9 binds to the extracellular epidermal growth factor-like repeat A (EGFA) domain of LDL-R and prevents recycling, resulting in the lysosomal degradation of the receptor. [89] PCSK9 thus reduces the availability of LDL-R, which in turn causes the levels of LDL-C in the circulation to increase (Fig. 4). PCSK9 gain-of-function mutations are linked to hypercholesterolemia, which is characterized by high levels of LDL-C and an increased risk of atherosclerosis, whereas loss-of-function (LOF) mutations are linked with hypocholesterolemia, which leads to low levels of LDL-C and a lower risk of CVD. [128,175] Therapeutic strategies targeting PCSK9 levels in the blood are therefore an effective approach against atherosclerosis. Fig. 5.

As discussed above, two monoclonal antibodies have already been approved for the targeted inhibition of PCSK9. Alternative approaches include the use of antisense oligonucleotides that interfere with the transcription of the PCSK9 gene [176] or the translation of PCSK9 mRNA. [177] An antisense oligonucleotide targeting the mouse PCSK 9 gene resulted in a 92% reduction in hepatic *PCSK9* mRNA levels in hyperlipidemic mice, boosting the levels of LDL-R substantially. [176] This caused a 52% reduction in total cholesterol, a 32% reduction in LDL, and a 54% reduction in HDL. Furthermore, siRNA molecules in lipidoid nanoparticles have been used to silence the PCSK9 gene by targeting the mRNA for degradation, resulting in lower levels of PCSK9, ApoB and LDL-C in mouse, rat and nonhuman primate models. [178] Built on this, the FDA recently approved siRNA-based gene therapy (Novartis Leqvio® (inclisiran) <https://www.novartis.com>) against PCSK9 to reduce levels of LDL-C. Yet another approach is to use CRISPR gene editing technologies to delete the PCSK9 gene altogether. Musunuru et al. administered CRISPR base editors using lipid nanoparticles (LNPs) in cynomolgus monkeys, and they found that a single dose resulted in a sustained 8 month decrease of PCSK9 and LDL-C of about 90% and 60%, respectively. [94].

The success of PCSK9-specific therapies has generated interest in the development of vaccines based on PCSK9 peptides. [130,179,129,133,131,137,89,134,136,132,135] One promising approach is the use of VLPs based on bacteriophage as a combined carrier and adjuvant for PCSK9 peptides. [130] Specifically, Q β -PCSK9 vaccines were produced by conjugating human PCSK9 peptides (PCSK9₆₈₋₇₆, PCSK9₁₅₃₋₁₆₃ or PCSK9₂₀₇₋₂₂₃) to bacteriophage Q β particles in vitro, and MS2-PCSK9 vaccines were produced by the genetic fusion of PCSK9 peptide sequences (PCSK9₁₅₃₋₁₆₃, PCSK9₁₈₈₋₂₀₀, PCSK9₂₀₈₋₂₂₂ and PCSK9₃₆₈₋₃₈₁) to the MS2 coat protein gene. Both VLP vaccines elicited high titers of IgG antibodies, blocking the binding of PCSK9 to LDL-R, and reducing total cholesterol and LDL-C levels. Mice receiving vaccine PCSK9Q β -003, consisting of the human PCSK9 B-cell epitope peptide (PCSK9₁₅₀₋₁₅₇) conjugated to Q β particles, produced high titers of IgG specific for the PCSK9-003 peptide and were characterized by lower total cholesterol levels, 30% lower free PCSK9 levels, higher levels of LDL-R, lower levels of LDL-C, and a reduction in aortic lesion area. [132,135].

PCSK9 and TT have also been combined with a nanoliposome carrier. In the first such example, the PCSK9 peptide (SIPWNLERITPVR) was combined with TT (AQYIKANSKFIGITEL), resulting in the elicitation of anti-PCSK9 antibodies that inhibited

PCSK9 activity. [89] The specific human PCSK9 sequence was searched in the uniprot database, and we found PCSK9₁₅₃₋₁₆₅ corresponds to SIPWNLERITPPR, but the authors have intentionally altered the sequence in accordance to the AFFiRiS group. This minor change to the sequence helps overcome B cell tolerance as the organism can mount an immune response against the “foreign” rather than “self” PCSK9 peptide. The tetanus toxin sequence could not be found in the uniprot database. In a mouse atherosclerosis model, this vaccine reduced lesion sizes by 39.13%, free PCSK9 levels by 58.5%, PCSK9–LDL-R binding by 69.86%, total cholesterol by 44.7%, and LDL-C by 57%. [134,136] Testing in rhesus macaques stimulated the production of anti-PCSK9 IgG that inhibited PCSK9–LDL-R binding by 33% with no adverse effects. [137] There has been one clinical trial of PCSK9 vaccines AT04A and AT06A from July 2015 to August 2017, which was completed successfully ([ClinicalTrials.gov](https://clinicaltrials.gov/ct2/show/study/NCT02508896) ID: NCT02508896). [180].

Heat shock proteins

Under normal physiological conditions, HSPs facilitate the folding and transport of proteins, [181] and thereby support cell proliferation, immune responses, and protect cells against apoptosis. [182] Humans have evolved cellular and humoral immunity against bacterial HSP65, the homolog of the 60-kDa human heat shock protein 60 (HSP60). [181] However, when cells are exposed to stress factors caused by smoking, hypertension or chronic infection, including the buildup of oxLDL on arterial walls, HSP60 expression is upregulated and may trigger an autoimmune response which is a major factor in the initiation and progression of atherosclerosis. [138,141] Mitochondria in endothelial cells release excess HSP60, which is secreted from the cell. The extracellular HSP60 (exHSP60) then binds to Toll-like receptor 4 (TLR4), inducing the production of the pro-inflammatory cytokines TNF α and IL-6. [182] As the disease progresses, anti-HSP60 antibodies induce complement-mediated or antibody-dependent cellular cytotoxicity (ADCC), resulting in the lysis of endothelial cells. [183].

Researchers have explored the use of vaccines to reinstate HSP60/HSP65 tolerance. However, the subcutaneous administration of HSP60/HSP65 was pro-atherogenic, resulting in larger atherosclerotic plaques. [147] In contrast, oral and nasal vaccines induced the desired tolerance. [145,184,139,142,144,143,140,148,146] In the most effective example, the size of atherosclerotic plaques was reduced by 80.7% and 83.3% following the oral administration of full-length HSP60 protein and the HSP60₂₅₃₋₂₆₈ peptide (SIQSIVPALEIANHR), respectively, in *Ldlr*^{-/-} mice. [140] This induced a significant increase in the abundance of CD4⁺CD25⁺Foxp3⁺ T_{reg} cells, TGF β and IL-10. Similar results were achieved by nasal immunization with recombinant *Helicobacter pylori* HSP60, showing that the reduction of atherosclerotic plaques size can be attributed to the upregulation of CD4⁺CD25⁺GARP⁺ T_{reg} cells that suppress Th1 and Th17 responses, [146] as well as CD4⁺CD25⁻LAP⁺ T_{reg} cells. [144] HSP60 tolerization may also involve the upregulation of myeloid derived suppressor cells (MDSCs). [147].

The mucosal administration of mycobacterial HSP65 delivered by *Lactococcus lactis* achieved a 39.83% reduction of atherosclerotic lesion area, which was attributed to the downregulation of IFN γ and the upregulation of IL-10. [142] Oral tolerance has also been

achieved in *ApoE*^{-/-} mice using recombinant mycobacterial HSP65. [148] As expected, the treated mice had significantly smaller aortic lesions, a larger number of T_{reg} cells in the spleen and draining lymph nodes, and a lower titer of anti-mbHSP65 antibodies compared to untreated controls.

Combinations of vaccines

ApoB-100 has been combined with other antigens such as HSP60, [185,162–164] *Chlamydia pneumonia* (Cpn), [162,164] IL-10, [108] CETP and PCSK9. [90] Most of these antigens have been discussed already with the exception of Cpn antigens, which are known to interact with MHC and may therefore lead to cytotoxic T lymphocyte (CTL) activation. [162] Cpn and HSP60 peptides have been incorporated into different ApoB-100 vaccine formulations with promising outcomes. A vaccine comprising human ApoB₆₈₈₋₇₀₇ (IEIGLEGKGFEPTEALFGK), human HSP60₁₅₃₋₁₆₃ (AELKKQSKPVT), human HSP60₃₀₃₋₃₁₂ (PGFGDNRKNQ), and a Cpn peptide was tested using the repetitive immunization multiple sites strategy (RIMMS) which achieved a reduction in atherosclerotic lesion area of 63.8%. [162] The same group also tested this vaccine in an oral dosing study and found a more modest reduction of atherosclerotic lesions of 46.5%. [164] The same researchers also developed a vaccine using ApoB₆₆₁₋₆₈₀ (FDPNNYLPKESMLKTTTLTAF) and human HSP60₁₅₃₋₁₆₃, which achieved a 41.3% reduction in early atherosclerotic lesion area, [185] and a 39.9% reduction in early plaque size development. [163] A combination of dendritic cells pulsed with ApoB-100 and the immunosuppressive cytokine IL-10 achieved a 70% reduction in aortic lesion area on the descending thoracic aorta. [108] Finally, we recently described an implant-based vaccine comprising the peptides human ApoB₃₁₆₃₋₃₁₈₂ (KTTKQSFDSLVSQAQYKKNKH), human CETP₄₆₁₋₄₇₆ (FGFPEHLLVDFLQSLS), and human PCSK9₂₀₇₋₂₂₃(NVPEEDGTRFHRQASKC) conjugated to bacteriophage Q β VLPs. [90] These trivalent slow-release implants elicited high titers of antigen-specific IgG, inhibiting all three protein targets and reducing total cholesterol levels.

CETP and HSP65 have also been combined to counteract the lipid homeostasis and inflammation pathways simultaneously. An Hsp65-CETP-PADRE-TT-CETP (HCPTC) fusion protein was administered intranasally to male New Zealand white rabbits, leading to a significant reduction in aortic atherosclerotic plaques size, the production of anti-CETP and anti-HSP65 IgG and protective IgA antibodies, lower levels of LDL-C and IFN γ , and an increase in the abundance of IL-10. [165].

Other protein targets

In addition to the major proteins discussed above, other targets have also been explored for vaccination, albeit to a lesser extent, including interleukins, CD99, apoC-III, ANGPTL3, S100A9, and angiotensin. [152,160,186,187,154,188,150,189–192,149,193,194,153,195,158,157,196,197,155,161,198,156,199–201,159,151] Interleukins play a major role in atherogenesis and act as key indicators of vaccine efficacy, but some vaccines have been designed to target these molecules directly, especially pro-inflammatory interleukins such as IL-1 α , [199] IL-12, [194] and IL-15. [190] The IL-1 α -C-Q β vaccine consisted of the full IL-1 α protein chemically conjugated to Q β VLPs, and was shown to elicit high titers of IL-1 α -specific antibodies, reducing plaques size in the descending aorta by 50% and in

the aortic root by 37%. [199] There was also a significant 42% and 32% decrease in the abundance of the inflammatory markers VCAM-1 and ICAM-1, respectively. Vaccination against IL-12 also achieved a 68.5% reduction in atherosclerotic lesion area, [194] whereas vaccination against IL-15 resulted in a 75% reduction in plaque size. [190].

CD99 is a membrane protein found on the surface of leukocytes that promotes the recruitment of macrophages and T cells. [200] The oral administration of a DNA vaccine consisting of *Salmonella typhimurium* transformed with mouse complementary DNA encoding the CD99 extracellular region inhibited CD99 expression in T cells and macrophages, increasing the number of CD8⁺ T cells that induced apoptosis when CD99 is overexpressed, reducing the size of carotid artery lesions by 69%. [200].

ApoC-III and angiopoietin-like protein 3 (ANGPTL3) can be grouped as the only two protein targets with focus on the reduction of triglyceride (TG) levels, a major risk factor for atherosclerosis. TG is transported by pro-atherogenic lipoproteins such as LDL and VLDL, but traditional statin therapies that lower LDL levels are ineffective at controlling TG levels; hence there is a need for more comprehensive therapeutic approaches. [191] ApoC-III is a protein expressed on hepatocytes whose overexpression leads to increased plasma TG levels and cardiovascular disease (CVD) risk. [202] In addition, ApoC-III increases the binding of LDL with the artery wall proteoglycans and activates adhesion molecules on monocytes which leads to more cholesterol retention in the arterial walls. [202] Chackerian et al. developed a VLP-based vaccine using ApoC-III and they saw a 51% reduction in TG levels. [196] Although a permanent reduction in ApoC-III levels by the vaccination with VLPs successfully reduced plasma TG levels in mice, the effects of an ApoC-III targeting vaccine on atherosclerosis and cardiovascular disease remain unknown. ANGPTL3 is a protein secreted by hepatocytes that inhibits lipoprotein lipase (LPL) which results in increased plasma TG. [192] ANGPTL3 loss-of-function mutations result in decreased LDL-C, TG, and CVD risk, making it a promising vaccination target. Unsurprisingly, vaccination against mouse ANGPTL3₃₂₋₄₁ (EPKSRFAMLD) resulted in high antibody titers, a reduction in serum triglycerides and LDL-C, and most importantly a reduction in atherosclerotic lesion sizes in an ApoE^{shl} mouse model. [192] Similarly, ANGPTL3₃₂₋₄₇ (EPKSRFAMLDDVKILA) conjugated to Q β VLPs led to high IgG titers accompanied by a 34% decrease in TG and an increase in LPL. [191] However, effectiveness of these vaccine candidates in preclinical models of atherosclerosis or cardiovascular disease models has not yet been reported.

S100A9 was recently used in our lab as a novel target for the treatment of atherosclerosis. [198] S100A9 is an inflammatory marker secreted by neutrophils of the innate immune system, and it is highly involved in vascular inflammation and atherothrombosis. The vaccine was made by fusing mouse S100A9₁₀₁₋₁₁₀ (RGHGHSHGKG) to the C-terminus of the Q β coat protein (CP) with a GSG linker and it was expressed using *E. coli*. Each nanoparticle had ~90 S100A9 peptides conjugated to the surface which led to high IgG titers. Once the immunogenicity of the soluble vaccine was confirmed, we developed slow release PLGA implants using hot-melt extrusion. The implants elicited high IgG titers with a slight Th1 bias which led to a 32% reduction in atherosclerotic lesion size.

Although angiotensin has not yet been thoroughly investigated as a target for the treatment of atherosclerosis, it is a main target for the treatment of hypertension, another major risk factor for atherosclerosis. This hormone has been the focus of many clinical trials, so we decided to include it in this project. The renin-angiotensin-aldosterone system (RAAS) is a major regulator of systemic blood pressure and blood volume. Briefly, renin is an enzyme secreted by the kidneys which results in the production of the angiotensin I (Ang I) hormone. Ang I is then cleaved to Ang II by the angiotensin-converting enzyme (ACE). [152] Ang II is an active hormone that causes vasoconstriction, high blood pressure, and aldosterone secretion when it binds to the angiotensin receptor subtype 1 (AT₁R). [152,156] The increase in blood pressure leads to the growth of vascular smooth muscle cells and oxidative stress which can contribute to the narrowing of the arterial wall. [203] Ang II is also involved in the recruitment of pro-inflammatory cytokines by activating AT₁R, which leads to the formation of atherosclerotic plaques. [159] Thus, Ang II has emerged as a novel target for atherosclerosis vaccination. In 2016, an AT₁R vaccine consisting of the ATR-001 peptide (CAFHYESQ) conjugated to Q β VLPs was administered subcutaneously to *ApoE*^{-/-} mice, reducing the plaque area of the aortic sinus by 28% and substantially attenuating VCAM-1 and MCP-1 expression. There was also a decrease in the expression of AT₁R and immunostaining revealed less macrophage infiltration in the aortic sinus. As this is the only preclinical study we could find, more research must be conducted to validate Ang II as a viable target for the treatment of atherosclerosis.

Other clinical trials

The only other vaccine trial found in the clinical trials database consisted of an oral vaccine that was composed of pig adipose tissue ([ClinicalTrials.gov](https://clinicaltrials.gov/ct2/show/study/NCT03042741) ID: [NCT03042741](https://clinicaltrials.gov/ct2/show/study/NCT03042741)). This atherosclerosis vaccine, named V6, has been through two small-scale phase II open label clinical trials and it has shown significant improvement in lipid profile in patients with overweight or obesity. However, the ID for this phase II trials or any previous trial are not available in the [ClinicalTrials.gov](https://clinicaltrials.gov/) website and are mentioned in the description of phase III trial and the results for this two small-scale phase II can be accessed only as academic publications. [204,205] This phase III trial started in February 2017 and is still in the recruiting stage. For practical purposes we considered this clinical trial as an ApoB/LDL vaccine during data analysis.

Conclusion

Our comprehensive analysis of the scientific literature, clinical trials and patents surrounding atherosclerosis vaccines over the last two decades has confirmed that the vaccines show promise but that more clinical trials and testing in animal models that closely replicate human lipid metabolism are necessary. We found that the vaccines target two major processes, namely the lowering of cholesterol levels in serum and the blocking of inflammatory responses by inhibiting the pro-inflammatory cascade. We also discovered four major targets (ApoB/LDL, CETP, PCSK9 and HSP60/HSP65), which have been explored extensively along all stages of vaccine innovation (patents, academic publications, and clinical trials). We conclude that the development of atherosclerosis vaccines has the


potential to alleviate the burden of cardiovascular diseases in a cost-effective manner, especially in developing countries, but we are far from a solution.

While the preclinical development pipeline focused on atherosclerosis vaccine candidates is moving rapidly, success in clinical trials is yet to be seen. With improved understanding of the underlying biology and advances in technology, i.e. nanotechnology delivery and adjuvant systems, the field is poised to make a clinical impact. Nanoparticles as carriers and/or adjuvants provide potent alternatives to the traditional adjuvants and delivery technology. Nanoparticles such as nanoliposomes and virus-like particles (VLPs) have already been researched in the setting of atherosclerosis vaccine development, and there is continued effort to move these candidate treatments toward clinical development and testing. Being a plug-and-play technology, vaccine design utilizing nanotechnology can draw from other application areas and implement designs to achieve potency while maintaining safety. Nanoparticles can target and stimulate innate immune cells and promote immunostimulation through promotion of cytokine/chemokine secretion. Targeted and carefully engineered formulations carrying adjuvants, co-stimulatory molecules, as well as antigens or epitopes can be designed to achieve potency while mitigating off-target toxic effects. The COVID-19 pandemic has highlighted that nanotechnology can enable rapid development of vaccines, and with the emergence of novel nanotechnologies, we are confident that platforms will become available that can help address the unique needs for a successful atherosclerosis vaccine, e.g., the need to target self in a chronic disease setting.

Funding

This work was supported in part by grants from the National Institute of Health (R01 HL137674 and R01CA253615 to N.F.S., R01CA253615-02S1 to M.A.M.-G.) and O.A.O.-R. acknowledges the UC MEXUS-CONACYT Postdoctoral Fellowship 2019–2020 no. FE-19-58.

Biographies



Miguel Moreno is a PhD student in NanoEngineering at the University of California, San Diego. Miguel obtained his B.S. in Nanoengineering at UCSD in 2020, and he plans to finish his PhD in 2025. His research focuses on vaccines and immunotherapies based on viral nanoparticles for treatment and prevention of atherosclerosis and cancer. He was awarded an NCI diversity supplement to support his research.



Dr. Ortega-Rivera earned his PhD in Mexico working with oncolytic viruses for cancer immunotherapy. He spent four years as post-doc in the Steinmetz Lab at the University of California, San Diego, where he developed multiple vaccines for cardiovascular diseases, cancer immunotherapy, and against infectious diseases (HPV and SARS-CoV-2). The vaccine development included the combination of virus-like particles and polymers for a slow-release approach as well as microneedle patches for self-administration. He has been honored with prestigious fellowships from CONACYT (Mexico) and UC-MEXUS (UC San Diego). His inventions have been licensed by biotech companies such as Mosaic ImmunoEngineering Inc. and Glycopep Therapeutics.



Dr. Steinmetz is a Professor of NanoEngineering at the University of California, San Diego and Founding Director of the Center for Nano-ImmunoEngineering and Co-Director for the Center of Engineering in Cancer. Dr. Steinmetz trained at The Scripps Research Institute and obtained her PhD from the John Innes Centre in the UK. Her early training was at the RWTH-Aachen University in Germany. Dr. Steinmetz is a Fellow of NAI, BMES, FIAAM, RSC, and AIMBE. She has published more than 200 peer-reviewed publications and is the inventor on more than 100 patents and patent applications, with several patents licensed to biotech companies.

Data Availability

Data will be made available on request.

References

- [1]. Cardiovascular diseases (CVDs) ([https://www.who.int/news-room/fact-sheets/detail/cardiovascular-diseases-\(cvds\)](https://www.who.int/news-room/fact-sheets/detail/cardiovascular-diseases-(cvds))).
- [2]. Gisterå A, Hansson GK, The immunology of atherosclerosis, *Nat. Rev. Nephrol* 13 (2017) 368–380. [PubMed: 28392564]
- [3]. Robinson JG, Heistad DD, Fox KAA, Atherosclerosis stabilization with PCSK-9 inhibition: an evolving concept for cardiovascular prevention, *Atherosclerosis* 243 (2015) 593–597. [PubMed: 26545013]
- [4]. Mallat Z, Binder CJ, The why and how of adaptive immune responses in ischemic cardiovascular disease, *Nat. Cardiovasc Res* 1 (2022) 431–444. [PubMed: 36382200]
- [5]. Wolf D, Ley K, Immunity and inflammation in atherosclerosis, *Circ. Res* 124 (2019) 315–327. [PubMed: 30653442]
- [6]. Boekholdt SM, et al. , Very low levels of atherogenic lipoproteins and the risk for cardiovascular events: a meta-analysis of statin trials, *J. Am. Coll. Cardiol* 64 (2014) 485–494. [PubMed: 25082583]
- [7]. Ridker PM, et al. , Antiinflammatory therapy with canakinumab for atherosclerotic disease, *N. Engl. J. Med* 377 (2017) 1119–1131. [PubMed: 28845751]
- [8]. Skålén K, et al. , Subendothelial retention of atherogenic lipoproteins in early atherosclerosis, *Nature* 417 (2002) 750–754. [PubMed: 12066187]

- [9]. Pentikäinen MO, Oörni K, Ala-Korpela M, Kovanen PT, Modified LDL - trigger of atherosclerosis and inflammation in the arterial intima, *J. Intern Med* 247 (2000) 359–370. [PubMed: 10762453]
- [10]. Hansson GK, Hermansson A, The immune system in atherosclerosis, *Nat. Immunol* 12 (2011) 204–212. [PubMed: 21321594]
- [11]. Endothelial expression of a mononuclear leukocyte adhesion molecule during atherogenesis - PubMed (<https://pubmed.ncbi.nlm.nih.gov/1990440/>).
- [12]. Robbins CS, et al. , Local proliferation dominates lesional macrophage accumulation in atherosclerosis, *Nat. Med* 19 (2013) 1166–1172. [PubMed: 23933982]
- [13]. Moore KJ, Sheedy FJ, Fisher EA, Macrophages in atherosclerosis: a dynamic balance, *Nat. Rev. Immunol* 13 (2013) 709–721. [PubMed: 23995626]
- [14]. Park YM, Febbraio M, Silverstein RL, CD36 modulates migration of mouse and human macrophages in response to oxidized LDL and may contribute to macrophage trapping in the arterial intima, *J. Clin. Invest* 119 (2009) 136–145. [PubMed: 19065049]
- [15]. Jonasson L, Holm J, Skalli O, Gabbiani G, Hansson GK, Expression of class II transplantation antigen on vascular smooth muscle cells in human atherosclerosis, *J. Clin. Invest* 76 (1985) 125–131. [PubMed: 3894417]
- [16]. Llodrá J, et al. , Emigration of monocyte-derived cells from atherosclerotic lesions characterizes regressive, but not progressive, plaques, *Proc. Natl. Acad. Sci. USA* 101 (2004) 11779–11784. [PubMed: 15280540]
- [17]. Bui MN, et al. , Autoantibody titers to oxidized low-density lipoprotein in patients with coronary atherosclerosis, *Am. Heart J* 131 (1996) 663–667. [PubMed: 8721636]
- [18]. Stemme S, et al. , T lymphocytes from human atherosclerotic plaques recognize oxidized low density lipoprotein, *Proc. Natl. Acad. Sci. USA* 92 (1995) 3893–3897. [PubMed: 7732003]
- [19]. Jonasson L, Holm J, Skalli O, Bondjers G, Hansson GK, Regional accumulations of T cells, macrophages, and smooth muscle cells in the human atherosclerotic plaque, *Arteriosclerosis* 6 (1986) 131–138. [PubMed: 2937395]
- [20]. Klingenberg R, et al. , Intranasal immunization with an apolipoprotein B-100 fusion protein induces antigen-specific regulatory T cells and reduces atherosclerosis, *Arterioscler., Thromb., Vasc. Biol* 30 (2010) 946–952. [PubMed: 20167655]
- [21]. Sjögren P, et al. , High plasma concentrations of autoantibodies against native peptide 210 of apoB-100 are related to less coronary atherosclerosis and lower risk of myocardial infarction, *Eur. Heart J* 29 (2008) 2218–2226. [PubMed: 18664466]
- [22]. Que X, et al. , Oxidized phospholipids are proinflammatory and proatherogenic in hypercholesterolaemic mice, *Nature* 558 (2018) 301–306. [PubMed: 29875409]
- [23]. Lehrer-Graiwer J, et al. , FDG-PET imaging for oxidized LDL in stable atherosclerotic disease: a phase II study of safety, tolerability, and anti-inflammatory activity, *JACC Cardiovasc Imaging* 8 (2015) 493–494. [PubMed: 25457756]
- [24]. Sabatine MS, et al. , Evolocumab and clinical outcomes in patients with cardiovascular disease, *N. Engl. J. Med* 376 (2017) 1713–1722. [PubMed: 28304224]
- [25]. Schwartz GG, et al. , Alirocumab and cardiovascular outcomes after acute coronary syndrome, *N. Engl. J. Med.* 379 (2018) 2097–2107. [PubMed: 30403574]
- [26]. Amgen. 2-part, Phase 2/3 Study to Assess the Safety, Tolerability and Efficacy of AMG 145 in Subjects With Homozygous Familial Hypercholesterolemia. Part A - Open-label, Single-arm, Multicenter Pilot Study to Evaluate Safety, Tolerability and Efficacy of AMG 145 in Subjects With Homozygous Familial Hypercholesterolemia. Part B - Double-blind, Randomized, Placebo-controlled, Multicenter Study to Evaluate Safety, Tolerability and Efficacy of AMG 145 in Subjects With Homozygous Familial Hypercholesterolemia (<https://clinicaltrials.gov/ct2/show/NCT01588496>) (2018).
- [27]. Amgen. A Double-Blind, Placebo Controlled, Multicenter Study to Assess the Effect of Evolocumab on Cognitive Function in Patients With Clinically Evident Cardiovascular Disease and Receiving Statin Background Lipid Lowering Therapy: A Study for Subjects Enrolled in the FOURIER (Study 20110118) Trial (<https://clinicaltrials.gov/ct2/show/NCT02207634>) (2017).
- [28]. Amgen. A Double-blind, Randomized, Multicenter Study to Evaluate Safety and Efficacy of AMG 145, Compared With Ezetimibe, in Hypercholesterolemic Subjects Unable to

Tolerate an Effective Dose of a HMG-CoA Reductase Inhibitor <https://clinicaltrials.gov/ct2/show/NCT01763905> (2020).

- [29]. Amgen. A Double-blind, Randomized, Multicenter Study to Evaluate the Safety and Efficacy of Evolocumab, Compared With Ezetimibe, in Hypercholesterolemic Japanese Subjects Unable to Tolerate an Effective Dose of a HMG-CoA Reductase Inhibitor Due to Muscle Related Side Effects (<https://clinicaltrials.gov/ct2/show/NCT02634580>) (2020).
- [30]. Amgen. A Double-blind, Randomized, Multicenter Study to Evaluate the Safety and Efficacy of Evolocumab, Compared With Ezetimibe, in Hypercholesterolemic Subjects Unable to Tolerate an Effective Dose of a HMG-CoA Reductase Inhibitor Due to Muscle Related Side Effects <https://clinicaltrials.gov/ct2/show/NCT01984424> (2018).
- [31]. Amgen. A Double-blind, Randomized, Multi-center, Placebo-controlled, Parallel-group Study to Determine the Effects of Evolocumab (AMG 145) Treatment on Atherosclerotic Disease Burden as Measured by Intravascular Ultrasound in Subjects Undergoing Coronary Catheterization <https://clinicaltrials.gov/ct2/show/NCT01813422> (2019).
- [32]. Amgen. A Double-blind, Randomized, Placebo and Ezetimibe Controlled, Multicenter Study to Evaluate Safety, Tolerability and Efficacy of AMG 145 on LDL-C in Combination With Statin Therapy in Subjects With Primary Hypercholesterolemia and Mixed Dyslipidemia (<https://clinicaltrials.gov/ct2/show/NCT01763866>) (2021).
- [33]. Amgen. A Double-blind, Randomized, Placebo and Ezetimibe-controlled, Multicenter Study to Evaluate Safety and Efficacy of Lipid Lowering Monotherapy With AMG 145 in Subjects With a 10-Year Framingham Risk Score of 10% or Less <https://clinicaltrials.gov/ct2/show/NCT01763827> (2020).
- [34]. Amgen. A Double-blind, Randomized, Placebo-controlled, Multicenter Study Assessing the Impact of Additional LDL-Cholesterol Reduction on Major Cardiovascular Events When Evolocumab (AMG 145) is Used in Combination With Statin Therapy In Patients With Clinically Evident Cardiovascular Disease <https://clinicaltrials.gov/ct2/show/NCT01764633> (2019).
- [35]. Amgen. A Double-Blind, Randomized, Placebo-controlled, Multicenter Study to Evaluate Long-Term Tolerability and Durable Efficacy of AMG 145 (Evolocumab) on LDL-C in Hyperlipidemic Subjects (<https://clinicaltrials.gov/ct2/show/NCT01516879>) (2018).
- [36]. Amgen. A Double-blind, Randomized, Placebo-controlled, Multicenter Study to Evaluate Safety and Efficacy of Evolocumab (AMG 145) in Combination With Statin Therapy in Diabetic Subjects With Hyperlipidemia or Mixed Dyslipidemia (BERSON) (<https://clinicaltrials.gov/ct2/show/NCT02662569>) (2019).
- [37]. Amgen. A Double-blind, Randomized, Placebo-controlled, Multicenter Study to Evaluate Safety, Tolerability and Efficacy of AMG 145 on LDL-C in Subjects With Heterozygous Familial Hypercholesterolemia <https://clinicaltrials.gov/ct2/show/NCT01763918> (2019).
- [38]. Amgen. A Double-blind, Randomized, Placebo-controlled, Multicenter Study to Evaluate Safety, Tolerability and Efficacy of Evolocumab (AMG 145) on LDL-C in Combination With Statin Therapy in Japanese Subjects With High Cardiovascular Risk and With Hyperlipidemia or Mixed Dyslipidemia (<https://clinicaltrials.gov/ct2/show/NCT01953328>) (2015).
- [39]. Amgen. A Double-blind, Randomized, Placebo-controlled, Multicenter Study to Evaluate Tolerability and Efficacy of AMG 145 on LDL-C in Subject With Heterozygous Familial Hypercholesterolemia <https://clinicaltrials.gov/ct2/show/NCT01375751> (2021).
- [40]. Amgen. A Multicenter, Controlled, Open-label Extension (OLE) Study to Assess the Long-term Safety and Efficacy of AMG 145 (Evolocumab) <https://clinicaltrials.gov/ct2/show/NCT01439880> (2019).
- [41]. Amgen. A Multicenter, Controlled, Open-label Extension (OLE) Study to Assess the Long-term Safety and Efficacy of AMG 145 <https://clinicaltrials.gov/ct2/show/NCT01854918> (2019).
- [42]. Amgen. A Multicenter, Open-label Study to Assess the Long-term Safety, Tolerability, and Efficacy of Evolocumab (AMG145) on LDL-C in Subjects With Severe Familial Hypercholesterolemia (<https://clinicaltrials.gov/ct2/show/NCT01624142>) (2019).
- [43]. Amgen. A RaNdomized Double-blInd Placebo ConTrolled Study Characterizing THe Effects of PCSK9 Inhibition On Arterial Wall Inflammation in Patients With Elevated Lp(a) (ANITSCHKOW) (<https://clinicaltrials.gov/ct2/show/NCT02729025>) (2019).

- [44]. Amgen. A Randomized, Actively Controlled, Open-label, Multicenter Study of Efficacy and Safety of Evolocumab Compared With Low Density Lipoprotein Cholesterol (LDL-C) Apheresis, Followed by Single-Arm Evolocumab Administration in Subjects Receiving LDL-C Apheresis Prior to Study Enrollment (<https://clinicaltrials.gov/ct2/show/NCT02585895>) (2017).
- [45]. Amgen. A Randomized, Actively Controlled, Open-label, Multicenter Study of Efficacy and Safety of Evolocumab Compared With Low Density Lipoprotein Cholesterol (LDL-C) Apheresis, Followed by Single-Arm Evolocumab Administration in Subjects Receiving LDL-C Apheresis Prior to Study Enrollment <https://clinicaltrials.gov/ct2/show/NCT02585895> (2017).
- [46]. Amgen. A Randomized, Multicenter Study to Evaluate Tolerability and Efficacy of AMG 145 on LDL-C, Compared With Ezetimibe, in Hypercholesterolemic Subjects Unable to Tolerate an Effective Dose of a HMG-CoA Reductase Inhibitor (<https://clinicaltrials.gov/ct2/show/NCT01375764>) (2021).
- [47]. Amgen. A Randomized, Placebo- and Ezetimibe-controlled, Dose-ranging Study to Evaluate Tolerability and Efficacy of AMG 145 on LDL-C in Hypercholesterolemic Subjects With a 10-year Framingham Risk Score of 10% or Less (<https://clinicaltrials.gov/ct2/show/NCT01375777>) (2021).
- [48]. Amgen. Double-blind, Randomized, Multicenter, Placebo-Controlled Study to Characterize the Efficacy, Safety, and Tolerability of 24 Weeks of Evolocumab for LDL-C Reduction in Pediatric Subjects 10 to 17 Years of Age With HeFH <https://clinicaltrials.gov/ct2/show/NCT02392559> (2020).
- [49]. Amgen. Double-blind, Randomized, Placebo-controlled, Single Site Study to Evaluate the Effects of Evolocumab (AMG 145) Treatment, Alone and in Combination With Atorvastatin, on Lipoprotein Kinetics (<https://clinicaltrials.gov/ct2/show/NCT02189837>) (2018).
- [50]. Goicoechea M Comparative Pilot Study of the Effectiveness of Evolocumab Versus LDL Apheresis in Patients With Hypercholesterolemia (<https://clinicaltrials.gov/ct2/show/NCT03429998>) (2018).
- [51]. Havenon A de. PCSK9 Inhibition in Patients With Symptomatic Intracranial Atherosclerosis <https://clinicaltrials.gov/ct2/show/NCT03507374> (2020).
- [52]. Johns Hopkins University. Effect of Evolocumab on Coronary Endothelial Function (<https://clinicaltrials.gov/ct2/show/NCT03500302>) (2020).
- [53]. Johns Hopkins University. Evolocumab in Acute Coronary Syndrome: A Double-Blind Randomized Placebo Controlled Study (<https://clinicaltrials.gov/ct2/show/NCT03515304>) (2022).
- [54]. LAPLACE-TIMI 57: Low-density Lipoprotein Cholesterol (LDL-C) Assessment With PCSK9 monoclonal Antibody Inhibition Combined With Statin therapy - Full Text View - ClinicalTrials.gov (<https://clinicaltrials.gov/ct2/show/NCT01380730>).
- [55]. MD NL Assessment of Atherosclerotic Plaque Characteristics Change by DCE-MRI With Alirocumab - a Pilot Study (<https://clinicaltrials.gov/ct2/show/NCT02992301>) (2019).
- [56]. Minneapolis Heart Institute Foundation. Effect of Alirocumab on Saphenous Vein Graft Atherosclerosis: The Alirocumab for Stopping Atherosclerosis Progression in Saphenous Vein Grafts (ASAP-SVG) Pilot Trial (<https://clinicaltrials.gov/ct2/show/NCT03542110>) (2021).
- [57]. Minno MD Subclinical Atherosclerosis in Patients With Familial Hypercholesterolemia Treated With Evolocumab@ <https://clinicaltrials.gov/ct2/show/NCT04313270> (2021).
- [58]. Novartis Pharmaceuticals. An Open Label, Active Comparator Extension Trial to Assess the Effect of Long Term Dosing of Inclisiran and Evolocumab Given as Subcutaneous Injections in Participants With High Cardiovascular Risk and Elevated LDL-C (ORION-3) <https://clinicaltrials.gov/ct2/show/NCT03060577> (2022).
- [59]. Radboud University Medical Center. Functional Improvement of Non-infarct related Coronary Artery Stenosis by Extensive LDL-C Reduction With a PCSK9 Antibody (<https://clinicaltrials.gov/ct2/show/NCT04141579>) (2022).
- [60]. Regeneron Pharmaceuticals. A Randomized, Double-Blind Study of the Efficacy and Safety of Alirocumab Added on to Atorvastatin Versus Ezetimibe Added on to Atorvastatin Versus Atorvastatin Dose Increase Versus Switch to Rosuvastatin in Patients Who Are Not Controlled on Atorvastatin (<https://clinicaltrials.gov/ct2/show/NCT01730040>) (2015).

- [61]. Regeneron Pharmaceuticals. A Randomized, Double-Blind Study of the Efficacy and Safety of REGN727 Added-on to Rosuvastatin Versus Ezetimibe Added-on to Rosuvastatin Versus Rosuvastatin Dose Increase in Patients Who Are Not Controlled on Rosuvastatin (<https://clinicaltrials.gov/ct2/show/NCT01730053>) (2020).
- [62]. Regeneron Pharmaceuticals. A Randomized, Double-Blind, Double-Dummy, Active-Controlled Study to Evaluate the Efficacy and Safety of REGN727/SAR236553 in Patients With Primary Hypercholesterolemia Who Are Intolerant to Statins (<https://clinicaltrials.gov/ct2/show/NCT01709513>) (2020).
- [63]. Regeneron Pharmaceuticals. A Randomized, Double-Blind, Placebo-Controlled Study to Evaluate the Efficacy and Safety of an Every Four Weeks Treatment Regimen of Alirocumab in Patients With Primary Hypercholesterolemia (<https://clinicaltrials.gov/ct2/show/NCT01926782>) (2017).
- [64]. Regeneron Pharmaceuticals. A Randomized, Double-Blind, Placebo-Controlled, Parallel-Group Study to Evaluate the Efficacy and Safety of Alirocumab in Patients With Homozygous Familial Hypercholesterolemia (<https://clinicaltrials.gov/ct2/show/NCT03156621>) (2021).
- [65]. Regeneron Pharmaceuticals. A Randomized, Double-Blind, Placebo-Controlled, Parallel-Group Study to Evaluate the Efficacy and Safety of REGN727/SAR236553 in Patients With Heterozygous Familial Hypercholesterolemia Not Adequately Controlled With Their Lipid-Modifying Therapy (<https://clinicaltrials.gov/ct2/show/NCT01709500>) (2015).
- [66]. Sanofi A. Multicenter, Randomized, Double-blind, Parallel-group, Placebo-controlled Study Evaluating the Efficacy and Safety of Three Doses of SAR236553 (REGN727) Over 12 Weeks in Patients With Primary Hypercholesterolemia and LDL-cholesterol 100 mg/dL (2.59 mmol/L) on Ongoing Stable Atorvastatin Therapy (<https://clinicaltrials.gov/ct2/show/NCT01812707>) (2016).
- [67]. Sanofi. A Randomized, Double-Blind, Active-Controlled, Parallel-Group Study to Evaluate the Efficacy And Safety of SAR236553/REGN727 Over 24 Weeks in Patients With Hypercholesterolemia (<https://clinicaltrials.gov/ct2/show/NCT01644474>) (2015).
- [68]. Sanofi. A Randomized, Double-blind, Parallel Group Study to Evaluate the Efficacy and Safety of Alirocumab (SAR236553/REGN727) Versus Ezetimibe in Asia in High Cardiovascular Risk Patients With Hypercholesterolemia Not Adequately Controlled With Their Statin Therapy (<https://clinicaltrials.gov/ct2/show/NCT02715726>) (2019).
- [69]. Sanofi. A Randomized, Double-Blind, Parallel Group Study to Evaluate the Efficacy and Safety of SAR236553/REGN727 Versus Ezetimibe in High Cardiovascular Risk Patients With Hypercholesterolemia Not Adequately Controlled With Their Statin Therapy (<https://clinicaltrials.gov/ct2/show/NCT01644188>) (2016).
- [70]. Sanofi. A Randomized, Double-Blind, Placebo-Controlled, Ascending Single-Dose Study of the Safety, Tolerability, Pharmacokinetics and Pharmacodynamics of Subcutaneously Administered Alirocumab in Chinese Healthy Subjects (<https://clinicaltrials.gov/ct2/show/NCT02979015>) (2017).
- [71]. Sanofi. A Randomized, Double-Blind, Placebo-Controlled, Parallel Group Study to Evaluate the Efficacy and Safety of Alirocumab in High Cardiovascular Risk Patients With Hypercholesterolemia Not Adequately Controlled With Their Lipid-Modifying Therapy in South Korea and Taiwan (<https://clinicaltrials.gov/ct2/show/NCT02289963>) (2017).
- [72]. Sanofi. A Randomized, Double-blind, Placebo-controlled, Parallel Group Study to Evaluate the Efficacy and Safety of Alirocumab in Patients With Hypercholesterolemia Not Adequately Controlled With Non-statin Lipid Modifying Therapy or the Lowest Strength of Statin (<https://clinicaltrials.gov/ct2/show/NCT02584504>) (2019).
- [73]. Sanofi. A Randomized, Double-Blind, Placebo-Controlled, Parallel Group Study to Evaluate the Efficacy and Safety of SAR236553/REGN727 in High Cardiovascular Risk Patients With Hypercholesterolemia Not Adequately Controlled With Their Lipid-Modifying Therapy (<https://clinicaltrials.gov/ct2/show/NCT01644175>) (2015).
- [74]. Sanofi. A Randomized, Double-Blind, Placebo-Controlled, Parallel Group Study to Evaluate the Efficacy and Safety of SAR236553/REGN727 in Patients With Heterozygous Familial Hypercholesterolemia and LDL-C Higher or Equal to 160mg/dL With Their Lipid-Modifying Therapy (<https://clinicaltrials.gov/ct2/show/NCT01617655>) (2016).

- [75]. Sanofi. A Randomized, Double-Blind, Placebo-Controlled, Parallel Group Study to Evaluate the Efficacy and Safety of SAR236553/REGN727 in Patients With Heterozygous Familial Hypercholesterolemia Not Adequately Controlled With Their Lipid-Modifying Therapy (<https://clinicaltrials.gov/ct2/show/NCT01623115>) (2016).
- [76]. Sanofi. A Randomized, Double-Blind, Placebo-Controlled, Parallel-Group Study Evaluating the Efficacy and Safety of Alirocumab in Patients With Primary Hypercholesterolemia Not Treated With a Statin (<https://clinicaltrials.gov/ct2/show/NCT02023879>) (2018).
- [77]. Sanofi. A Randomized, Double-Blind, Placebo-Controlled, Parallel-Group Study to Evaluate the Effect of Alirocumab (SAR236553/REGN727) on the Occurrence of Cardiovascular Events in Patients Who Have Recently Experienced an Acute Coronary Syndrome (<https://clinicaltrials.gov/ct2/show/NCT01663402>) (2019).
- [78]. Sanofi. A Randomized, Open-label, Blinded Intravascular Ultrasound Analysis, Parallel Group, Multicenter Study to Evaluate the Effect of Praluent® (Alirocumab) on Coronary Atheroma Volume in Japanese Patients Hospitalized for Acute Coronary Syndrome With Hypercholesterolemia Not Adequately Controlled With Statin (<https://clinicaltrials.gov/ct2/show/NCT02984982>) (2019).
- [79]. Sanofi. A Phase 1 Study of the Effects of Subcutaneous Doses of Alirocumab on Lipid and Lipoprotein Metabolism in Adults With Mildly Elevated LDL-Cholesterol (<https://clinicaltrials.gov/ct2/show/NCT01959971>) (2017).
- [80]. Sanofi. An 8-Week Open-Label, Sequential, Repeated Dose-Finding Study to Evaluate the Efficacy and Safety of Alirocumab in Children and Adolescents With Heterozygous Familial Hypercholesterolemia Followed by an Extension Phase (<https://clinicaltrials.gov/ct2/show/NCT02890992>) (2019).
- [81]. Sanofi. Long-term Safety and Tolerability of SAR236553 (REGN727) in High Cardiovascular Risk Patients With Hypercholesterolemia Not Adequately Controlled With Their Lipid Modifying Therapy: A Randomized, Double-Blind, Placebo-Controlled Study (<https://clinicaltrials.gov/ct2/show/NCT01507831>) (2015).
- [82]. Sanofi. Open-Label Extension Study of EFC12492, R727-CL-1112, EFC12732 and LTS11717 Studies to Assess the Long-Term Safety and Efficacy of Alirocumab in Patients With Heterozygous Familial Hypercholesterolemia (<https://clinicaltrials.gov/ct2/show/NCT01954394>) (2018).
- [83]. University Hospital Inselspital, Berne. Effects of the PCSK9 Antibody AliroCuMab on Coronary Atherosclerosis in Patients With Acute Myocardial Infarction: A Serial, Multivessel, Intravascular Ultrasound, Near-Infrared Spectroscopy And Optical Coherence Tomography Imaging Study (<https://clinicaltrials.gov/ct2/show/NCT03067844>) (2022).
- [84]. University Hospital Inselspital, Berne. EVOlocumab for Early Reduction of LDL-cholesterol Levels in Patients With Acute Coronary Syndromes (EVOPACS) - A Randomized, Double-blind, Placebo-controlled Multicenter Study (<https://clinicaltrials.gov/ct2/show/NCT03287609>) (2019).
- [85]. Washington University School of Medicine. Effect of Alirocumab on Reverse Cholesterol Transport in Humans (<https://clinicaltrials.gov/ct2/show/NCT03014830>) (2018).
- [86]. Ranade P High Cholesterol: PCSK9s. ICER (<https://icer.org/assessment/high-cholesterol-2015/>).
- [87]. Shin MD, et al. , COVID-19 vaccine development and a potential nanomaterial path forward, Nat. Nanotechnol 15 (2020) 646–655. [PubMed: 32669664]
- [88]. Chackerian B, Fietze KM, Moving towards a new class of vaccines for non-infectious chronic diseases, Expert Rev. Vaccin 15 (2016) 561–563.
- [89]. Momtazi-Borojeni AA, Jaafari MR, Badiie A, Sahebkar A, Long-term generation of antiPCSK9 antibody using a nanoliposome-based vaccine delivery system, Atherosclerosis 283 (2019) 69–78. [PubMed: 30797988]
- [90]. Ortega-Rivera OA, Pokorski JK, Steinmetz NF, A single-dose, implant-based, trivalent virus-like particle vaccine against “cholesterol checkpoint” proteins, Adv. Ther 4 (2021) 2100014.
- [91]. Walter U, Santamaria P, CD8+ T cells in autoimmunity, Curr. Opin. Immunol 17 (2005) 624–631. [PubMed: 16226438]
- [92]. Chyu K-Y, et al. , CD8+ T cells mediate the athero-protective effect of immunization with an apob-100 peptide, PLOS ONE 7 (2012) e30780. [PubMed: 22347402]

- [93]. Chyu K-Y, et al. , Immunization using ApoB-100 peptide-linked nanoparticles reduces atherosclerosis, *JCI Insight* 7 (2022).
- [94]. Musunuru K, et al. , In vivo CRISPR base editing of PCSK9 durably lowers cholesterol in primates, *Nature* 593 (2021) 429–434. [PubMed: 34012082]
- [95]. Zhou X, Caligiuri G, Hamsten A, Lefvert AK, Hansson GK, LDL immunization induces T-cell-dependent antibody formation and protection against atherosclerosis, *Arterioscler. Thromb. Vasc. Biol* 21 (2001) 108–114. [PubMed: 11145941]
- [96]. Chyu K-Y, et al. , Timing affects the efficacy of LDL immunization on atherosclerotic lesions in apo E (–/–) mice, *Atherosclerosis* 176 (2004) 27–35. [PubMed: 15306171]
- [97]. Zhou X, Robertson A-KL, Rudling M, Parini P, Hansson GK, Lesion development and response to immunization reveal a complex role for CD4 in atherosclerosis, *Circ. Res* 96 (2005) 427–434. [PubMed: 15662027]
- [98]. van Puijvelde GHM, et al. , Induction of oral tolerance to oxidized low-density lipoprotein ameliorates atherosclerosis, *Circulation* 114 (2006) 1968–1976. [PubMed: 17060383]
- [99]. Asgary S, Saberi S-A, Azampanah S, Effect of immunization against ox-LDL with two different antigens on formation and development of atherosclerosis, *Lipids Health Dis* 6 (2007) 32. [PubMed: 18036243]
- [100]. Habets KLL, et al. , Vaccination using oxidized low-density lipoprotein-pulsed dendritic cells reduces atherosclerosis in LDL receptor-deficient mice, *Cardiovasc. Res* 85 (2010) 622–630. [PubMed: 19819882]
- [101]. Zhong Y, et al. , CD4+LAP + and CD4 +CD25 +Foxp3 + regulatory T cells induced by nasal oxidized low-density lipoprotein suppress effector T cells response and attenuate atherosclerosis in ApoE–/– mice, *J. Clin. Immunol* 32 (2012) 1104–1117. [PubMed: 22552859]
- [102]. Martos-Folgado I, et al. , MDA-LDL vaccination induces athero-protective germinal-center-derived antibody responses, *Cell Rep* 41 (2022).
- [103]. Fredrikson GN, et al. , Inhibition of atherosclerosis in apoE-null mice by immunization with apoB-100 peptide sequences, *Arterioscler. Thromb. Vasc. Biol* 23 (2003) 879–884. [PubMed: 12649092]
- [104]. Chyu K-Y, et al. , Immunization using an Apo B-100 related epitope reduces atherosclerosis and plaque inflammation in hypercholesterolemic apo E (–/–) mice, *Biochem Biophys. Res Commun* 338 (2005) 1982–1989. [PubMed: 16288717]
- [105]. Fredrikson GN, et al. , Atheroprotective immunization with MDA-modified apo B-100 peptide sequences is associated with activation of Th2 specific antibody expression, *Autoimmunity* 38 (2005) 171–179. [PubMed: 16040338]
- [106]. Fredrikson GN, Björkbacka H, Söderberg I, Ljungcrantz I, Nilsson J, Treatment with apo B peptide vaccines inhibits atherosclerosis in human apo B-100 transgenic mice without inducing an increase in peptide-specific antibodies, *J. Intern Med* 264 (2008) 563–570. [PubMed: 18783480]
- [107]. Wigren M, et al. , Evidence for a role of regulatory T cells in mediating the atheroprotective effect of apolipoprotein B peptide vaccine, *J. Intern. Med* 269 (2011) 546–556. [PubMed: 21114693]
- [108]. Hermansson A, et al. , Immunotherapy With tolerogenic apolipoprotein B-100-loaded dendritic cells attenuates atherosclerosis in hypercholesterolemic mice, *Circulation* 123 (2011) 1083–1091. [PubMed: 21357823]
- [109]. Herbin O, et al. , Regulatory T-cell response to apolipoprotein B100-derived peptides reduces the development and progression of atherosclerosis in mice, *Arterioscler., Thromb., Vasc. Biol* 32 (2012) 605–612. [PubMed: 22223728]
- [110]. Tse K, et al. , Atheroprotective vaccination with MHC-II restricted peptides from ApoB-100, *Front Immunol* 4 (2013) 493. [PubMed: 24416033]
- [111]. Pierides C, Bermudez-Fajardo A, Fredrikson GN, Nilsson J, Oviedo-Orta E, Immune responses elicited by apoB-100-derived peptides in mice, *Immunol. Res* 56 (2013) 96–108. [PubMed: 23345063]

- [112]. Kimura T, et al. , Atheroprotective vaccination with MHC-II-restricted ApoB peptides induces peritoneal IL-10-producing CD4 T cells, *Am. J. Physiol. Heart Circ. Physiol* 312 (2017) H781–H790. [PubMed: 28087520]
- [113]. Kobiyama K, et al. , A clinically applicable adjuvant for an atherosclerosis vaccine, *Eur. J. Immunol* 48 (2018) 1580–1587. [PubMed: 29932463]
- [114]. Kimura T, et al. , Regulatory CD4+ T cells recognize MHC-II-restricted peptide epitopes of apolipoprotein B, *Circulation* 138 (2018) 1130–1143. [PubMed: 29588316]
- [115]. Rittershaus CW, et al. , Vaccine-induced antibodies inhibit CETP activity in vivo and reduce aortic lesions in a rabbit model of atherosclerosis, *Arterioscler., Thromb., Vasc. Biol* 20 (2000) 2106–2112. [PubMed: 10978256]
- [116]. Gaofu Q, et al. , Long-lasting specific antibodies against CETP induced by subcutaneous and mucosal administration of a 26-amino acid CETP epitope carried by heat shock protein 65 kDa in the absence of adjuvants, *Vaccine* 22 (2004) 3187–3194. [PubMed: 15297073]
- [117]. Gaofu Q, et al. , Vaccinating rabbits with a cholesteryl ester transfer protein (CETP) B-Cell epitope carried by heat shock protein-65 (HSP65) for inducing anti-CETP antibodies and reducing aortic lesions in vivo, *J. Cardiovasc Pharm* 45 (2005) 591–598.
- [118]. Gaofu Q, et al. , Antibody against cholesteryl ester transfer protein (CETP) elicited by a recombinant chimeric enzyme vaccine attenuated atherosclerosis in a rabbit model, *Life Sci* 77 (2005) 2690–2702. [PubMed: 15963532]
- [119]. Mao D, et al. , Intramuscular immunization with a DNA vaccine encoding a 26-amino acid CETP epitope displayed by HBc protein and containing CpG DNA inhibits atherosclerosis in a rabbit model of atherosclerosis, *Vaccine* 24 (2006) 4942–4950. [PubMed: 16697088]
- [120]. Yuan X, et al. , Intranasal immunization with chitosan/pCETP nanoparticles inhibits atherosclerosis in a rabbit model of atherosclerosis, *Vaccine* 26 (2008) 3727–3734. [PubMed: 18524427]
- [121]. Thomas LJ, et al. , Co-administration of a CpG adjuvant (VaxImmune, CPG 7909) with CETP vaccines increased immunogenicity in rabbits and mice, *Hum. Vaccin* 5 (2009) 79–84. [PubMed: 18670196]
- [122]. Qi G, et al. , A chimeric peptide of intestinal trefoil factor containing cholesteryl ester transfer protein B cell epitope significantly inhibits atherosclerosis in rabbits after oral administration, *Peptides* 32 (2011) 790–796. [PubMed: 21192999]
- [123]. Liaw Y-W, et al. , A vaccine targeted at CETP alleviates high fat and high cholesterol diet-induced atherosclerosis and non-alcoholic steatohepatitis in rabbit, *PLoS One* 9 (2014) e111529. [PubMed: 25486007]
- [124]. Aghebati T, et al. , Anti-atherosclerosis effect of different doses of CETP vaccine in rabbit model of atherosclerosis, *Biomed. Pharm* 81 (2016) 468–473.
- [125]. Aghebati T, et al. , A novel atheroprotective role of MF59-like adjuvant when co-administered with CETP vaccine in rabbit model of atherosclerosis, *Iran. J. Basic Med Sci* 19 (2016) 1345–1352. [PubMed: 28096968]
- [126]. Gutiérrez-Vidal R, et al. , Therapeutic intranasal vaccine HB-ATV-8 prevents atherogenesis and non-alcoholic fatty liver disease in a pig model of atherosclerosis, *Arch. Med. Res* 49 (2018) 456–470. [PubMed: 30792163]
- [127]. Aghebati T, et al. , Development of an effective liposomal cholesterol ester transfer protein (CETP) vaccine for protecting against atherosclerosis in rabbit model, *Pharm. Dev. Technol* 25 (2020) 432–439. [PubMed: 31852350]
- [128]. Chackerian B, Remaley A, Vaccine strategies for lowering LDL by immunization against proprotein convertase subtilisin/kexin type 9, *Curr. Opin. Lipido* 27 (2016) 345–350.
- [129]. Galabova G, et al. , Peptide-based anti-PCSK9 vaccines - an approach for long-term LDLc management, *PLoS One* 9 (2014) e114469. [PubMed: 25474576]
- [130]. Crossey E, et al. , A cholesterol-lowering VLP vaccine that targets PCSK9, *Vaccine* 33 (2015) 5747–5755. [PubMed: 26413878]
- [131]. Landlinger C, et al. , The AT04A vaccine against proprotein convertase subtilisin/kexin type 9 reduces total cholesterol, vascular inflammation, and atherosclerosis in APOE*3Leiden.CETP mice, *Eur. Heart J* 38 (2017) 2499–2507. [PubMed: 28637178]

- [132]. Pan Y, et al. , A therapeutic peptide vaccine against PCSK9, *Sci. Rep.* 7 (2017) 12534. [PubMed: 28970592]
- [133]. Kawakami R, et al. , Development of vaccine for dyslipidemia targeted to a proprotein convertase subtilisin/kexin type 9 (PCSK9) epitope in mice, *PLoS One* 13 (2018) e0191895. [PubMed: 29438441]
- [134]. Momtazi-Borojeni AA, Jaafari MR, Badiie A, Banach M, Sahebkar A, Therapeutic effect of nanoliposomal PCSK9 vaccine in a mouse model of atherosclerosis, *BMC Med* 17 (2019) 223. [PubMed: 31818299]
- [135]. Wu D, et al. , PCSK9Q β –003 vaccine attenuates atherosclerosis in apolipoprotein E-deficient mice, *Cardiovasc Drugs Ther* 35 (2021) 141–151. [PubMed: 32725442]
- [136]. Momtazi-Borojeni AA, Jaafari MR, Afshar M, Banach M, Sahebkar A, PCSK9 immunization using nanoliposomes: preventive efficacy against hypercholesterolemia and atherosclerosis, *Arch. Med Sci* 17 (2021) 1365–1377. [PubMed: 34522266]
- [137]. Momtazi-Borojeni AA, et al. , Pre-clinical evaluation of the nanoliposomal antiPCSK9 vaccine in healthy non-human primates, *Vaccines* 9 (2021) 749. [PubMed: 34358164]
- [138]. Maron R, et al. , Mucosal administration of heat shock protein-65 decreases atherosclerosis and inflammation in aortic arch of low-density lipoprotein receptor-deficient mice, *Circulation* 106 (2002) 1708–1715. [PubMed: 12270867]
- [139]. Harats D, Yacov N, Gilburd B, Shoenfeld Y, George J, Oral tolerance with heat shock protein 65 attenuates *Mycobacterium tuberculosis*-induced and high-fat-diet-driven atherosclerotic lesions, *J. Am. Coll. Cardiol* 40 (2002) 1333–1338. [PubMed: 12383583]
- [140]. van Puijvelde GHM, et al. , Induction of oral tolerance to HSP60 or an HSP60-peptide activates T cell regulation and reduces atherosclerosis, *Arterioscler. Thromb. Vasc. Biol* 27 (2007) 2677–2683. [PubMed: 17901374]
- [141]. Xiong Q, Li J, Jin L, Liu J, Li T, Nasal immunization with heat shock protein 65 attenuates atherosclerosis and reduces serum lipids in cholesterol-fed wild-type rabbits probably through different mechanisms, *Immunol. Lett* 125 (2009) 40–45. [PubMed: 19505506]
- [142]. Jing H, et al. , Oral administration of *Lactococcus lactis* delivered heat shock protein 65 attenuates atherosclerosis in low-density lipoprotein receptor-deficient mice, *Vaccine* 29 (2011) 4102–4109. [PubMed: 21497632]
- [143]. Long J, et al. , Nasal immunization with different forms of heat shock protein-65 reduced high-cholesterol-diet-driven rabbit atherosclerosis, *Int Immunopharmacol* 13 (2012) 82–87. [PubMed: 22465386]
- [144]. Li H, Ding Y, Yi G, Zeng Q, Yang W, Establishment of nasal tolerance to heat shock protein-60 alleviates atherosclerosis by inducing TGF- β -dependent regulatory T cells, *J. Huazhong Univ. Sci. Technol. Med Sci* 32 (2012) 24–30.
- [145]. Grundtman C, et al. , *Mycobacterial* heat shock protein 65 (mbHSP65)-induced atherosclerosis: Preventive oral tolerization and definition of atheroprotective and atherogenic mbHSP65 peptides, *Atherosclerosis* 242 (2015) 303–310. [PubMed: 26233917]
- [146]. Zhong Y, et al. , Intranasal immunization with heat shock protein 60 induces CD4(+) CD25(+) GARP(+) and type 1 regulatory T cells and inhibits early atherosclerosis, *Clin. Exp. Immunol* 183 (2016) 452–468. [PubMed: 26452441]
- [147]. Hu Y, et al. , Effects of oral and subcutaneous administration of HSP60 on myeloid-derived suppressor cells and atherosclerosis in ApoE $^{-/-}$ mice, *Biochem Biophys. Res Commun* 498 (2018) 701–706. [PubMed: 29107690]
- [148]. Wick C, Onestingel E, Demetz E, Dietrich H, Wick G, Oral tolerization with mycobacterial heat shock protein 65 reduces chronic experimental atherosclerosis in aged mice, *Gerontology* 64 (2018) 36–48. [PubMed: 28910785]
- [149]. Gardiner SM, et al. , Active immunization with angiotensin I peptide analogue vaccines selectively reduces the pressor effects of exogenous angiotensin I in conscious rats, *Br. J. Pharm* 129 (2000) 1178–1182.
- [150]. Downham MR, et al. , Evaluation of two carrier protein–angiotensin I conjugate vaccines to assess their future potential to control high blood pressure (hypertension) in man, *Br. J. Clin. Pharmacol* 56 (2003) 505–512. [PubMed: 14651724]

- [151]. Zhu F, et al. , Target organ protection from a novel angiotensin II receptor (AT1) vaccine ATR12181 in spontaneously hypertensive rats, *Mol. Immunol* 3 (2006) 8.
- [152]. Ambühl PM, et al. , A vaccine for hypertension based on virus-like particles: preclinical efficacy and phase I safety and immunogenicity, *J. Hypertens* 25 (2007) 63–72. [PubMed: 17143175]
- [153]. Hong F, et al. , A vaccine for hypertension based on peptide AngI-R: a pilot study, *Int. J. Cardiol* 148 (2011) 76–84. [PubMed: 19932924]
- [154]. Chen X, et al. , Effectiveness and safety of a therapeutic vaccine against angiotensin II receptor type 1 in hypertensive animals, *Hypertension* 61 (2013) 408–416. [PubMed: 23184378]
- [155]. Nakagami F, et al. , Decrease in blood pressure and regression of cardiovascular complications by angiotensin ii vaccine in mice, *PLoS One* 8 (2013) e60493. [PubMed: 23544146]
- [156]. Ou X, et al. , Construction, expression and immunogenicity of a novel anti-hypertension angiotensin II vaccine based on hepatitis A virus-like particle, *Hum. Vaccin. Immunother* 9 (2013) 1191–1199. [PubMed: 23412424]
- [157]. Li L-D, et al. , Effect of active immunization against angiotensin II type 1 (AT1) receptor on hypertension & arterial remodelling in spontaneously hypertensive rats (SHR), *Indian J. Med. Res* 139 (2014) 619–624. [PubMed: 24927350]
- [158]. Koriyama H, et al. , Long-term reduction of high blood pressure by angiotensin II DNA vaccine in spontaneously hypertensive rats, *Hypertension* 66 (2015) 167–174. [PubMed: 26015450]
- [159]. Zhou Y, et al. , ATRQ β -001 vaccine prevents atherosclerosis in apolipoprotein E-null mice, *J. Hypertens* 34 (2016) 474–485. [PubMed: 26771341]
- [160]. Azegami T, et al. , Intranasal vaccination against angiotensin II type 1 receptor and pneumococcal surface protein A attenuates hypertension and pneumococcal infection in rodents, *J. Hypertens* 36 (2018) 387–394. [PubMed: 28832363]
- [161]. Nakamaru R, et al. , A novel angiotensin II peptide vaccine without an adjuvant in mice, *J. Hypertens* 39 (2021) 181–189. [PubMed: 32667158]
- [162]. Lu X, et al. , Impact of multiple antigenic epitopes from ApoB100, hHSP60 and *Chlamydomonas pneumoniae* on atherosclerotic lesion development in Apobtm2SgyLdlrtm1Her J mice, *Atherosclerosis* 225 (2012) 56–68. [PubMed: 22959702]
- [163]. Mundkur L, et al. , Mucosal tolerance to a combination of ApoB and HSP60 peptides controls plaque progression and stabilizes vulnerable plaque in Apobtm2SgyLdlrtm1Her/J mice, *PLOS ONE* 8 (2013) e58364. [PubMed: 23505495]
- [164]. Mundkur L, et al. , Oral dosing with multi-antigenic construct induces atheroprotective immune tolerance to individual peptides in mice, *Int. J. Cardiol* 175 (2014) 340–351. [PubMed: 24962340]
- [165]. Jun L, et al. , Effects of nasal immunization of multi-target preventive vaccines on atherosclerosis, *Vaccine* 30 (2012) 1029–1037. [PubMed: 22192848]
- [166]. Amirfakhryan H, Vaccination against atherosclerosis: an overview, *Hell. J. Cardiol* 61 (2020) 78–91.
- [167]. Podrez EA, Schmitt D, Hoff HF, Hazen SL, Myeloperoxidase-generated reactive nitrogen species convert LDL into an atherogenic form in vitro, *J. Clin. Invest* 103 (1999) 1547–1560. [PubMed: 10359564]
- [168]. Schiopu A, et al. , Recombinant human antibodies against aldehyde-modified apolipoprotein B-100 peptide sequences inhibit atherosclerosis, *Circulation* 110 (2004) 2047–2052. [PubMed: 15451805]
- [169]. Fredrikson GN, et al. , Identification of immune responses against aldehydemodified peptide sequences in ApoB associated with cardiovascular disease, *Arterioscler., Thromb., Vasc. Biol* 23 (2003) 872–878. [PubMed: 12649091]
- [170]. Roy P, Ali AJ, Kobiyama K, Ghosheh Y, Ley K, Opportunities for an atherosclerosis vaccine: from mice to humans, *Vaccine* 38 (2020) 4495–4506. [PubMed: 31964554]
- [171]. Schiopu A, et al. , Recombinant antibodies to an oxidized low-density lipoprotein epitope induce rapid regression of atherosclerosis in apobec-1(–/–)/low-density lipoprotein receptor(–/–) mice, *J. Am. Coll. Cardiol* 50 (2007) 2313–2318. [PubMed: 18068040]

- [172]. Barter PJ, Kastelein JJP, Targeting cholesteryl ester transfer protein for the prevention and management of cardiovascular disease, *J. Am. Coll. Cardiol* 47 (2006) 492–499. [PubMed: 16458126]
- [173]. Barter PJ, et al. , Cholesteryl ester transfer protein: a novel target for raising HDL and inhibiting atherosclerosis, *Arterioscler. Thromb. Vasc. Biol* 23 (2003) 160–167. [PubMed: 12588754]
- [174]. Affiris AG Safety, Immunogenicity and Dose Response of ATH03, a New Vaccine Against the Cholesterol Ester Transfer Protein (CETP), in Healthy Male Subjects With High Density Lipoprotein Cholesterol (HDLc) Blood Concentrations Equal or Below 80 mg/dl <https://clinicaltrials.gov/ct2/show/NCT01284582> (2012).
- [175]. Catapano AL, Pirillo A, Norata GD, New pharmacological approaches to target PCSK9, *Curr. Atheroscler. Rep* 22 (2020) 24. [PubMed: 32495301]
- [176]. Graham MJ, et al. , Antisense inhibition of proprotein convertase subtilisin/kexin type 9 reduces serum LDL in hyperlipidemic mice, *J. Lipid Res* 48 (2007) 763–767. [PubMed: 17242417]
- [177]. Rinaldi C, Wood MJA, Antisense oligonucleotides: the next frontier for treatment of neurological disorders, *Nat. Rev. Neurol* 14 (2018) 9–21. [PubMed: 29192260]
- [178]. Frank-Kamenetsky M, et al. , Therapeutic RNAi targeting PCSK9 acutely lowers plasma cholesterol in rodents and LDL cholesterol in nonhuman primates, *Proc. Natl. Acad. Sci. USA* 105 (2008) 11915–11920. [PubMed: 18695239]
- [179]. Fattori E, et al. , Immunization against proprotein convertase subtilisin-like/kexin type 9 lowers plasma LDL-cholesterol levels in mice, *J. Lipid Res* 53 (2012) 1654–1661. [PubMed: 22611251]
- [180]. Affiris AG A Single-blind Phase 1 Study Assessing the Safety, Immunogenicity and Low Density Lipoprotein Cholesterol (LDLc)-Lowering Activity of 2 Different Proprotein Convertase Subtilisin/Kexin Type 9 (PCSK9) Targeting AFFITOPE® Vaccines in Healthy Subjects <https://clinicaltrials.gov/ct2/show/NCT02508896> (2017).
- [181]. Xu Q, Role of heat shock proteins in atherosclerosis, *Arterioscler., Thromb., Vasc. Biol* 22 (2002) 1547–1559. [PubMed: 12377729]
- [182]. Duan Y, et al. , Heat shock protein 60 in cardiovascular physiology and diseases, *Front Mol. Biosci* 7 (2020) 73. [PubMed: 32426370]
- [183]. Grundtman C, Kreutmayer SB, Almanzar G, Wick MC, Wick G, Heat shock protein 60 and immune inflammatory responses in atherosclerosis, *Arterioscler. Thromb. Vasc. Biol* 31 (2011) 960–968. [PubMed: 21508342]
- [184]. Hagiwara M, et al. , Sublingual vaccine with GroEL attenuates atherosclerosis, *J. Dent. Res* 93 (2014) 382–387. [PubMed: 24554540]
- [185]. Lu X, et al. , Immunization with a combination of ApoB and HSP60 epitopes significantly reduces early atherosclerotic lesion in Apobtm2SgyLdlrtm1Her/J mice, *Atherosclerosis* 212 (2010) 472–480. [PubMed: 20609438]
- [186]. Bocksch L, et al. , C-terminal apolipoprotein E-derived peptide, Ep1.B, displays anti-atherogenic activity, *Atherosclerosis* 194 (2007) 116–124. [PubMed: 17126342]
- [187]. Caligiuri G, et al. , Phosphorylcholine-targeting immunization reduces atherosclerosis, *J. Am. Coll. Cardiol* 50 (2007) 540–546. [PubMed: 17678738]
- [188]. De Haro J, et al. , Attenuation of early atherosclerotic lesions by immunotolerance with β 2 glycoprotein I and the immunomodulatory effectors interleukin 2 and 10 in a murine model, *J. Vasc. Surg* 62 (2015) 1625–1631. [PubMed: 25041990]
- [189]. Dunér P, et al. , Immunization of apoE^{-/-} mice with aldehyde-modified fibronectin inhibits the development of atherosclerosis, *Cardiovasc Res* 91 (2011) 528–536. [PubMed: 21493703]
- [190]. van Es T, et al. , IL-15 aggravates atherosclerotic lesion development in LDL receptor deficient mice, *Vaccine* 29 (2011) 976–983. [PubMed: 21115056]
- [191]. Fowler A, Sampson M, Remaley AT, Chackerian B, A VLP-based vaccine targeting ANGPTL3 lowers plasma triglycerides in mice, *Vaccine* 39 (2021) 5780–5786. [PubMed: 34474934]
- [192]. Fukami H, et al. , Vaccine targeting ANGPTL3 ameliorates dyslipidemia and associated diseases in mouse models of obese dyslipidemia and familial hypercholesterolemia, *Cell Rep. Med* 2 (2021) 100446. [PubMed: 34841293]
- [193]. Hauer AD, et al. , Vaccination against VEGFR2 attenuates initiation and progression of atherosclerosis, *Arterioscler. Thromb. Vasc. Biol* 27 (2007) 2050–2057. [PubMed: 17600223]

- [194]. Hauerd A, et al. , Blockade of interleukin-12 function by protein vaccination attenuates atherosclerosis, *Circulation* 112 (2005) 1054–1062. [PubMed: 16103256]
- [195]. Koizumi Y, Kurita-Ochiai T, Oguchi S, Yamamoto M, Nasal immunization with *Porphyromonas gingivalis* outer membrane protein decreases *P. gingivalis*-induced atherosclerosis and inflammation in spontaneously hyperlipidemic mice, *Infect. Immun* 76 (2008) 2958–2965. [PubMed: 18426881]
- [196]. Lindquist IE, Crossey E, Amar M, Remaley A, Chackerian B, A vaccine-based strategy for reducing cardiovascular disease risk factors, *J. Am. Coll. Cardiol* 69 (2017).
- [197]. Lu X, et al. , Immunization with a combination of 2 peptides derived from the C5a receptor significantly reduces early atherosclerotic lesion in *Ldlr(tm1Her) Apob(tm2Sgy)* J mice, *Arterioscler. Thromb. Vasc. Biol* 32 (2012) 2358–2371. [PubMed: 22837469]
- [198]. Ortega-Rivera OA, Shin MD, Moreno-Gonzalez MA, Pokorski JK, Steinmetz NF, A single-dose Q β VLP vaccine against S100A9 protein reduces atherosclerosis in a preclinical model, *Adv. Ther* 5 (2022) 2200092.
- [199]. Tissot AC, et al. , A VLP-based vaccine against interleukin-1 α protects mice from atherosclerosis, *Eur. J. Immunol* 43 (2013) 716–722. [PubMed: 23254454]
- [200]. van Wanrooij EJA, et al. , Vaccination against CD99 inhibits atherogenesis in low-density lipoprotein receptor-deficient mice, *Cardiovasc Res* 78 (2008) 590–596. [PubMed: 18250143]
- [201]. Xia M, et al. , Immunization of *Chlamydia pneumoniae* (Cpn)-infected *apobtm2SgyLdlrtm1Her/J* mice with a combined peptide of Cpn significantly reduces atherosclerotic lesion development, *PLoS One* 8 (2013) e81056. [PubMed: 24349031]
- [202]. Borén J, Packard CJ, Taskinen M-R, The roles of ApoC-III on the metabolism of triglyceride-rich lipoproteins in humans, *Front. Endocrinol* 11 (2020).
- [203]. Alexander RW, Hypertension and the pathogenesis of atherosclerosis, *Hypertension* 25 (1995) 155–161. [PubMed: 7843763]
- [204]. Bourinbaiair AS, Jirathitikal V, Effect of oral immunization with pooled antigens derived from adipose tissue on atherosclerosis and obesity indices, *Vaccine* 28 (2010) 2763–2768. [PubMed: 20117273]
- [205]. Bourinbaiair AS, Jirathitikal V, Safety and efficacy trial of adipose-tissue derived oral preparation V-6 Immunitor (V-6): results of open-label, two-month, follow-up study, *Lipids Health Dis* 9 (2010) 14. [PubMed: 20122177]




Fig. 1. Lipid-related, environmental, and cell-dependent factors that influence atherosclerotic plaque formation

. LDL retention initiates atherosclerosis. The subendothelial accumulation of lipoproteins leads to the upregulation of adhesion molecules on the endothelial surface and the recruitment of monocytes to the lesion. Monocytes transmigrate into the subendothelial space and differentiate into macrophages in response to macrophage-colony stimulating factor (M-CSF) and granulocyte-macrophage colony stimulating factor (GM-CSF) produced by endothelial cells. Smooth muscle cells can also transdifferentiate into macrophage-like cells. Scavenger-receptor-mediated uptake of lipoproteins by macrophages leads to the formation of foam cells. Printed with permission from BioRender.




Fig. 2. The development of vaccines to treat atherosclerosis. (A) The number of patent registrations (top), academic publications (middle), and registered clinical trials (bottom) between the years 2000 and 2022. (B) Word clouds showing the frequency of different targets (percentage representation) and their rankings in patents, academic publications, and clinical trials (excludes monoclonal antibody therapies).




Fig. 3. Proatherogenic properties of CETP.

CETP acts as a proatherogenic protein by promoting the transfer of cholesteryl esters from the antiatherogenic HDL to the proatherogenic LDL and VLDL. As CETP concentration increases, there is a lipid imbalance between the lipoproteins which leads to atherogenesis. CETP, cholesteryl ester transfer protein; HDL, high-density lipoprotein; LDL, low-density lipoprotein; VLDL, very-low-density lipoprotein. Printed with permission from BioRender.




Fig. 4. The PCSK9 mechanism of action.

Under normal physiological conditions (top), LDL-C is removed from blood circulation by LDL-R on the surface of hepatocytes. LDL-C is degraded inside the hepatocyte, and LDL-R is recycled back to the cell surface where it can bind to more LDL-C. In the presence of PCSK9 (bottom), both LDL-C and PCSK9 bind to LDL-R. LDL-C and LDL-R undergo lysosomal degradation, whereas PCSK9 is recycled back to the cell surface where it can continue to deplete the receptors. PCSK9, proprotein convertase subtilisin/kexin type 9; LDL-C, low-density lipoprotein cholesterol; LDL-R, low-density lipoprotein receptor. Printed with permission from BioRender.




Fig. 5. Role of HSP60 in atherosclerosis autoimmunity.

HSP60 is present in the mitochondria of healthy endothelial cells. As these cells are presented with stress factors such as smoking, chronic infections, and plaque buildup, HSP60 gets over-expressed and released from the cell. The exHSP60 gets recognized by the immune system as foreign and anti-HSP60 lead to ADCC. The exHSP60 is also recognized by TLR4 which leads to the production of pro-inflammatory cytokines TNF α and IL-6. HSP60, heat shock protein 60; exHSP60, extracellular heat shock protein 60, ADCC, antibody-dependent cellular cytotoxicity; TLR4, toll-like receptor 4; TNF α , tumor necrosis factor alpha; IL-6, interleukin 6. Printed with permission from BioRender.

Table 1

Major antigens considered during the development of atherosclerosis vaccines. Red shading indicates antigens tested in clinical trials. The rest have only been tested in preclinical models.

Lipid-related antigen	Non-lipid-related antigen	Pathogen-related antigen
Low density lipoprotein(LDL)	Heat shock protein (HSP60/HSP65)	<i>Chlamydia pneumoniae</i>
ApoB-100-derived peptides	Angiotensin II	<i>Porphyromonas gingivalis</i>
Cholesteryl ester transfer protein (CETP)	Interleukin 1 α /12/15	
Proprotein convertase subtilisin/kexin 9 (PCSK9)-derived peptides	CD99	
ApoE-derived peptides	Fibronectin	
ApoC-III-derived peptides	Vascular endothelial growth factor receptor 2 (VEGFR2)	
Phosphocholine head group	Complement 5a protein (C5a)	
	T-cell receptor β variable 31-derived peptide (TRBV31)	

Table 2

Preclinical studies of atherosclerosis vaccines 2000–2022. UniProt ApoB (human): P04114. UniProt HSP60 (human): P10809. UniProt ApoB (mouse): E9Q414. UniProt CETP (human): P11597. UniProt tetanus toxin: P04958. UniProt PCSK9 (human): Q8NBP7. UniProt PCSK9 (mouse): Q80W65.

Lipid-Related Antigens							
Protein Target	Year	Antigens	Carrier	Adjuvant	Model	Results	Reference
Low density lipoprotein (LDL)	2001	Homologous atherosclerotic plaque homogenate/homologous MDA-LDL	N/A	Complete and incomplete Freund's adjuvant	<i>ApoE^{-/-}</i> mice	46% lesion size ↓, T-cell-dependent IgG ↑	[95]
	2004	Murine native LDL	N/A	Murine IL-12	<i>ApoE^{-/-}</i> mice	46% plaque size ↓, oxLDL IgG ↑	[96]
	2005	Homologous MDA-LDL	N/A	Complete and incomplete Freund's adjuvant	<i>ApoE^{-/-}</i> and CD4/ <i>ApoE^{-/-}</i> mice	41% lesion size ↓, MDA-LDL IgM and IgG ↑	[97]
	2006	oxLDL	N/A	Dimethyl dioctadecyl ammonium bromide	<i>Ldlr^{-/-}</i> mice	71.2% plaque size ↓, CD4 ⁺ CD25 ⁺ Foxp3 ⁺ T _{reg} ↑, TGFβ ↑	[98]
	2007	MDA-LDL or Cu-LDL	N/A	Freund's adjuvant	New Zealand white rabbits	Fatty streak lesions ↓, LDL antibodies ↑, LDL-cholesterol ↓	[99]
	2010	oxLDL-pulsed DCs	N/A	N/A	<i>Ldlr^{-/-}</i> mice	87% plaque size ↓, T cell-dependent Cuox-LDL IgG↑	[100]
	2012	Human copper-oxLDL	N/A	N/A	<i>ApoE^{-/-}</i> mice	47.6% lesion size ↓, CD4 ⁺ LAP ⁺ and CD4 ⁺ CD25 ⁺ Foxp3 ⁺ T _{reg} ↑, TGFβ↑	[101]
	2022	MDA-LDL	N/A	Complete Freund's adjuvant	<i>Ldlr^{-/-}</i> mice	atherosclerotic lesion area ↓, memory B cells ↑, germinal center B cells ↑, IgM/IgG1/IgG2b/IgG2c ↑	[102]
Apolipoprotein B 100 (ApoB-100)	2003	MDA modified human ApoB-100 peptide 143 (aa 2131–2150: IALDDAKINFEKLSQLQTY) and peptide 210 (aa 3136–3155: KTKQSFDSLVSQAQYKKNKH)	Cationized bovine serum albumin (cBSA)	Alum (aluminum hydroxide)	<i>ApoE^{-/-}</i> mice	60% atherosclerotic lesion area ↓, MDA peptides IgG ↑	[103]
	2005	Human ApoB-100 peptide-1 (aa: EEEMLENVSLVCPKDATTRFK) and human ApoB-100 peptide-2 (aa: ATRFKHLRKYYTYNEAESS)	N/A	Alum	<i>ApoE^{-/-}</i> mice	42% aortic atherosclerotic lesion area ↓, peptide-specific IgG and IgM ↑	[104]
	2005	MDA-modified human ApoB-100 peptide 45 (aa 688–707: IEIGLEGKGFEPTLEALFGK) and peptide 74 (aa 1123–1142: VISIPRLQAEARSEILAHWS)	cBSA	Alum	<i>ApoE^{-/-}</i> mice	48% and 31% atherosclerotic lesion area ↓, peptide-specific IgG ↑, Th2 shift	[105]
	2008	Native human ApoB-100 peptide 45 (aa 661–680: IEIGLEGKGFEPTLEALFGK)	cBSA	Alum	<i>Ldlr^{-/-}</i> / human apoB-100 transgenic mice	66% atherosclerotic lesion area ↓, native LDL and copper-ox LDL IgM ↑	[106]

Lipid-Related Antigens							
Protein Target	Year	Antigens	Carrier	Adjuvant	Model	Results	Reference
		and peptide 210 (aa 3136–3155: KTTKQSFDSL SVK A QYKKNKH)					
	2010	Human ApoB peptide 210 (aa 3136–3155: KTTKQSFDSL SVK A QYKKNKH)	N/A	CTB (B subunit of cholera toxin)	<i>ApoE</i> ^{-/-} mice	35% aortic lesion area ↓, peptide specific IgG ⁺ , Foxp3 mRNA ↑	[20]
	2011	Human ApoB peptide 210 (aa 3136–3155: KTTKQSFDSL SVK A QYKKNKH)	cBSA	Alum	<i>ApoE</i> ^{-/-} mice	37% atherosclerotic lesion area ↓ CD4 ⁺ CD25 ⁺ T cell and CD4 ⁺ CD25 ⁺ Foxp3, Treg↑	[107]
	2011	Human ApoB-100 and IL-10	Dendritic cells	N/A	huB100 ⁺ (tg) × <i>Ldlr</i> ^{-/-} mice	70% lesion area ↓, ApoB100 IgG ↑, CD4 ⁺ ↓, IFN γ ↓	[108]
	2012	Mixture of human ApoB-100 peptide 210 (aa 3136–3155: KTTKQSFDSL SVK A QYKKNKH), peptide 240 (aa 3586–3605: FPDLGQEV ALNANTKNQKIR), and MDA-P210	N/A	N/A	<i>ApoE</i> ^{-/-} mice	40% lesion area ↓, 30% CD4 ⁺ CD25 ⁺ Foxp3 ⁺ T _{reg} ↑, 45% T-cell ↓	[109]
	2012	Human ApoB100 peptide 210 (aa 3136–3155: KTTKQSFDSL SVK A QYKKNKH)	cBSA	Alum	<i>ApoE</i> ^{-/-} mice	57% atherosclerotic lesion area ↓, peptide-specific IgG and IgM ↑, CD8 ⁺ CD62L ⁺ and CD8 ⁺ CD25 ⁺ IL-10 ⁺ T cells ↑	[92]
	2013	Murine ApoB-100 (aa 3501–3516: SQEYSGSVANEANVYL) and murine ApoB-100 (aa 978–993: TGAYSNASSTESASY)	N/A	Complete and incomplete Freund's adjuvant	<i>ApoE</i> ^{-/-} mice	40% lesion size ↓, peptide-specific IgG ↑	[110]
	2013	Human ApoB-100-derived peptides P2 (amino acid sequence not specified), P45 (aa 661–680: IEIGLEKGFEP TLEALFGK) and P210 (aa 3136–3155: KTTKQSFDSL SVK A QYKKNKH)	BSA and dendritic cells	N/A	<i>ApoE</i> ^{-/-} mice	CD4 ⁺ CD25 ⁺ FoxP3 ⁺ T _{reg} ↑, peptide-specific IgG1 ↑, IL-10 ↑	[111]
	2017	Murine ApoB peptides P101 (aa 705–720: FGKQGFPPDSVKNALY, P102 (aa 441–456: TLXALSHAVNSYFDVD, and P103 (aa 3953–3968: LYYKEDKTSLSAAS)	N/A	Complete and incomplete Freund's adjuvant	<i>ApoE</i> ^{-/-} mice	40% atherosclerotic lesion area ↓, peptide-specific IgG1 and IgG2c ↑, FoxP3 ⁺ CD4 ⁺ T cells ↑	[112]
	2018	Murine ApoB peptide P6 (aa 978–993: TGAYSNASSTESASY)	N/A	Addavax	<i>ApoE</i> ^{-/-} mice	57% lesion area ↓	[113]
	2018	Murine ApoB peptide P18 (aa 3030–3044: SLFFSAQPFEITAST)	N/A	Complete and incomplete Freund's adjuvant	<i>ApoE</i> ^{-/-} mice	~35% lesion size ↓, CD25 ⁺ FoxP3 ⁺ CD4 ⁺ T _{reg} ↑, peptide-specific IgG1 and IgG2c ↑	[114]
	2022	Human ApoB100 peptide 210 (aa 3136–3155: KTTKQSFDSL SVK A QYKKNKH)	Peptide amphiphile micelles (PAMs)	N/A	<i>A2Kb-Tg ApoE</i> ^{-/-} mice	atherosclerotic lesion area ↓, CD4 + effector memory T cells ↓, CD8 + central memory T cells ↑, CD4 +CD25 +FoxP3 + T _{reg} ↑, LDL-C ↓	[93]
CETP	2000	Human CETP (aa 461–476: FGPEHLLVDLFLQSL) and tetanus toxin (aa 830–843: QYIKANSKFIGITE)	N/A	Complete and incomplete Freund's adjuvant	New Zealand white rabbits	39.6% aortic lesion area ↓, CETP antibodies ⁺ , CETP activity ↓, HDL ↑, LDL ↓	[115]

Lipid-Related Antigens							
Protein Target	Year	Antigens	Carrier	Adjuvant	Model	Results	Reference
	2004	Human CETP (aa 451–476: RDGFLLQLQMDFGFPEHLLVDLFLQSLs)	HSP65	N/A	C57BL/6 J mice	anti-CETP antibodies ↑	[116]
	2005	Human CETP (aa 448–476: IITRDGFLLLQMDFGFPEHLLVDLFLQSLs)	HSP65	Alum	New Zealand white rabbits	30.8% aortic lesion area ↓, anti-CETP antibodies ↑, HDL-C ↑, LDL-C ↓	[117]
	2005	Asparaginase (AnsB), tetanus toxin (aa 831–854: YIKANSKFIGITELKKLESKINKV), and human CETP (aa 448–476: IITRDGFLLLQMDFGFPEHLLVDLFLQSLs)	N/A	Alum	New Zealand white rabbits	47.6% atherosclerotic plaque area ↓, anti-CETP antibodies ↑, HDL-C ↑, LDL-C ↓	[118]
	2006	pCR-X8-HBc-CETP	hepatitis B core (HBc)	HBc	New Zealand white rabbits	80.6% aortic lesion area ↓, anti-CETP antibodies ↑, HDL-C ↑, LDL-C ↓	[119]
	2008	pCR-X8-HBc-CETP	Chitosan	HBc	New Zealand white rabbits	59.2% aortic lesion area ↓, anti-CETP IgG ↑, LDL-C ↓	[120]
	2009	Human CETP (aa 461–476: FGPEHLLVDLFLQSLs), tetanus toxin (aa 830–843: QYIKANSKFIGITE) and PADRE T cell epitope (aK-Cha-VAAWTLKAa)	N/A	Alhydrogel and VaxImmune	New Zealand white rabbits and BALB/c mice	anti-CETP antibodies ↑	[121]
	2011	Human CETP (aa 461–476: FGPEHLLVDLFLQSLs) and tetanus toxin (aa QYIKANSKFIGITE)	Human intestinal trefoil factor (TFF3)	N/A	New Zealand white rabbits	atherosclerotic lesion area ↓, anti-CETP antibodies ↑, plasma CETP activity ↓, HDL-C ↑, LDL-C ↓, TNFα and IFNγ ↓, IL-10 and TGFβ ↑	[122]
	2014	Fc-CETP6	N/A	Complete and incomplete Freund's adjuvant	New Zealand white rabbits	atherosclerotic lesion area ↓, anti-CETP antibodies ↑, CETP activity ↓, HDL-C ↑, ApoA-I ↑, ox-LDL ↓	[123]
	2016	Human CETP (aa 461–476: FGPEHLLVDLFLQSLs) and tetanus toxoid (aa 830–845: QYIKANSKFIGITELK)	N/A	Complete and incomplete Freund's adjuvant	New Zealand white rabbits	anti-CETP antibodies ↑, CETP activity ↓	[124]
	2016	Human CETP (aa 461–476: FGPEHLLVDLFLQSLs) and tetanus toxoid (aa CQYIKANSKFIGITE)	N/A	MF59	New Zealand white rabbits	atherosclerotic lesion area ↓, IFNγ ↑, IL-4 ↓	[125]
	2018	CETP (aa H486–S496)	Micellar nanoparticle	N/A	Pigs (large white × landrace)	atherosclerotic lesion area ↓, anti-CETP IgG ↑	[126]
	2020	Human CETP (aa 461–476: FGPEHLLVDLFLQSLs) and tetanus toxoid (830–845: QYIKANSKFIGITELK)	Liposome	CpG oligonucleotide	White rabbits	atherosclerotic lesion area ↓, anti-TT-CETP antibodies ↑, IL-4 ↓, IFNγ ↑	[127]
PCSK9	2012	Human PCSK9 protein	N/A	CpG oligonucleotide	BALB/c mice	anti-hPCSK9 antibodies ↑, total cholesterol and HDL-C and LDL-C ↓, plasma mPCSK9 ↓, LDLR ↑	[128]
	2014	PCSK9 peptides (amino acid sequence not specified)	KLH	Alhydrogel	BALB/c mice	anti-hPCSK9 and anti-mPCSK9 antibodies ↑, total cholesterol ↓, LDLR ↑, LDI ↑, LDI ↓	[129]

Lipid-Related Antigens							
Protein Target	Year	Antigens	Carrier	Adjuvant	Model	Results	Reference
	2015	Human PCSK9 (aa 207–223; NVPEEDGTRFHRQASKC)	Qβ VLPs	Qβ VLPs	BALB/c mice and macaque monkeys	peptide-specific and recombinant human PCSK9 IgG ↑, 55% total cholesterol ↓, 50% free PCSK9 ↓, 15% LDL-C ↓	[130]
	2017	Human and mouse PCSK9 protein (aa 153–692)	N/A	N/A	APOE*3Leiden.CETP transgenic mice	64% lesion size ↓, PCSK9 antibodies ↑, plasma muPCSK9 ↓, 53% total cholesterol ↓, LDL-C ↓	[131]
	2017	PCSK9 peptide V150–157 (aa: FAQSIPWN)	Aluminum hydroxide gel	Qb VLP	<i>Ldlr</i> ^{−/−} mice	30% free PCSK9 ↓, cholesterol and LDL-C ↓, LDLR ↑	[132]
	2018	Mouse PCSK9 antigen (aa 682–690: CRSRPSAKA)	KLH	complete or incomplete Freund's adjuvant	<i>ApoE</i> ^{−/−} mice	Anti-PCSK9 antibody ↑, LDLR ↑, total cholesterol ↓	[133]
	2019	PCSK9 (aa: SIPWNLERITPVR) and tetanus toxin (aa: AQYIKANSKFIGITEL)	Nanoliposome	Alum	BALB/c mice	PCSK9 peptide IgG ↑, 52.2% plasma PCSK9 ↓, 50.2% PCSK9-LDLR binding ↓	[89]
	2019	PCSK9 (aa: SIPWNLERITPVR) and tetanus toxin (aa: AQYIKANSKFIGITEL)	Nanoliposome	Alum	C57BL/6 mice	39.13% lesion size ↓, PCSK9 peptide IgG ↑, 58.5% plasma PCSK9 ↓, 44.8% PCSK9-LDLR binding ↓, 44.7% total cholesterol ↓, 51.7% LDL-C ↓, LDLR ↑	[134]
	2020	PCSK9–003 peptide (aa: FAQSIPWN)	Qb VLP	Qb VLP	<i>ApoE</i> ^{−/−} mice	aortic lesion area ↓, peptide-specific antibody ↑, total cholesterol and LDL-C ↓, LDLR ↑, macrophage infiltration ↓	[135]
	2021	PCSK9 (aa: SIPWNLERITPVR) and tetanus toxin (aa: AQYIKANSKFIGITEL)	Nanoliposome	Alum	C57BL/6 mice	24.25% lesion size ↓, PCSK9 peptide IgG ↑, 40% plasma PCSK9 ↓, 69.86% PCSK9-LDLR binding ↓, 38.13% total cholesterol ↓, 57% LDL-C ↓, LDLR ↑	[136]
	2021	PCSK9 (aa: SIPWNLERITPVR) and tetanus toxin (aa: AQYIKANSKFIGITEL)	Nanoliposome	Alum	Rhesus macaque monkeys	PCSK9 IgG ↑, 33% PCSK9-LDLR binding ↓	[137]
Non-Lipid Related Antigens							
HSP60/65	2002	<i>Mycobacterium</i> HSP-65	N/A	N/A	<i>Ldlr</i> ^{−/−} mice	lesion size ↓, anti-HSP antibody ↓, macrophages and CD4 ⁺ T cells and IFNγ ↓, IL-10 ↑	[138]
	2002	<i>Mycobacterium</i> HSP-65	N/A	N/A	<i>Ldlr</i> ^{−/−} mice	lesion size ↓, IL-4 ↑	[139]
	2007	<i>Mycobacterium</i> HSP60 and HSP60 peptide (aa 253–268: SIQSIVPALEIANHR)	N/A	N/A	<i>Ldlr</i> ^{−/−} mice	83.3% plaque size ↓, CD4 ⁺ CD25 ⁺ Foxp3 ⁺ T _{reg} ↑, TGFβ and IL-10 ↑	[140]

Lipid-Related Antigens							
Protein Target	Year	Antigens	Carrier	Adjuvant	Model	Results	Reference
Angiotensin	2009	<i>Mycobacterium</i> HSP-65	N/A	CTB	New Zealand white rabbits	atherosclerotic lesion area ↓, IL-10 ↑, total cholesterol and LDL-C and HDL-C ↓,	[141]
	2011	<i>Mycobacterium</i> HSP-65	<i>Lactococcus lactis</i>	N/A	<i>Ldlr</i> ^{-/-} mice	39.83% atherosclerotic lesion area ↓, IFN γ ↓, IL-10 ↑	[142]
	2012	DNA encoding Hsp65 (Hsp65 DNA) and <i>Mycobacterium tuberculosis</i> Hsp65 protein	N/A	N/A	New Zealand White rabbits	lesion area ↓, anti-Hsp65 IgG and IgA ↑, IFN γ ↓, IL-10 ↑, total cholesterol and LDL-C ↓	[143]
	2012	recombinant H. Pylori HSP60	N/A	N/A	<i>ApoE</i> ^{-/-} mice	53.7% atherosclerotic plaque size ↓, CD4 ⁺ CD25 ⁺ LAP ⁺ and CD4 ⁺ CD25 ⁺ Foxp3 ⁺ T _{reg} ↑, 50% IFN γ ↓, TGF β ↑	[144]
	2015	Recombinant mycobacterial HSP65	N/A	N/A	<i>ApoE</i> ^{-/-} mice	lesion size ↓, splenic T _{reg} ↑, IL-10 ↑	[145]
	2016	Recombinant <i>Helicobacter pylori</i> HSP60	N/A	N/A	<i>ApoE</i> ^{-/-} mice	30.6% atherosclerotic plaque size ↓, CD4 ⁺ CD25 ⁺ GARP ⁺ and CD4 ⁺ CD25 ⁺ FoxP3 ⁺ T _{reg} ↑, IFN γ ↓, IL-10 and TGF β ↑	[146]
	2018	HSP-60 (organism not specified)	N/A	N/A	<i>ApoE</i> ^{-/-} mice	atherosclerotic plaque size ↓, MDSCs ⁺ , IFN γ ↓, IL-10 ⁺	[147]
	2018	Recombinant mycobacterial HSP65	N/A	incomplete Freund's adjuvant	<i>ApoE</i> ^{-/-} mice	atherosclerotic lesion area ↓, total cholesterol ↓, T _{reg} ↑	[148]
	2000	Angiotensin I (AngI) and Angiotensin II (AngII) analogues	Tetanus toxoid	Alum	Sprague-Dawley rats	anti-Ang IgG ↑, AngI response ↓	[149]
	2003	Angiotensin I (AngI) analogue	KLH	Alum	Sprague-Dawley rats	anti-AngI IgG and IgM ↑, blood pressure ↓	[150]
	2006	Angiotensin II type I receptor peptide ATR12181 (aa: AFHYESR)	Tetanus toxoid	Complete and incomplete Freund's adjuvant	Spontaneously hypertensive Rats	peptide-specific antibody ↑, blood pressure ↓	[151]
	2007	Angiotensin II (aa 1–8: DRVYIHPF)	Q β VLP	Q β VLP	Spontaneously hypertensive Rats	anti-AngII IgG ↑, blood pressure ↓, AngII ↑ (antibody-bound AngII)	[152]
	2011	Angiotensin I (aa: DRVYIHPFSL)	N/A	Alum	Spontaneously hypertensive Rats	anti-AngI-R and anti-AngII antibody ↑, blood pressure ↓, AngI and AngII ↓	[153]
	2013	Angiotensin II type I receptor peptide ATR-001 (aa: CAFHYESQ)	Q β VLP	Q β VLP	Spontaneously hypertensive Rats	peptide-specific antibody ↑, blood pressure ↓, AT ₁ R ↓	[154]
	2013	Angiotensin II (peptide not specified)	KLH	Complete and incomplete Freund's adjuvant	Spontaneously hypertensive Rats	anti-Ang II antibodies ↑, blood pressure ↓, IFN γ and IL-4 ↑	[155]

Lipid-Related Antigens						
Protein Target	Year	Antigens	Carrier	Adjuvant	Model	Results
	2013	pHAV-4Ang IIs	HAVLP	HAVLP	Spontaneously hypertensive Rats	anti-AngII IgG ↑, blood pressure ↓, Ang II ↓ [156]
	2014	Angiotensin II type 1 receptor peptides ATR12181 (aa: AFHYESR), ATR12185 (aa: ESRNSTL), and ATR 10014 (aa: IQDDCPK)	Tetanus toxoid	Complete and incomplete Freund's adjuvant	Spontaneously hypertensive Rats	peptide-specific antibody ↑, ATR12181 blood pressure ↓ [157]
	2015	pcDNA3.1-HBc-Ang II	HBc	HBc	Spontaneously hypertensive Rats	anti-Ang II antibody ↑, blood pressure ↓, AngII ↓, IFNγ and IL-2 ↑ [158]
	2016	Angiotensin II type 1 receptor peptide (aa: CAFHYESQ)	Qβ VLP	Qβ VLP	<i>ApoE</i> ^{-/-} mice	28% plaque area ↓, AT1R ↓, macrophages ↓ [159]
	2018	Angiotensin II type 1 receptor peptide (GenBank accession no. NM_030985; amino acids 181–187)	Pneumococcal surface protein A (PspA)	Cyclic diguanylate monophosphate (di-GMP)	Spontaneously hypertensive Rats	anti-AT1R IgG ↑, blood pressure ↓, AngII-AT1R binding ↓ [160]
	2021	AJP001 (aa: ELKLIFLHRLKRLKRLK) – angiotensin II	N/A	N/A	BALB/c	anti-Ang II antibody ↑, blood pressure ↓, IFNγ and IL-4 and IL-10 ↑, AngII ↑ [161]
Combination	2010	ApoB-100 (aa 661–680: IEIGLEKGFEPTLEALFGK) and human HSP60 (aa 153–163: AELKKQSKPVT)	N/A	Keyhole limpet hemocyanin (KLH)	ApoB ^Δ (tm2Sgy)/LDLr ^Δ (tm1Her/J) mice	41.3% atherosclerotic lesion area ↓, peptide-specific IgG ↑, macrophage and CD4 ⁺ and dendritic cell ↓, IFNγ and TNFα ↓, IL-10 ↑ 167
	2012	Human ApoB peptide (aa 688–707: IEIGLEKGFEPTLEALFGK), human HSP60 peptide (aa 153–163: AELKKQSKPVT), human HSP60 peptide (aa 303–312: PGFGDNRRNQ), and <i>Chlamydia pneumoniae</i> peptides (aa 67–74: GDYVFDRI and aa 283–291: QAVANGGAI)	N/A	Alum	ApoB ^Δ (tm2Sgy)/LDLr ^Δ (tm1Her) J mice	63.8% atherosclerotic lesion area ↓, peptide-specific IgG ↑, CD4 ⁺ ↑ [162]
	2013	ApoB-100 (aa 661–680: IEIGLEKGFEPTLEALFGK) and human HSP60 (aa 153–163: AELKKQSKPVT)	N/A	KLH	ApoB ^Δ (tm2Sgy)/LDLr ^Δ (tm1Her) mice	39.9% atherosclerotic lesion area ↓, T-regulatory cells and macrophage ↑, CD4 ⁺ [163]
	2014	Human ApoB-100 (aa 688–707: IEIGLEKGFEPTLEALFGK), human HSP60 (aa 153–163: AELKKQSKPVT), and <i>Chlamydia</i> peptides (aa 67–74: GDYVFDRI and aa 284–292: QAVANGGAI)	N/A	N/A	ApoB ^Δ (tm2Sgy) / LDLr ^Δ (tm1Her/J) mice	CTL A4 ⁺ and CD4 ⁺ CD25 ⁺ Foxp3 ⁺ T _{reg} ↑, 46.5% lesion area ↓, CD4 ⁺ CD25 ⁺ Foxp3 ⁺ T _{reg} ↑, CD11c ⁺ cells ↑ [164]
ApoB-100, CETP, and PCSK9	2021	Human ApoB (aa 3163–3182: KTTKQSFDSLVSVAQYKKKKH), Human CETP (aa 461–476: FGFEHLVDLQSL), and Human PCSK9 (aa 207–223: NVPEEDGTRFHRQASKC)	Qβ VLP	Qβ VLP	C57BL/6 J mice	peptide-specific IgG ↑, CETP and ApoB and PCSK9 ↓, total cholesterol ↓ [90]

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Lipid-Related Antigens							
Protein Target	Year	Antigens	Carrier	Adjuvant	Model	Results	Reference
CETP and HSP60	2012	Hsp65 DNA, Hsp65-CETP-PADRE-TT-CETP (HCPTC protein), or Hsp65 DNA + HCPTC protein	N/A	N/A	New Zealand white rabbits	aortic lesion area ↓, anti-CETP and anti-HSP65 antibodies ↑, LDL-C ↓, IFNγ ↓, IL-10 ↑	[165]