Self-Adaptive Kernels as Evolving Measures: Transport Adaptive Control

Mathematical Foundations of Self-Adaptive Kernels, version 2: a complete mathematical treatment

Delhi Technological University

October 2026

Version 2. This document supersedes version 1, *Mathematical Foundations of Self-Adaptive Kernels Achieving Lyapunov Convergence*. Section 2 states what survived from version 1 and what was withdrawn. This is research in progress and has not been peer reviewed. Code, tests, experiments and review records: https://github.com/Dutta-06/Lyapunov-Convergence-Kernels-Showcase.

Copyright © 2026 Odwitiyo Dutta. All rights reserved.

Abstract

This is a complete mathematical treatment of adaptive tracking control with Gaussian radial-basis-function (RBF) networks whose weights, centres and widths are adapted online. Part I is the projection-based adaptive RBF controller of version 1. An audit found its stability argument to be a correct but standard composition of known results, with two claims wrong as written (a non-convex "projection" for centre separation and an implicit-function-theorem argument that is not a proof); it contains no new theorem. We state it with explicit assumptions, a Hessian-free disturbance bound and an RMS bound, and list the errata to version 1. Part II develops transport adaptive control: the network is an empirical measure on a lifted parameter space, and the adaptive law is its transport. Theorems A–D and F are proved; A, B, C and F are also checked by tests. Theorem E is proved conditional on Theorem 1 of Morgan and Narendra (1977), in the form stated in that paper's abstract; its hypotheses were not checked against the full text. Proposition G is formal and checked directly. Theorem H shows that the averaged mean-field loop with saturated features, σ-modification and dither is mean-field Langevin dynamics and converges globally and exponentially to the regularised optimum; this is proved for the averaged system, the reduction to it is formal, and the real loop is a conjecture. At finite gain the averaged loop is in general not a gradient flow of any free energy in the $W_{G^{-1}}$ metric: Proposition H.7 (frozen-measure part proved) exhibits a semi-gradient term, whose performative defect is non-reciprocal in the checked instances ($n = 1, 2$), and an orbit-area circulation kernel ($n \geq 2$). With a second-order averaging term they predict a closed-loop effect for which a targeted literature search found no precedent: traversing the reference orbit backwards changes the learned kernels by $O(1/k_e)$, although the occupation measure is unchanged. At $k_e = 40$ the prediction matches to 3% in size and 11% as a vector (cosine 0.994). Proposition H.8, proved assuming the chain rule for the perturbed flow, gives input-to-state stability of the averaged system.

Contents

1 Introduction **4**
- 1.1 The idea . . . 4
- 1.2 Why the classical theory needs it . . . 4
- 1.3 What the closed loop actually is . . . 4
- 1.4 Conventions and verification . . . 5
- 1.5 Contributions and their status . . . 5
- 1.6 Where the new mathematics lies . . . 5
- 1.7 Organisation . . . 5

2 Relation to version 1 **7**
- 2.1 What version 1 claimed . . . 7
- 2.2 The audit's verdict . . . 8
- 2.3 Errata to version 1 . . . 10
- 2.4 What survives and what is withdrawn . . . 11
- 2.5 How Part II answers the structural defects . . . 11

I The projection-based controller **13**

3 Setting and standing assumptions **13**

4 The algorithm **14**

5 Lemmas and the boundedness theorem **14**
- 5.1 Lemmas . . . 14
- 5.2 Theorem 1: uniform ultimate boundedness . . . 16

6 The no-bias claim, stated precisely **17**

7 Prior art **17**

8 What the experiments show **18**
- 8.1 Robustness mechanism versus geometry . . . 18
- 8.2 Kernel-regression baselines . . . 19
- 8.3 The bound against data . . . 20
- 8.4 Summary . . . 22

II Transport adaptive control **25**

9 Setting **25**

10 Theorem A: identification **27**

11 Theorem B: the optimal-assignment Lyapunov function **29**

12 Theorem C: optimal-transport σ-modification **33**

13 Theorem D: the mean-field closed loop **38**

14 Theorem E: label-free local convergence under tangential excitation **40**

15 Theorem F: the transport cost of adaptation **44**

16 Proposition G: the slow geometry **45**

17 Theorem H: Langevin adaptive control and global convergence to the regularised optimum **50**

18 Finite gain: semi-gradient and non-reciprocal structure **56**
18.1 Frozen-measure structure of the exactly averaged loop 56
18.2 The oscillation correction and the orientation of the reference orbit 59
18.3 Input-to-state stability of $(\mathfrak{A})$ and the route to Conjecture H.6 61
18.4 Evidence from the real closed loop 63

III Evidence and outlook **64**

19 Numerical evidence **64**
19.1 Theorem B: the label-free Lyapunov function 64
19.2 Theorem C: OT σ-modification 64
19.3 Theorem D: convergence in m 65
19.4 Theorem E: label-free convergence under excitation 66
19.5 Theorem F: transport cost 67
19.6 Proposition G: slow geometry 68
19.7 Theorem H: Langevin adaptive control 69
19.8 Remark H.0 and Proposition H.7: the exactly averaged closed loop 70
19.9 Remark H.7a: the orientation of the reference orbit 71

20 Research program **72**

A Verification map **74**

1 Introduction

1.1 The idea

An adaptive approximator with nonlinearly parametrised features can be read as an *empirical measure* $\mu = \frac{1}{m}\sum_i \delta_{\vartheta_i}$ on a lifted parameter space. Here the features are Gaussian RBF kernels with adaptive weights, centres and widths. The adaptive law transports this measure, and the control output $\hat{f} = \int \Phi \, \mathrm{d}\mu$ is linear in μ. Stability, robustness modifications, excitation, the many-kernel limit and the slow evolution of kernel geometry all acquire formulations on Wasserstein space. So does the global optimality of kernel placement. These formulations do not depend on how the kernels are labelled.

1.2 Why the classical theory needs it

This work identifies two structural defects of the classical labelled analysis, that of version 1 of this manuscript and of the literature it builds on.

- The classical Lyapunov analyses depend on labels. A network that is a relabelling of the ideal has zero function error but a large parameter error.
- There is no object that describes where the kernels go, or whether their placement is good.

A third apparent defect was that constants grow with the number of kernels. It turned out to be an artefact of the normalisation. The mean-field normalisation removes it for labelled and label-free analyses alike (Remark A.1).

1.3 What the closed loop actually is

- At infinite feedback gain, the averaged closed loop is a Wasserstein gradient flow. With dither it is a mean-field Langevin dynamics (Theorem H).
- At finite gain it is in general *not* a gradient flow of any free energy in the $W_{G^{-1}}$ metric. This already holds for $n = 1$, so it is not a two-dimensional effect (Proposition H.7). The first correction has two parts at a frozen measure, and a third along the real loop.
  - The first part is a *semi-gradient* term. The adaptive law differentiates the loss on the closed-loop state distribution while ignoring that this distribution depends on the parameters, as in performative prediction. Like semi-gradient TD, the resulting update is not the gradient of any objective. Its performative defect is present for every n, and it is non-reciprocal in the checked instances ($n = 1$ and $n = 2$).
  - The second part is an antisymmetric interaction between kernels ($n \geq 2$). Its kernel is an integral of $\sum_k \det[\nabla\tilde{\Phi}_k(\vartheta), \nabla\tilde{\Phi}_k(\vartheta')]$ over the region that the reference orbit encloses (Stokes).
  - Along the real loop, a third part comes from the parameters' oscillation within each period. It is a second-order averaging term (Remark H.7a).
- Together these predict a new closed-loop effect: traversing the reference orbit backwards changes the learned kernels by $O(1/k_e)$, although the occupation measure is unchanged. At $k_e = 40$ the prediction matches to 3% in size and 11% as a vector (cosine 0.994). The vector error halves with each doubling of k_e (`experiments/exp_orbit_orientation.py`).
- The gradient structure survives as an input-to-state Lyapunov function (Proposition H.8). This reduces the global conjecture for the real loop to an averaging estimate.

1.4 Conventions and verification

Status labels. Every result of Part II is labelled *proved*, *formal* or *conjectural*, and the label is given in a status line after the result.

- "Proved" means proved here. "Formal" means a derivation without error control.
- "Checked" means a test evaluates the statement, or every inequality in it, along simulated trajectories or on sampled states.
- The novelty column of Table 1 follows the verdicts in `LITERATURE_TRANSPORT.md`.

In Part I, every step of the stability argument is either proved or verified numerically by `tests/test_theorem_to_code.py`.

Verification and review.

- Every inequality the tests can evaluate is checked numerically along simulated trajectories, and every number in this document is produced by a script in the accompanying repository (the experiments are in `experiments/`); no table is edited by hand.
- The theory went through three rounds of automated multi-agent adversarial review. Every confirmed issue and its fix is logged in `review/REVIEW_LOG.md`, and the raw reports are kept beside it.
- The claim-by-claim novelty audit is in `LITERATURE_TRANSPORT.md`.

Appendix A maps each checked statement to the test or experiment that checks it.

1.5 Contributions and their status

Table 1 lists the results of Part II with their status, in the conventions of Section 1.4, and with the novelty verdict of the literature audit.

1.6 Where the new mathematics lies

The new mathematics this viewpoint requires lies in the open problems of Section 20.

1. Infinite-horizon averaging for McKean–Vlasov closed loops. Propositions H.7 and H.8 reduce it to controlling a near-identity averaging transformation along the moving measure.
2. Uniform-in-time propagation of chaos with a plant in the loop.
3. The theory of the non-reciprocal, semi-gradient mean-field dynamics that the closed loop actually produces.
4. Global convergence without dither.

1.7 Organisation

Section 2 relates this document to version 1: what version 1 claimed, the audit's verdict, the errata, what survives and what is withdrawn, and how Part II answers the structural defects of Section 1.2. Part I treats the projection-based controller: the setting and standing assumptions (Section 3), the algorithm (Section 4), the lemmas and the uniform-ultimate-boundedness theorem (Section 5), the "no bias" claim stated precisely (Section 6), prior art and the honest novelty statement (Section 7), and the experiments (Section 8). Part II develops transport adaptive control in Sections 9–18: the setting (Section 9), Theorems A–F (Sections 10–15), Proposition G (Section 16), Theorem H (Section 17) and the finite-gain structure of Propositions H.7 and H.8 (Section 18). Part III collects the numerical evidence (Section 19) and the research program (Section 20). Appendix A is the verification map.

Table 1: Contributions of Part II and their honest status. Status conventions: Section 1.4.

result	statement	status	novelty (audit)
A	The adaptive law is the particle form of the Wasserstein gradient flow of a μ-linear potential. Projected rates coincide only for a per-kernel projection ($m \geq 2$).	proved and checked	interpretive; ingredients known [8], [13], [33], [45]
B	$V_{\mathrm{OT}} = \frac{1}{2}\|e\|^2 + \frac{1}{2}W^2$ satisfies a Dini inequality with a two-regime matched remainder. It gives conditional RMS bounds along (L), and along (L_σ) with a leak term.	proved and checked, including a moving ideal, $\varepsilon \neq 0$, switches, an active projection, and the failure of the plain bound along (L_σ)	packaging new; ingredients known [55], [1], [44], [10]
C	OT σ-modification: exact dissipation for selection solutions; a gap-protected strict inequality with $\sqrt{C} \leq b + 2\eta$; a local exponential estimate, m-uniform for a single-atom prior and $O(m^{-1/2})$ otherwise; reshuffling by the labelled leak	proved and checked; the C(iv) instance satisfies every hypothesis	novel as a control construction; C(i) and the obstruction are known [53], [5]
D	Local well-posedness of the mean-field loop; a Dobrushin estimate; a tube bound and global existence under a self-consistent growth condition; the mean-field limit on the existence interval	proved; consistency in m checked numerically (the constant is not checked)	technique anticipated [17, 47, 48]
E	Label-free local exponential convergence under tangential PE, in a uniform tube, at fixed m; exponential merging of solutions	proved, conditional on [39, Thm. 1] as stated in its abstract (its hypotheses were not checked against the full text); linearisation checked; convergence observed	partially anticipated [10], [22]
F	Along (L), the Wasserstein length of learning is $\leq \lambda_G^{1/2} L_1 \int \|e\|$; leak versions	proved and checked, including the leak clause, a tight leak instance with $e \equiv 0$ and a tight velocity instance	elementary
G	Averaged joint descent on the occupation loss. The slow geometry follows the gradient flow of the profile loss in the metric induced on the slow manifold, which is plain variable projection only when that manifold is flat ($S \ll 1$).	formal; checked directly, including a counterexample to plain VarPro at $S = O(1)$	a fixed-m, closed-loop analogue of the two-timescale/VarPro training of [38], [4]
H	The averaged mean-field loop with saturated features, σ-modification and dither is mean-field Langevin dynamics. It converges globally and exponentially to the regularised optimum, with the log-Sobolev constant given in control quantities.	proved for ($\mathfrak{A}$); the reduction to ($\mathfrak{A}$) is formal (Remark H.0); real loop: Conjecture H.6	a bridge to [9], [41]; new in adaptive control
H.7	At a frozen measure, the exactly averaged loop is ($\mathfrak{A}$) plus an explicit $O(1/k_e)$ correction: a semi-gradient term, whose performative defect is present for every n and non-reciprocal in the checked instances ($n = 1, 2$), and an orbit-area circulation kernel ($n \geq 2$). Along the real loop, a second-order averaging term is added (Remark H.7a).	Frozen-measure part proved. Stokes identity and parity hold to machine precision. The expansion is checked through its $O(1/k_e)$ convergence. Non-reciprocity of the exact second variation is checked to 3% (asserted; 1–1.5% measured). The full first-order theory predicts the orientation dependence of the learned kernels in the real closed loop to 3% in size and 11% as a vector (cosine 0.994, $k_e = 40$).	no precedent found (targeted search in the third review round); ingredients known (higher-order averaging; performative prediction [46]; semi-gradient TD [58], [56])
H.8	($\mathfrak{A}$) is input-to-state stable under drift perturbations, with gain $1/(\tau\rho_\tau)$	proved, assuming the chain rule for the perturbed flow	elementary given [9]

2 Relation to version 1

Version 1 of this document, *Mathematical Foundations of Self-Adaptive Kernels Achieving Lyapunov Convergence* (14 pages), presented the controller of Part I. An audit (`ASSESSMENT.md`) re-examined its theory, its code and its experiments; every numerical statement of the audit was regenerated by running the code, and the scripts and raw files are named next to each claim. A revised theory (`THEORY_REVISED.md`, Part I of this document) replaced its stability argument. This section records what version 1 claimed, the audit's verdict, the errata, what survives and what is withdrawn, and how Part II answers the structural defects of Section 1.2.

2.1 What version 1 claimed

Version 1 was subtitled *Uniform Ultimate Boundedness via Projection-Based Adaptive RBF Control Without Persistent Bias of Leaky Modification.* It treated the plant $\dot{x} = f(x,t) + u$ on a compact domain with the certainty-equivalence control $u = \dot{x}_m - k_e e - \hat{f}(x,t)$ and the network $\hat{f} = W^\top \phi(x, c, \sigma)$, adapted weights, centres and widths by projected gradient laws, and used the Lyapunov candidate $V = \frac{1}{2} e^\top e + \frac{1}{2}\operatorname{tr}(\tilde{W}^\top \Gamma_W^{-1} \tilde{W}) + \frac{1}{2}\tilde{c}^\top \Gamma_c^{-1} \tilde{c} + \frac{1}{2}\tilde{\sigma}^\top \Gamma_\sigma^{-1} \tilde{\sigma}$. Its claims were the following.

- *Central contribution* (abstract): "a rigorous proof that a fully shape-shifting kernel architecture — with online adaptation of weights, centers, and bandwidths — achieves Uniform Ultimate Boundedness (UUB) of the tracking error without the persistent steady-state bias introduced by classical leaky modification schemes (σ-modification, e-modification)".
- *Key mechanism* (abstract, §1.3): "the strict decoupling of offline Tikhonov regularization, which guarantees the well-posedness of the ideal parameter target, from the online smooth projection operator, which provides hard parameter boundedness", with an implicit-function-theorem (IFT) argument giving finite velocity bounds V_W, V_c, V_σ for the ideal parameters.
- *A "critical and non-trivial" result* (abstract, §7.2): the projection's hard cap on parameter energy "converts the standard boundary-only Lyapunov descent argument into a global interior argument", proving convergence to the infimal invariant set without LaSalle's invariance principle or Barbalat's lemma, with the ultimate bound

$$\limsup_{t\to\infty} \|e(t)\| \le \sqrt{R_{\mathrm{UUB}}^2 + 2V_{\max,\Theta}}.$$

- Centre separation $\|c_i - c_j\| \ge \delta > 0$, "maintained via the projection set construction" (§1.2); and a corollary that a standard persistent-excitation condition on the regressor is insufficient to guarantee parameter convergence, so that alternative mechanisms, such as integral concurrent learning (ICL), are required (§8).
- §9 summarised five architectural choices and the failure mode each was said to close: $\sigma_{\min} > 0$ (Hessian blow-up near $\sigma \to 0$); offline Tikhonov regularisation (circular self-consistency of the projection geometry); a smooth convex projection operator (a cancellation gap when the projection is active); the projection's hard energy cap (the LaSalle and Barbalat requirements); and stability from the Lyapunov function and the projection alone (the circularity of ICL, which needs bounded states).
- §10 stated that "the theoretical core of this manuscript is mathematically complete and internally consistent", and that the remaining items were "completeness additions and empirical validation requirements …, not theoretical repairs": a constructive feasibility theorem; empirical validation (Lorenz-driven tracking against σ- and e-modification "to prove bias elimination", a Monte Carlo sweep "verifying R_{UUB} bound is not violated", an

Table 2: Novelty of the ingredients of version 1 (audit).

ingredient	prior source	status of the claim of version 1
Adaptive RBF control with projection-type robustification	Sanner & Slotine 1992 [52]; Ioannou & Sun 1996 [30]	known
Adapting centres/widths by first-order Taylor expansion with bounded remainder	Lewis, Yesildirek & Liu 1996 [36]; Polycarpou 1996 [49]; Fabri & Kadirkamanathan 1996 [16]; Ge & Wang 2002 [21]	known; version 1 cites none of these
Smooth projection with ε boundary layer and inequality (P)	Pomet & Praly 1992 [50]; Lavretsky & Gibson 2011 [34]; Lavretsky & Wise 2013 [35]	known; the showcase README called it "the core theoretical contribution"
Projection introduces no bias in $\dot{V}$ when $\theta^* \in \Omega$	Ioannou & Sun 1996 [30, §4.4]	known
Level-set ("global interior") UUB argument	Khalil [31, Thm. 4.18]; Lewis et al. 1999 [37]	known; called "critical and non-trivial"
Simultaneous adaptation of W, c, σ for Gaussian RBFs with all three projected	—	the only combinational novelty; minor

ablation of fixed against adaptive centres); and a novelty framing in which the architecture "eliminates this tradeoff entirely", the projection introduces "zero bias into the Lyapunov derivative", and the result is "not a marginal improvement but a structural result enabled by the strict decoupling of offline regularization from online projection".

2.2 The audit's verdict

The theory is a correct but standard composition of known results (Taylor-linearised inner-layer adaptation, boundary-layer projection, level-set UUB), presented with language that overstates its novelty and its conclusion. Two of its claims are wrong as written (non-convex "projection" for centre separation; an IFT argument that is not a proof) and one was not implemented (the Γ-weighted projection). The experiments contained a hand-chosen "theoretical" threshold, a confounded headline comparison against baselines that failed representationally, GP baselines handicapped by stale predictions and a mis-placed inducing grid, invalid significance tests on autocorrelated samples, and the public figure omitted the one benchmark ($n = 3$) where the method is weak and the one baseline (NTK) that beats it. After revision, the implemented algorithm is exactly the proved one and is machine-checked; the honest experiments show a real, consistent advantage of projection over σ- and e-modification on identical geometry (5/5 seeds in every cell) and a modest but consistent gain from adaptive geometry, while the a-priori bound is shown to be 915× to 10 954× loose.

Novelty. Table 2 compares the ingredients of version 1 with prior work. Net: no new theorem. What can honestly be claimed after revision (Section 7) is a machine-checked theorem-to-code map, an explicit Hessian-free disturbance bound, the RMS form of the bound, and a controlled mechanism-versus-geometry experiment.

Integrity findings. Table 3 lists the integrity findings of the audit, each verified by running code, with the evidence and the fix. File names refer to the accompanying repository.

Recommendation. The audit recommended that version 1 not be submitted in its form, but re-framed as an empirical-plus-verified-implementation paper: the factorial result is a clean,

Table 3: Integrity findings of the audit of version 1 and its code (each verified by running code).

#	finding	evidence	fix
I1	Implemented centre update $\neq$ proved update: a pairwise repulsion term broke the projection inequality the proof needs	`tests/test_theorem_to_code.py` (oracle plant): inequality violated at 6054 of 20000 samples with the legacy term, 0 of 20000 without; the term also failed its own purpose (min separation 0.07 vs $\delta = 0.5$)	removed from the algorithm (`enforce_separation=False` default); tracking unchanged on the headline run (mean steady-state error $0.0767 \to 0.0774$, max $0.46 \to 0.40$)
I2	Projection applied the isotropic formula to any Γ; the proof needs the Γ-weighted operator	`tests/test_controller_paths.py`	Γ-weighted projection implemented; fast/dense paths agree to 10^{-12}
I3	"R_{UUB} threshold" in the Monte Carlo figure was the constant 0.5	`tier1_monte_carlo.py` line 69: "A chosen arbitrary bound"	`exp_monte_carlo_honest.py`: only exactly checkable statements (200/200 hard-bound and ISS checks)
I4	"Theoretical R_{UUB}" in the k_e sweep used invented constants and an underived formula	`tier2_ruub_sweep.py`, former `estimate_D_max`	`exp_oracle_ruub.py`: all constants computable; a-priori bound 915×–10 954× loose; RMS bound with measured disturbance holds 48/48
I5	Headline figure: fixed-grid baselines with a grid not covering the reference; their error (≈ 15) exceeded open loop, so the figure showed representational failure, not "bias"	`results/tier1_lorenz_tracking.png` vs ablation figure (projection W-only fails identically)	`exp_bias_factorial.py`: 3 mechanisms $\times$ 3 geometries, covering grid, 5 seeds
I6	Wilcoxon tests on ∼12 000 time samples with lag-1 autocorrelation 0.9999; the README's "$p > 0.3$ vs Exact GP" is contradicted by its own data ($p = 1.6 \times 10^{-11}$)	recomputed from `results/*.npy`	seed-level statistics only
I7	$n = 3$ benchmark and NTK baseline removed from the public figure (commits "exclude n=3 and NTK")	`results/n3/*.npy`, `results/*/ntk_*.npy`; `plot_comparison_hd_complete.py`	complete table regenerated: proposed ranks 3/5, 2/5, 3/5, 2/5
I8	GP baselines held a stale prediction for 50–100 steps; sparse GP inducing points on a diagonal line over $x \in [-3, 3]$ while the reference spans ± 20	`sim1`–`sim3`	`kernel_baselines_fair.py`: per-step prediction, 9×3 covering grid, seeds
I9	"$O(mn)$ per step" claim false with the $O(m^2 n)$ repulsion	`projection.py`	true after I1
I10	"Concurrent Learning" hook used the current residual only	former `controller.py`	removed
I11	Lyapunov monitor requires θ^*; never used by any experiment, so "Lyapunov descent verification" was never run	`integrator.py`	oracle module and tests perform it on a representable plant

reproducible finding, and the theory should be the revised one (Sections 3–6) with the prior work cited.

2.3 Errata to version 1

Table 4 lists the errata to version 1. Section numbers in the second column refer to version 1; "README" refers to the public showcase README that accompanied it.

Table 4: Errata to version 1.

#	section	statement in version 1	status
E1	1.2, 9	Centre separation $\|c_i - c_j\| \geq \delta$ "maintained via the projection set construction" with "convex projection boundaries"	**Wrong.** The separation set is non-convex; no projection onto it has property (P). The implemented repulsion term violated (P) in 30% of samples and did not keep the separation (min distance 0.07 vs $\delta = 0.5$). Removed from the algorithm; not needed by the proof.
E2	1.3	Offline Tikhonov regularisation of the ideal target: "This strict separation eliminates the circularity"	**Irrelevant to the proof.** Optimality of θ^* is never used (Section 3, remarks on the assumptions).
E3	1.3	IFT gives finite velocity bounds V_W, V_c, V_σ	**Not a proof.** The Hessian of a Tikhonov-regularised non-convex loss at a minimiser is PSD, not necessarily PD; the IFT hypothesis is asserted, not established. Replaced by Assumption A4.
E4	2.1	$D_{\max}$ from the Taylor remainder, "a finite geometric scalar"	**Valid but not explicit.** Replaced by the explicit, Hessian-free Lemma 2. The code's former `estimate_D_max` $= \varepsilon(1 + 2W_{\max}c_{\max})$ had no derivation and is deleted.
E5	4	$\mathrm{tr}(\tilde{W}^\top \Gamma_W^{-1} \tilde{W})$ with Γ_W "positive-definite learning rate matrices"	**Type error** for $\Gamma_W \in \mathbb{R}^{mn \times mn}$; the correct form is $\mathrm{vec}(\tilde{W})^\top \Gamma_W^{-1} \mathrm{vec}(\tilde{W})$. Harmless for scalar Γ.
E6	5	Projection inequality "holds as an absolute mathematical certainty — the cancellation gap does not exist"	**True only under A3** (θ^* in the inner set) and only for the Γ-weighted operator; the previous code applied the isotropic formula to any Γ. Fixed and tested.
E7	7.2	"A critical and non-trivial consequence of the projection operator"; its hard energy cap "converts what would be a boundary-only argument … into a global interior argument"	**Standard** level-set / comparison argument (Khalil [31, Thm. 4.18]). Correct, not new; also "strict monotone descent through every intermediate level set until the infimal set is penetrated" overstates: entry into every level above $c_{\inf}$ is finite-time, entry into $c_{\inf}$ itself only asymptotic.
E8	abstract, 7.3, 10.3	Projection gives UUB "without the persistent steady-state bias" (abstract); "no modification bias is introduced" (7.3); "zero bias into the Lyapunov derivative", "not a marginal improvement but a structural result" (10.3)	**Overclaimed.** The ultimate bound $\sqrt{R_{\mathrm{UUB}}^2 + 2V_{\max,\Theta}}$ contains the parameter-energy term; see Section 6. The tighter, bias-free statement is the RMS bound (iii).
E9	8	"a standard Persistent Excitation (PE) condition on the regressor ϕ is insufficient to guarantee $\tilde{\Theta}(t) \to 0$"; convergence "requires alternative mechanisms", such as ICL	Parameter convergence is not treated; the remark is speculation and the code's former "Concurrent Learning" hook used only the current residual (not recorded data) and is removed.
E10	README	"100% of the independent simulation trials converged strictly below the calculated R_{UUB} threshold"	**False.** The threshold 0.5 was a hand-chosen constant (`tier1_monte_carlo.py`: "A chosen arbitrary bound"). Replaced by `exp_monte_carlo_honest.py` / `exp_oracle_ruub.py`.
E11	README	"$O(mn)$ time"	**Was false** with the $O(m^2 n)$ pairwise repulsion; true now that it is off by default.

2.4 What survives and what is withdrawn

What survives.

- *The corrected theorem.* Theorem 1 of Section 5 holds under the explicit Assumptions A1–A5, for an algorithm identical to the implemented one. Its pointwise ultimate bound (ii) is the conclusion of version 1; it is the textbook level-set argument, and its radius contains the parameter-energy term $V_{\max,\Theta}$. Its RMS bound (iii) does not contain that term and is the form of the result that can be confronted with data; its reference bound (iv) holds whatever the adaptation does. The ingredients are the projection property (P) (Lemma 1), the explicit Hessian-free disturbance bound (Lemma 2) and the Lyapunov derivative (Lemma 3), verified numerically along simulated trajectories. Nothing in the stability theorem is new (Section 7). The "no bias" claim survives only in the precise form of Section 6: projection removes the *Lyapunov-derivative* bias of leakage, not the ultimate *tracking* bound's dependence on the parameter set.
- *The projection-versus-leakage finding* (Section 8). On identical geometry, projection beats both σ- and e-modification in every cell of a mechanism-by-geometry factorial, 5/5 seeds, with paired confidence intervals excluding zero. Within projection, adaptive centres improve on the fixed grid by 0.012 (5/5) and adaptive widths by a further 0.035 (5/5). So the method's real advantage is the robustness mechanism, with geometry adaptation a smaller second-order gain, close to the reverse of the story told by the original figure.
- *The comparison with kernel regression, with its limits* (Section 8). Against kernel-regression baselines the method is never the most accurate. Its edge is cost: it is cheaper per step than every GP baseline at every n and than NTK except at $n = 3$, and in the fair $n = 1$ rerun, with per-step prediction for every method, it has the cheapest step. Every kernel-regression baseline (GP and NTK, including the fair $n = 1$ rerun) is trained on noise-free samples of the true $f(x, t)$, which is equivalent to measuring the state derivative exactly, while the adaptive law uses only the tracking error.

What is withdrawn.

- Earlier versions of the public README stated that the method's steady-state error was "statistically indistinguishable (Wilcoxon signed-rank $p > 0.3$) from both the Exact GP and the Sparse GP", that "100% of the independent simulation trials converged strictly below the calculated R_{UUB} threshold", and announced "Curse of Dimensionality Solved". Those statements are withdrawn: the significance test was invalid, the threshold was an arbitrary constant, and the omitted $n = 3$ and NTK results contradict them.
- The "zenith" language is dropped: "absolute mathematical certainty", "curse of dimensionality solved", "structural result". None of it survives the data. Claims of a "structural" advantage over leaky modifications are withdrawn.
- The claim that centre separation is "maintained via the projection set construction" is withdrawn (erratum E1); no centre-separation constraint is needed by the proof.

2.5 How Part II answers the structural defects

Section 1.2 named two structural defects of the classical labelled analysis, that of version 1 and of the literature it builds on: its Lyapunov analyses depend on labels, and it has no object that describes where the kernels go or whether their placement is good. Part II addresses both by working with the measure $\mu = \frac{1}{m}\sum_i \delta_{\vartheta_i}$, which does not see the labels, and with its transport.

- *Labels.* The Lyapunov function of Theorem B, $V_{\mathrm{OT}} = \frac{1}{2}\|e\|^2 + \frac{1}{2}W(\mu_t, \mu_t^*)^2$, measures the parameter error by the optimal-assignment (Wasserstein) distance W, a minimum over

all assignments of kernels to ideal atoms (Section 9). The labelled σ-modification is not a functional of μ; Theorem C replaces it by an optimal-transport σ-modification, and Proposition C.5 shows that the labelled leak reshuffles kernels. Theorem E gives label-free local exponential convergence under tangential persistent excitation (proved conditional on a cited theorem).

- *Where the kernels go.* The measure μ_t is that object. By Theorem A the adaptive law is the particle form of the Wasserstein gradient flow of a μ-linear potential; Theorem D treats the mean-field (many-kernel) loop; Theorem F bounds the Wasserstein length of learning by the integrated tracking error; Proposition G describes the slow geometry (formal). Whether the placement is good is the subject of Theorem H: with saturated features, σ-modification and dither, the averaged mean-field loop converges globally and exponentially to the regularised optimum. This is proved for $(\mathfrak{A})$; the reduction to $(\mathfrak{A})$ is formal, and the real loop is Conjecture H.6, which Propositions H.7 and H.8 reduce to an averaging estimate.
- *Constants growing with m.* The third apparent defect is an artefact of the normalisation; the mean-field normalisation removes it for labelled and label-free analyses alike (Remark A.1).

The status of each of these results is that of Table 1.

Part I
The projection-based controller

3 Setting and standing assumptions

This part replaces the stability argument of version 1 with a statement whose assumptions are explicit, whose algorithm is exactly the one in `src/core/` of the accompanying repository, and whose every step is either proved here or verified numerically by `tests/test_theorem_to_code.py`. The errata to version 1 are listed in Section 2; Section 7 positions the result against prior work.

Notation. $\|\cdot\|$ is the Euclidean norm for vectors and the Frobenius norm for matrices; $\mathrm{vec}(\cdot)$ is the row-major vectorisation used throughout the code; there are m kernels and n states.

Plant, reference and error.

$$\dot{x} = f(x,t) + u, \qquad e = x - x_m(t), \qquad u = \dot{x}_m - k_e e - \hat{f}(x), \tag{3.1}$$

$$\hat{f}(x) = W^\top \varphi(x, c, \sigma), \qquad \varphi_i(x) = \exp\Big(-\frac{\|x - c_i\|^2}{2\sigma_i^2}\Big). \tag{3.2}$$

Assumption A1 (operation on a compact set)**.** There is a compact set $X \subset \mathbb{R}^n$ with $x(t) \in X$ for all $t \geq 0$, and $F_{\max} := \sup_{x \in X,\, t \geq 0} \|f(x,t)\| < \infty$. *(Version 1 assumes "a compact spatial domain X" in the same way. For globally bounded f the assumption is discharged by Theorem 1(iv); for the Lorenz plant it is a genuine assumption.)*

Assumption A2 (reference)**.** x_m is C^1 with x_m, $\dot{x}_m$ bounded, and $\dot{x}_m$ is available to the controller.

Assumption A3 (admissible ideal parameters)**.** There is a curve $\theta^*(t) = (W^*(t), c^*(t), \sigma^*(t))$ lying for all t in the *inner* projection sets

$$\|W^*\| \leq W_{\max} - \varepsilon, \qquad \|c^*\| \leq c_{\max} - \varepsilon, \qquad \sigma_{\min} + \varepsilon \leq \sigma_i^* \leq \sigma_{\max} - \varepsilon.$$

Assumption A4 (ideal-parameter regularity)**.** $\theta^*(\cdot)$ is C^1 with $\|\dot{W}^*\| \leq V_W$, $\|\dot{c}^*\| \leq V_c$, $\|\dot{\sigma}^*\| \leq V_\sigma$. (For a time-invariant plant one may take θ^* constant, $V_W = V_c = V_\sigma = 0$.)

Assumption A5 (gains and geometry)**.** $k_e > 0$; Γ_W, Γ_c symmetric positive definite on $\mathbb{R}^{mn}$ (a zero Γ freezes that group, see Remark 3); Γ_σ diagonal positive; $0 < \varepsilon < \min\big(W_{\max},\, c_{\max},\, (\sigma_{\max} - \sigma_{\min})/2\big)$; $\sigma_{\min} > 0$.

Remarks on the assumptions.

- **The proof never uses optimality of θ^*.** θ^* is an analytic device; any admissible C^1 curve works, and the approximation quality of the network enters the result only through the *actual* size of the disturbance $d(t)$ defined below, not through the bound. The Tikhonov-regularised definition of θ^* in version 1 and its implicit-function-theorem (IFT) argument for A4 are therefore unnecessary — and, as written, not a proof (erratum E3, Section 2).
- **No centre-separation constraint is needed.** The set $\{c : \|c_i - c_j\| \geq \delta \ \forall\, i \neq j\}$ is not convex, so no projection onto it has property (P) below; the claim of version 1 that separation is "maintained via the projection set construction" is withdrawn (erratum E1, Section 2). The bounded-Jacobian property that the proof needs follows from $\sigma \geq \sigma_{\min}$ alone.

4 The algorithm

The algorithm is identical to the one implemented in `src/core/controller.py`:

$$\begin{aligned} \dot{W} &= \mathrm{Proj}_W\big(W,\ \Gamma_W\,\varphi e^\top\big), \\ \dot{c} &= \mathrm{Proj}_c\big(c,\ \Gamma_c\,\varphi_c'^\top W e\big), \\ \dot{\sigma} &= \mathrm{Proj}_\sigma\big(\sigma,\ \Gamma_\sigma\,\varphi_\sigma'^\top W e\big), \end{aligned} \tag{4.1}$$

with $\varphi_c' = \partial\varphi/\partial\,\mathrm{vec}(c) \in \mathbb{R}^{m\times mn}$ (row i has the single non-zero block $\varphi_i(x-c_i)^\top/\sigma_i^2$) and $\varphi_\sigma' = \mathrm{diag}\big(\varphi_i\|x-c_i\|^2/\sigma_i^3\big)$.

Norm-ball projection (W and c), boundary layer of width ε.

$$\mathrm{Proj}(\theta,\Gamma y) = \begin{cases} \Gamma y - h(\theta)\,\Gamma\theta\,\dfrac{\theta^\top\Gamma y}{\theta^\top\Gamma\theta} & \text{if } h(\theta)>0 \text{ and } \theta^\top\Gamma y>0, \\ \Gamma y & \text{otherwise,} \end{cases} \tag{4.2}$$

$$h(\theta) = \begin{cases} 0 & \text{for } \|\theta\| \le \theta_{\max}-\varepsilon, \\ \big((\|\theta\|-\theta_{\max}+\varepsilon)/\varepsilon\big)^2 & \text{in between,} \\ 1 & \text{for } \|\theta\| \ge \theta_{\max}. \end{cases} \tag{4.3}$$

Box projection (σ, component-wise). The update component is multiplied by $\mathrm{clip}\big((\sigma_i-\sigma_{\min})/\varepsilon, 0, 1\big)$ when it points below the lower face and by $\mathrm{clip}\big((\sigma_{\max}-\sigma_i)/\varepsilon, 0, 1\big)$ when it points above the upper face.

Discrete time. The simulations integrate with fixed-step RK4 and re-clip the parameters to their sets after every stage (`src/core/integrator.py`). The theorem is a continuous-time statement; the clip is a safety net that is inactive if the step is small enough (property (P) already removes the outward component on the boundary).

5 Lemmas and the boundedness theorem

5.1 Lemmas

Lemma 1 (projection property (P))**.** *Under A3 and A5, for every θ in its set and every y,*

$$\tilde\theta^\top\Gamma^{-1}\,\mathrm{Proj}(\theta,\Gamma y) \le \tilde\theta^\top y, \qquad \tilde\theta := \theta-\theta^*. \tag{P}$$

Proof. If the projection is inactive the two sides are equal. Otherwise

$$\tilde\theta^\top\Gamma^{-1}\big(\mathrm{Proj}\ -\Gamma y\big) = -h\,\frac{\theta^\top\Gamma y}{\theta^\top\Gamma\theta}\,\tilde\theta^\top\theta,$$

with $h>0$ and $\theta^\top\Gamma y>0$, and $\tilde\theta^\top\theta = \|\theta\|^2 - \theta^{*\top}\theta \ge \|\theta\|\big(\|\theta\|-\|\theta^*\|\big) \ge 0$ because $h>0$ forces $\|\theta\| \ge \theta_{\max}-\varepsilon \ge \|\theta^*\|$. For the box, component i, with s_i the clip factor applied to that component: $\tilde\sigma_i(s_i-1)y_i \le 0$ because $s_i \le 1$, and on the lower face $y_i<0$ while $\tilde\sigma_i = \sigma_i-\sigma_i^* \le \sigma_{\min}+\varepsilon-\sigma_i^* \le 0$ (symmetrically on the upper face). □

Status. Proved.

Checked by: `tests/test_projection.py` (randomised; isotropic and non-isotropic Γ; vector, matrix and box), together with the counterexample showing that (P) fails when θ^* is outside the inner set.

Lemma 2 (lumped disturbance)**.** *Define*

$$d := f - \hat{f} + \tilde{W}^\top \varphi + W^\top \varphi_c' \,\tilde{c} + W^\top \varphi_\sigma' \,\tilde{\sigma}. \tag{5.1}$$

Then identically

$$\dot{e} = -k_e e - \tilde{W}^\top \varphi - W^\top \varphi_c' \,\tilde{c} - W^\top \varphi_\sigma' \,\tilde{\sigma} + d, \tag{5.2}$$

and under A1, A3,

$$\|d\| \le D_{\max} := F_{\max} + 3W_{\max}\sqrt{m} + W_{\max}\big(2c_{\max}J_c + \sqrt{m}\,(\sigma_{\max} - \sigma_{\min})J_\sigma\big), \tag{5.3}$$
$$J_c = \mathrm{e}^{-1/2}/\sigma_{\min}, \qquad J_\sigma = 2\mathrm{e}^{-1}/\sigma_{\min}.$$

Proof. The identity is the definition of d substituted into $\dot{e} = -k_e e + f - \hat{f}$. For the bound: $\|f\| \le F_{\max}$; $\|\hat{f}\| \le \|W\|\|\varphi\| \le W_{\max}\sqrt{m}$ $(0 < \varphi_i \le 1)$; $\|\tilde{W}\| \le 2W_{\max}$; $\|\tilde{c}\| \le 2c_{\max}$; $\|\tilde{\sigma}\| \le \sqrt{m}\,(\sigma_{\max} - \sigma_{\min})$; the rows of φ_c' have disjoint supports, so

$$\|\varphi_c'\|_2 = \max_i \frac{\varphi_i \|x - c_i\|}{\sigma_i^2} \le \sup_{r \ge 0} \frac{r\,\mathrm{e}^{-r^2/(2\sigma^2)}}{\sigma^2} = \frac{\mathrm{e}^{-1/2}}{\sigma} \le J_c,$$

and $\|\varphi_\sigma'\|_2 = \max_i \varphi_i \|x - c_i\|^2/\sigma_i^3 \le 2\mathrm{e}^{-1}/\sigma \le J_\sigma$. □

Status. Proved.

Checked by: `tests/test_rbf_kernel.py` checks the two suprema by brute force.

Remark 1 (local form)**.** Writing $\varphi^* = \varphi(x; c^*, \sigma^*)$ and expanding to first order in $(\tilde{c}, \tilde{\sigma})$ gives

$$d = \varepsilon + W^{*\top} O_{\mathrm{ho}} + \tilde{W}^\top \varphi_c' \,\tilde{c} + \tilde{W}^\top \varphi_\sigma' \,\tilde{\sigma}, \qquad \varepsilon := f - W^{*\top}\varphi^*,$$

with O_{ho} the Taylor remainder — the decomposition of version 1. Near convergence ($\tilde{\theta} \to 0$) this shows $\|d\| \lesssim \|\varepsilon\| + O(\|\tilde{\theta}\|^2)$, which is why the controller performs far better than $D_{\max}$ suggests; but it does not yield a smaller global constant than Lemma 2, which needs no Hessian and no Taylor expansion.

Lemma 3 (Lyapunov derivative)**.** *Let*

$$V = \tfrac{1}{2} e^\top e + \tfrac{1}{2}\,\mathrm{vec}(\tilde{W})^\top \Gamma_W^{-1}\,\mathrm{vec}(\tilde{W}) + \tfrac{1}{2}\tilde{c}^\top \Gamma_c^{-1}\tilde{c} + \tfrac{1}{2}\tilde{\sigma}^\top \Gamma_\sigma^{-1}\tilde{\sigma}.$$

Along trajectories of the algorithm,

$$\begin{aligned}
\dot{V} &= -k_e\|e\|^2 + e^\top d + \mathrm{cross} - \Omega^*, \\
\mathrm{cross} &= \langle \tilde{W},\, \Gamma_W^{-1}\dot{W} - \varphi e^\top \rangle + \tilde{c}^\top\big(\Gamma_c^{-1}\dot{c} - \varphi_c'^\top W e\big) + \tilde{\sigma}^\top\big(\Gamma_\sigma^{-1}\dot{\sigma} - \varphi_\sigma'^\top W e\big) \le 0, \\
\Omega^* &= \langle \tilde{W},\, \Gamma_W^{-1}\dot{W}^* \rangle + \tilde{c}^\top \Gamma_c^{-1}\dot{c}^* + \tilde{\sigma}^\top \Gamma_\sigma^{-1}\dot{\sigma}^*, \qquad |\Omega^*| \le C_\Omega, \\
C_\Omega &:= 2W_{\max}\|\Gamma_W^{-1}\|_2 V_W + 2c_{\max}\|\Gamma_c^{-1}\|_2 V_c + \sqrt{m}\,(\sigma_{\max} - \sigma_{\min})\|\Gamma_\sigma^{-1}\|_2 V_\sigma.
\end{aligned} \tag{5.4}$$

Proof. Differentiate V, substitute the identity of Lemma 2 for $\dot{e}$, and use $e^\top \tilde{W}^\top \varphi = \langle \tilde{W},\, \varphi e^\top \rangle$, $e^\top W^\top \varphi_c' \,\tilde{c} = \tilde{c}^\top \varphi_c'^\top W e$, $e^\top W^\top \varphi_\sigma' \,\tilde{\sigma} = \tilde{\sigma}^\top \varphi_\sigma'^\top W e$. That cross ≤ 0 is Lemma 1 applied to the three groups; the bound on Ω^* is Cauchy–Schwarz with A3, A4. □

Status. Proved.

Checked by: `tests/test_theorem_to_code.py`: the identity (to machine precision) and cross ≤ 0 (at every sample) are verified along simulated trajectories, on a plant that is exactly representable with known constant θ^*.

5.2 Theorem 1: uniform ultimate boundedness

Theorem 1 (uniform ultimate boundedness)**.** *Assume A1–A5, let*

$$R_{\mathrm{UUB}} := \frac{D_{\max}}{2k_e} + \sqrt{\frac{D_{\max}^2}{4k_e^2} + \frac{C_\Omega}{k_e}},$$

and let $V_{\max,\Theta}$ be the supremum over the projection sets of the parameter part of V.

(i) ***Descent outside a ball.*** *$\dot V \le -k_e\|e\|^2 + D_{\max}\|e\| + C_\Omega$, hence $\dot V < 0$ whenever $\|e\| > R_{\mathrm{UUB}}$.*

(ii) ***Pointwise ultimate bound.*** *Every solution satisfies $V(t) \le \max\{V(0), c_{\inf}\}$ with $c_{\inf} := \frac{1}{2}R_{\mathrm{UUB}}^2 + V_{\max,\Theta}$, every level $c > c_{\inf}$ is reached in finite time, and*

$$\limsup_{t\to\infty}\|e(t)\| \le \sqrt{R_{\mathrm{UUB}}^2 + 2V_{\max,\Theta}}.$$

(iii) ***RMS bound (the quantitatively meaningful one).*** *For every $T_1 < T_2$,*

$$\mathrm{RMS}_{[T_1,T_2]}(\|e\|) \le \frac{D_{\max}}{2k_e} + \sqrt{\frac{D_{\max}^2}{4k_e^2} + \frac{C_\Omega}{k_e} + \frac{V(T_1)}{k_e(T_2-T_1)}},$$

so $\limsup_{T_2\to\infty}\mathrm{RMS}_{[T_1,T_2]}(\|e\|) \le R_{\mathrm{UUB}}$ — without the parameter-energy term of (ii). The same inequality holds with $D_{\max}$ and C_Ω replaced by the trajectory-wise suprema of $\|d(t)\|$ and $|\Omega^*(t)|$ on $[T_1,T_2]$.

(iv) ***Reference bound (adaptation-free).*** *Under A1 and the hard parameter bounds alone, $\dot e = -k_e e + (f - \hat f)$ with $\|f - \hat f\| \le F_{\max} + W_{\max}\sqrt{m} =: \tilde F$, so*

$$\|e(t)\| \le \mathrm{e}^{-k_e t}\|e(0)\| + \tilde F/k_e \qquad \text{for all } t,$$

whatever the adaptation does.

Proof. (i) Lemma 3 with Cauchy–Schwarz on $e^\top d$.

(ii) If $V > c_{\inf}$ then, since the parameter part of V is at most $V_{\max,\Theta}$, $\frac{1}{2}\|e\|^2 > \frac{1}{2}R_{\mathrm{UUB}}^2$, so $\|e\| > R_{\mathrm{UUB}}$ and $\dot V < 0$; more precisely, with $\Delta := V - c_{\inf} > 0$ one has $\|e\|^2 \ge R_{\mathrm{UUB}}^2 + 2\Delta$ and therefore

$$\dot V \le -g\Big(\sqrt{R_{\mathrm{UUB}}^2 + 2\Delta}\Big) =: -\psi(\Delta), \qquad g(r) = k_e r^2 - D_{\max} r - C_\Omega,$$

with g increasing for $r \ge R_{\mathrm{UUB}}$ and ψ of class $\mathcal{K}$. The comparison lemma gives $V(t) \le \max\{V(0), c_{\inf}\}$, finite-time entry into every sublevel set $\{V \le c\}$ with $c > c_{\inf}$, and $\limsup V \le c_{\inf}$, i.e. the stated bound on $\|e\|$ ($\|e\|^2 \le 2V$).

(iii) Integrate (i):

$$\begin{aligned} k_e\int_{T_1}^{T_2}\|e\|^2\,\mathrm{d}t &\le V(T_1) - V(T_2) + D_{\max}\int_{T_1}^{T_2}\|e\|\,\mathrm{d}t + C_\Omega(T_2-T_1) \\ &\le V(T_1) + D_{\max}\sqrt{T_2-T_1}\sqrt{\int_{T_1}^{T_2}\|e\|^2\,\mathrm{d}t} + C_\Omega(T_2-T_1) \end{aligned}$$

by Cauchy–Schwarz; with $S := \mathrm{RMS}_{[T_1,T_2]}(\|e\|)$ this reads

$$k_e S^2 - D_{\max} S - \big(C_\Omega + V(T_1)/(T_2-T_1)\big) \le 0,$$

and S is below the positive root.

(iv) Variation of constants. □

Status. Proved.

What (ii)–(iv) say together. Part (ii) is the conclusion of version 1; it is the textbook level-set argument (Khalil [31, Thm. 4.18]; Lewis et al. [37]) and its radius contains the parameter-energy reservoir $V_{\max,\Theta}$ — a bias-like term — so it does **not** establish that projection yields a tighter tracking bound than σ-modification. Part (iii) is tighter and is the statement that can be checked without unknown constants: replace $D_{\max}$ by the measured $\sup\|d\|$, see `experiments/exp_oracle_ruub.py`, where it is satisfied in 48/48 runs (Section 8). Part (iv) shows that mere boundedness of e is already a consequence of projection plus a stable error channel, independent of the adaptation law; the content of adaptive control is in making the *actual* $d(t)$ small, which no a-priori constant captures. In the oracle experiments the a-priori R_{UUB} exceeds the observed RMS error by factors of 915 to 10954.

Remark 2 (no LaSalle / Barbalat)**.** Version 1's claim that the argument needs neither LaSalle's invariance principle nor Barbalat's lemma is correct, and standard: UUB statements never need them. Nothing here claims asymptotic convergence of e or of $\tilde{\theta}$.

Remark 3 (frozen groups)**.** If $\Gamma_c = 0$ (fixed centres), drop the $\tilde{c}$ term from V; the identity of Lemma 2 still holds with $-W^\top \varphi_c' \tilde{c}$ absorbed into d, with the same bound. Likewise for σ.

Remark 4 (what is not proved)**.** Parameter convergence (it requires persistent excitation of a nonlinear-in-parameter regressor — not addressed); exact tracking; any statement about the size of the ideal approximation error ε.

6 The no-bias claim, stated precisely

With σ-modification,

$$\dot{V} \le -k_e\|e\|^2 - \tfrac{1}{2}\sigma_{\mathrm{mod}}\|\tilde{\theta}\|_\Gamma^2 + \tfrac{1}{2}\sigma_{\mathrm{mod}}\|\theta^*\|_\Gamma^2 + D\|e\| + C_\Omega:$$

the leak adds a positive constant and removes the parameter-energy reservoir, which gives exponential convergence of V to a ball whose radius grows with $\sigma_{\mathrm{mod}}\|\theta^*\|^2$. With projection,

$$\dot{V} \le -k_e\|e\|^2 + D\|e\| + C_\Omega:$$

no positive constant is added, but nothing dissipates parameter energy, which is why the pointwise bound (ii) of Theorem 1 keeps $V_{\max,\Theta}$. Projection therefore removes the *Lyapunov-derivative* bias of leakage (true, and known since Pomet–Praly [50] and Ioannou–Sun [30]), not the dependence of the ultimate *tracking* bound on the parameter set (false as stated in Sections 7.3 and 10.3 of version 1). Empirically the two mechanisms are compared on identical geometry in `experiments/exp_bias_factorial.py` (Section 8).

7 Prior art

- Adaptive Gaussian-network control with robust modification: Sanner and Slotine [52].
- Adaptation of inner-layer parameters via first-order Taylor expansion of the feature map with bounded remainder — the exact device of Sections 4 and 5: Lewis, Yesildirek and Liu [36]; Polycarpou [49]; Ge and Wang [21]; dynamic-structure RBF networks with tuned centres and widths: Fabri and Kadirkamanathan [16].
- Smooth projection with a boundary layer and the inequality (P), including the "no bias when $\theta^* \in \Omega$" observation: Pomet and Praly [50]; Ioannou and Sun [30, §4.4]; Lavretsky and Gibson [34]; Lavretsky and Wise [35, Ch. 11].
- σ- and e-modification and their bias: Ioannou and Kokotovic [29]; Narendra and Annaswamy [40].

- UUB via level sets: Khalil [31, Thm. 4.18]; Lewis, Jagannathan and Yesildirek [37].
- Concurrent learning (recorded-data stacks, finite excitation): Chowdhary and Johnson [11, 12].

Table 5 sets the ingredients of version 1 against these sources.

Table 5: Novelty of the ingredients of the stability argument.

Ingredient	Prior source	Status of the claim of version 1
Adaptive RBF control with projection-type robustification	Sanner and Slotine [52]; Ioannou and Sun [30]	known
Adapting centres/widths by first-order Taylor expansion with bounded remainder	Lewis, Yesildirek and Liu [36]; Polycarpou [49]; Fabri and Kadirkamanathan [16]; Ge and Wang [21]	known; version 1 cites none of these
Smooth projection with ε boundary layer and inequality (P)	Pomet and Praly [50]; Lavretsky and Gibson [34]; Lavretsky and Wise [35]	known; the public showcase README calls it "the core theoretical contribution"
Projection introduces no bias in $\dot{V}$ when $\theta^* \in \Omega$	Ioannou and Sun [30, §4.4]	known
Level-set ("global interior") UUB argument	Khalil [31, Thm. 4.18]; Lewis et al. [37]	known; called "critical and non-trivial"
Simultaneous adaptation of W, c, σ for Gaussian RBFs with all three projected	—	the only combinational novelty; minor

Net: no new theorem.

What is new here. Nothing in the stability theorem: Theorem 1(i)–(ii) is the composition of known pieces (Taylor-linearised inner-layer adaptation, boundary-layer projection and level-set UUB), applied to Gaussian RBFs with simultaneously adaptive weights, centres and widths. The contribution of the accompanying repository, after the revision of version 1, is (a) a theorem-to-code map in which the implemented and the proved algorithm coincide and are machine-checked (Lemmas 1–3 verified numerically along trajectories), (b) the explicit Hessian-free disturbance bound of Lemma 2 and the RMS bound (iii) of Theorem 1, which is the form of the result that can actually be confronted with data, and (c) a controlled empirical separation of "projection vs leakage" from "adaptive vs fixed geometry" (Section 6; `results/factorial/`). Claims of a "structural" advantage over leaky modifications are withdrawn.

8 What the experiments show

Every numerical statement in this section was regenerated from the scripts of the accompanying repository; the result file and the generating script are named with each table. Statistics are computed over seeds, not over autocorrelated time samples.

8.1 Robustness mechanism versus geometry

The factorial experiment `experiments/exp_bias_factorial.py` separates the two factors that the original headline figure (`results/tier1_lorenz_tracking.png`) confounded: there, the fixed-grid baselines used a grid not covering the reference, and their error (≈ 15) exceeded open loop, so the figure showed representational failure, not "bias". Here three robustness mechanisms (projection, σ-modification, e-modification, the leaky ones implemented on the same architecture

in `src/baselines/leaky_geometry.py`) are crossed with three geometries (fixed grid with only W adapted; adaptive W, c; adaptive W, c, σ) on Lorenz tracking with a covering grid, $m = 27$, $k_e = 10$ and 5 seeds (Tables 6 and 7, Figure 1).

Table 6: Robustness mechanism × geometry (Lorenz tracking, $m = 27$, $k_e = 10$). Seeds: 5; metric = steady-state $\|e\|$ over the last 40% of a 30 s run; mean ± 95% CI over seeds; max SS = maximum over the window, mean over seeds; "div" = diverged runs. Source: `results/factorial/table.md`.

geometry	mechanism	mean SS $\|e\|$	median SS	max SS	max $\|W\|_F$	min sep	div
fixed (W)	projection	0.1584 ± 0.0072	0.1288	0.559	47.5	1.365	0
fixed (W)	σ-mod	0.7327 ± 0.0223	0.5647	2.720	38.5	1.469	0
fixed (W)	e-mod	0.5619 ± 0.0306	0.4264	2.192	26.7	1.469	0
adaptive (W, c)	projection	0.1468 ± 0.0038	0.1020	0.751	40.9	0.039	0
adaptive (W, c)	σ-mod	0.4004 ± 0.0050	0.3468	1.863	32.3	0.078	0
adaptive (W, c)	e-mod	0.2327 ± 0.0024	0.1993	0.718	28.1	0.099	0
adaptive (W, c, σ)	projection	0.1119 ± 0.0043	0.0852	0.512	38.5	0.172	0
adaptive (W, c, σ)	σ-mod	0.3862 ± 0.0057	0.3422	1.655	30.9	0.215	0
adaptive (W, c, σ)	e-mod	0.2385 ± 0.0029	0.2103	0.820	27.7	0.239	0

Table 7: Paired (same seed) differences of mean steady-state $\|e\|$ for the factorial experiment. Top: projection minus baseline. Bottom: geometry effect within the projection mechanism. Source: `results/factorial/table.md`.

geometry	baseline	mean diff	95% CI	projection better in
fixed (W)	σ-mod	−0.5743	±0.0232	5/5 seeds
fixed (W)	e-mod	−0.4035	±0.0310	5/5 seeds
adaptive (W, c)	σ-mod	−0.2536	±0.0021	5/5 seeds
adaptive (W, c)	e-mod	−0.0858	±0.0025	5/5 seeds
adaptive (W, c, σ)	σ-mod	−0.2743	±0.0043	5/5 seeds
adaptive (W, c, σ)	e-mod	−0.1266	±0.0031	5/5 seeds
comparison		mean diff	95% CI	adaptive better in
adaptive (W, c) − fixed (W)		−0.0116	±0.0056	5/5
adaptive (W, c, σ) − adaptive (W, c)		−0.0349	±0.0026	5/5

Projection beats both leaky baselines in every cell, in 5/5 seeds, with paired confidence intervals excluding zero. Within projection, adaptive centres improve on the fixed grid by 0.0116 ± 0.0056 (5/5 seeds) and adaptive widths by a further 0.0349 ± 0.0026 (5/5 seeds). So the method's real advantage is the robustness mechanism, with geometry adaptation a smaller second-order gain — close to the reverse of the story told by the original headline figure.

8.2 Kernel-regression baselines

Table 8 is the comparison against kernel-regression baselines from saved single runs, with nothing excluded (in particular the benchmark $n = 3$ and the NTK baseline are included). The proposed method is never first: it ranks 3/5, 2/5, 3/5 and 2/5 at $n = 1, 2, 3, 6$. NTK is first at $n = 2$ and $n = 6$ (mean steady-state $\|e\|$ 0.0015 and 0.0013, against 0.0372 and 0.0044 for the proposed method), and the sparse GP is first at $n = 3$ (3.2729), where the error of




Figure 1: Mechanism × geometry on Lorenz tracking (same seeds, same covering grid, $m = 27$): $\|e(t)\|$ against time, logarithmic scale, median of the 5 seeds for projection, σ-modification and e-modification; shaded bands: minimum to maximum over the 5 seeds. Panels: fixed (W), adaptive (W, c), adaptive (W, c, σ). Generated by `experiments/exp_bias_factorial.py`.

the proposed method is 7.7190. Its genuine edge is cost against the GP baselines: the mean update time of the proposed method is 0.10, 0.22 and 0.60 ms per step at $n = 1, 2, 6$ and 4.11 ms at $n = 3$, against 52.92–1888.80 ms per update for the GP baselines, which update every 50–100 steps. NTK takes 1.59–13.22 ms per update; it updates every 50 steps at $n = 1$ and every step at $n = 2, 3, 6$ (`experiments/sim4_ntk_control.py`, `experiments/sim4_ntk_n2.py`, `experiments/sim4_ntk_n3.py`, `experiments/sim4_ntk_n6.py`), so it is cheaper than the proposed method only at $n = 3$ (2.35 against 4.11 ms per step). The comparison is not information-matched: every baseline, here and in the fair re-run below, is trained on noise-free samples of the true $f(x, t)$, which amounts to measuring the state derivative exactly, whereas the adaptive law uses only the tracking error e. With one run per cell no valid significance test is available.

In the original baseline scripts (`sim1`–`sim3`) the GP baselines held a stale prediction for 50–100 steps, and the inducing points of the sparse GP lay on a diagonal line over $x \in [-3, 3]$ while the reference spans ± 20. The fair re-run at $n = 1$ (`experiments/kernel_baselines_fair.py`, `experiments/run_fair_n1.py`, `experiments/plot_fair_n1.py`) uses per-step prediction for all methods, a covering inducing grid and 3 seeds (Table 9, Figure 2). The mean steady-state $\|e\|$ is 0.0591 for the exact GP, 0.1867 for the sparse GP, 0.1978 for the proposed method, 0.3087 for the deep GP and 0.4700 for NTK; the median online step is 6.309, 1.353, 0.060, 4.461 and 0.212 ms respectively. Once it is allowed to predict every step, the exact GP is 3–4× more accurate than the proposed method (mean steady-state $\|e\|$ 0.0591 against 0.1978), and more accurate in every seed (the proposed method has the lower mean in 0/3 seeds). The proposed method is on par with a fixed-grid sparse GP (the proposed method has the lower mean in 2/3 seeds) at 22.6× lower cost (median step 0.060 ms against 1.353 ms); its step is the cheapest of the five methods. This is the version of the comparison to quote.

8.3 The bound against data

Oracle bound study. The oracle experiment (`experiments/exp_oracle_ruub.py`) uses a plant that is exactly representable with constant θ^* ($m = 8$, $n = 2$), so every constant of Theorem 1 is computable; with θ^* constant, $C_\Omega = 0$ and $R_{\mathrm{UUB}} = D_{\max}/k_e$, reported as D_{theory}/k_e with $D_{\max}$ from Lemma 2. The "full-state bound" is $\sqrt{R_{\mathrm{UUB}}^2 + 2V_{\max,\Theta}}$ of Theorem 1(ii), and D_{meas} is the measured $\sup_{t\in[10,30]}\|d(t)\|$ over the evaluation window. Table 10 and Figure 3 cover $k_e \in \{2, 5, 10, 20\}$ and four learning-rate scales, 3 seeds each. The Lyapunov identity of Lemma 3 holds to 3.55×10^{-15} (maximal $|\dot{V} - (-k_e\|e\|^2 + e^\top d + \mathrm{cross})|$), property (P) is never violated, the hard bounds held, and the RMS bound of Theorem 1(iii) with the measured disturbance holds in 48/48 runs and tracks $1/k_e$. The a-priori R_{UUB} is 26.1–261.1 against observed RMS errors of 0.0087–0.0908; the ratio of the a-priori bound to the observed RMS error ranges from

Table 8: Higher-dimensional comparison, complete (nothing excluded). Single trajectory per cell; steady state = last 40% of 30 s. Step time = mean wall-clock time of the method's update call (GP: per update, every 50–100 steps; NTK: per update, every 50 steps at $n = 1$ and every step at $n = 2, 3, 6$; proposed: per step). No valid significance test is available (one run per cell). Generated by `experiments/plot_comparison_hd_complete.py`; source: `results/comparison_complete_table.md`.

benchmark	method	mean SS $\|e\|$	median SS	max SS	rank (mean SS)	mean update time (ms)
$n = 1$ (1-D plant, Lorenz ref.)	Proposed (Proj)	0.2342	0.1496	1.290	3/5	0.10
	Exact GP	0.2206	0.1415	1.016	1/5	273.78
	Sparse GP	0.2285	0.1914	0.874	2/5	52.92
	Deep GP	0.3963	0.2994	1.436	4/5	94.34
	NTK	0.4286	0.3318	1.324	5/5	13.22
$n = 2$ (Van der Pol)	Proposed (Proj)	0.0372	0.0137	0.310	2/5	0.22
	Exact GP	0.0555	0.0370	0.278	3/5	498.50
	Sparse GP	0.1771	0.0929	0.886	5/5	150.30
	Deep GP	0.1758	0.1169	1.097	4/5	415.37
	NTK	0.0015	0.0009	0.006	1/5	1.59
$n = 3$ (Lorenz plant)	Proposed (Proj)	7.7190	5.7197	30.476	3/5	4.11
	Exact GP	10.2774	8.8437	34.467	5/5	1888.80
	Sparse GP	3.2729	2.5099	19.176	1/5	264.64
	Deep GP	9.5344	8.3305	33.755	4/5	802.93
	NTK	5.7642	4.2944	18.240	2/5	2.35
$n = 6$ (coupled Duffing)	Proposed (Proj)	0.0044	0.0013	0.026	2/5	0.60
	Exact GP	0.0841	0.0695	0.279	3/5	668.01
	Sparse GP	0.1444	0.0872	0.670	5/5	523.83
	Deep GP	0.0953	0.0553	0.342	4/5	1607.48
	NTK	0.0013	0.0010	0.004	1/5	4.75

Table 9: Fair 1-D kernel comparison: per-step prediction for every method, covering inducing grid, seed-level statistics. Seeds: 3 (different reference and plant initial conditions). Steady state = last 40% of 30 s. Step time = median full online step (prediction + amortised training) after warm-up, same process type, 2 BLAS threads. Source: `results/fair_n1/table.md`.

method	mean SS $\|e\|$ (mean $\pm$ 95% CI over seeds)	max SS $\|e\|$ (mean)	median step (ms)	mean step (ms)	seeds where proposed has lower mean SS
Proposed (Proj)	0.1978 ± 0.0503	1.147	0.060	0.069	—
Exact GP	0.0591 ± 0.0061	0.780	6.309	13.936	0/3
Sparse GP (9×3 grid)	0.1867 ± 0.0210	0.699	1.353	2.732	2/3
Deep GP	0.3087 ± 0.1121	1.262	4.461	5.495	2/3
NTK	0.4700 ± 0.0250	1.438	0.212	0.503	3/3




Figure 2: 1-D benchmark, fair protocol. Left: $\|e(t)\|$ (median over seeds, 300-step average), logarithmic scale, for the proposed method (Proj), the exact GP, the sparse GP (9×3 grid), the deep GP and NTK. Right: mean steady-state $\|e\|$ ($\pm$ 95% CI over seeds). Generated by `experiments/plot_fair_n1.py`; numbers in Table 9.

915× to 10954×.




Figure 3: Oracle bound study, logarithmic axes. Left: scaling with k_e at learning-rate scale 1.0 — observed RMS $\|e\|$, observed maximal steady-state $\|e\|$, the RMS bound with the measured disturbance, the a-priori R_{UUB} (Lemma 2) and the a-priori full-state bound. Right: scaling with the learning-rate scale at $k_e = 5$ — observed RMS $\|e\|$, the RMS bound with the measured disturbance and the a-priori R_{UUB}. Generated by `experiments/exp_oracle_ruub.py`; numbers in Table 10.

Monte Carlo over initial conditions. On a plant that is not exactly representable only exactly checkable statements are tested (`experiments/exp_monte_carlo_honest.py`, Table 11): all 200 trajectories are finite, the hard parameter bounds held at every recorded step in 200/200 trials, and the a-posteriori ISS bound held in 200/200 trials (maximal ratio 0.979). No a-priori R_{UUB} threshold is drawn: $D_{\max}$ and C_Ω are not computable for this plant (the threshold 0.5 of the original figure was an arbitrary constant, not a calculated bound).

8.4 Summary

- On identical geometry, projection beats σ- and e-modification in every cell and in every seed (5/5), with paired confidence intervals excluding zero.

Table 10: Oracle R_{UUB} study (exactly representable plant, constant θ^*, $m = 8$, $n = 2$). Window $[10, 30]$ s; mean over 3 seeds. All bounds in the same units as $\|e\|$. "P viol" = samples with cross > 0 (must be 0); "inside" = hard bounds held. RMS bound (measured disturbance) satisfied in all 48 runs: True. Maximal identity error $|\dot{V} - (-k_e\|e\|^2 + e^\top d + \mathrm{cross})|$: 3.55×10^{-15}. Source: `results/oracle_ruub/table.md`.

k_e	Γ scale	RMS $\|e\|$	max SS $\|e\|$	RMS bound (D_{meas}, finite window)	D_{meas}/k_e	D_{theory}/k_e	full-state bound	ratio theory/RMS	P viol	inside
2	0.5	0.0908	0.1704	0.6791	0.5435	261.1	261.9	2875×	0	True
2	1	0.0616	0.1215	0.6801	0.6123	261.1	261.5	4239×	0	True
2	2	0.0383	0.0796	0.7139	0.6808	261.1	261.3	6824×	0	True
2	5	0.0238	0.0476	1.1766	1.1649	261.1	261.2	10954×	0	True
5	0.5	0.0533	0.1054	0.3170	0.1979	104.4	106.4	1958×	0	True
5	1	0.0392	0.0708	0.2793	0.2123	104.4	105.4	2664×	0	True
5	2	0.0270	0.0539	0.2601	0.2240	104.4	104.9	3866×	0	True
5	5	0.0157	0.0329	0.2925	0.2780	104.4	104.6	6662×	0	True
10	0.5	0.0398	0.0793	0.1970	0.0981	52.2	56.1	1313×	0	True
10	1	0.0266	0.0519	0.1596	0.1003	52.2	54.2	1965×	0	True
10	2	0.0194	0.0369	0.1395	0.1057	52.2	53.2	2688×	0	True
10	5	0.0117	0.0240	0.1335	0.1188	52.2	52.6	4466×	0	True
20	0.5	0.0285	0.0522	0.1290	0.0501	26.1	33.2	915×	0	True
20	1	0.0197	0.0389	0.0987	0.0494	26.1	29.9	1328×	0	True
20	2	0.0132	0.0260	0.0799	0.0503	26.1	28.0	1971×	0	True
20	5	0.0087	0.0168	0.0676	0.0534	26.1	26.9	3011×	0	True

Table 11: Monte Carlo over random initial conditions (Van der Pol, $m = 16$, $k_e = 5.0$, 200 trials). Source: `results/monte_carlo_honest/summary.md`.

check	result
all trajectories finite	200/200
hard parameter bounds held at every recorded step	200/200
a-posteriori ISS bound $\|e(t)\| \le \mathrm{e}^{-k_e(t-T_1)}\|e(T_1)\| + \sup\|f - \hat{f}\|/k_e$ held	200/200 (max ratio 0.979)
steady-state max $\|e\|$ (last 25%): median / 95th pct / max	0.0568 / 0.0938 / 0.1075
steady-state mean $\|e\|$: median / 95th pct / max	0.0302 / 0.0484 / 0.0593
$\sup\|f - \hat{f}\|$ in steady state / k_e: median / max	0.0627 / 0.1475
transient max $\|e\|$: max over trials	4.903

- Adaptive geometry is a smaller but consistent gain: 0.0116 for adaptive centres and a further 0.0349 for adaptive widths in mean steady-state $\|e\|$, 5/5 seeds each.
- Against kernel-regression baselines the proposed method is never the most accurate. Its edge is cost against the GP baselines (NTK is cheaper only at $n = 3$), and in the fair re-run at $n = 1$ it has the cheapest step of the five methods.
- In the fair re-run at $n = 1$ the exact GP is 3–4$\times$ more accurate than the proposed method.
- In the oracle study the a-priori bound exceeds the observed RMS error by factors of 915 to 10954, while the RMS bound with the measured disturbance holds in 48/48 runs.

Part II
Transport adaptive control

9 Setting

Plant, reference, control, error.

$$\dot{x} = f(x,t) + u, \qquad e = x - x_m(t), \qquad u = \dot{x}_m - k_e e - \hat{f}(x).$$

Lifted parameter. $\vartheta = (a, c, s) \in \mathbb{R}^n \times \mathbb{R}^n \times \mathbb{R}_{>0}$, and $p := 2n + 1$. The feature is $\Phi(x;\vartheta) := a\,\varphi(x;c,s) \in \mathbb{R}^n$, where $\varphi(x;c,s) = \exp\bigl(-\|x-c\|^2/(2s^2)\bigr)$.

Network as a measure.

$$\hat{f}(x) = \frac{1}{m}\sum_{i=1}^{m} \Phi(x;\vartheta_i) = \int \Phi(x;\vartheta)\,\mathrm{d}\mu(\vartheta), \qquad \mu := \frac{1}{m}\sum_{i=1}^{m} \delta_{\vartheta_i}.$$

The map $\mu \mapsto \hat{f}$ is linear.

Metric. $G = \mathrm{diag}(g_a I_n, g_c I_n, g_s) \succ 0$ is the gain matrix. Distances are measured in $\|v\|_{G^{-1}}^2 = v^\top G^{-1} v$, and $\lambda_G := \lambda_{\max}(G)$.

Admissible box and projection.

- **The box.** $\Theta_K := \{\|a\| \le a_{\max}\} \times \{\|c\| \le c_{\max}\} \times [s_{\min}, s_{\max}]$.
- **Layer and inner set.** ε_K is the width of the boundary layer. Shrinking each bound of Θ_K by ε_K gives the inner set Θ_K^ε, with interior $\operatorname{int}\Theta_K^\varepsilon$.
- **The operator.** $\mathrm{Proj}_K(\vartheta, y)$ is the per-particle, factor-wise projection (implemented in `src/core/projection.py` in the accompanying repository). It uses Pomet–Praly norm-ball operators [50] for a and c, with a quadratic ramp $h \in [0,1]$ (defined in Section 4), and a box operator for s, with a linear ramp.
- **Regularity.** The operator is locally Lipschitz in (ϑ, y). The norm-ball factors are C^1 at the inner edge of their layers. The box factor, and every factor at the outer edge, is only Lipschitz. The proofs use only Lipschitz continuity and ramp values in $[0,1]$.

Property (P). For $\vartheta \in \Theta_K$, $\vartheta' \in \Theta_K^\varepsilon$ and all y,

$$\langle \vartheta - \vartheta',\, G^{-1}\,\mathrm{Proj}_K(\vartheta, Gy)\rangle \le \langle \vartheta - \vartheta',\, y\rangle. \tag{P}$$

On $\partial\Theta_K$ the projected velocity has no outward component, so Θ_K is forward invariant (Nagumo; Pomet–Praly 1992 [50]).

Lemma P.1 (Proj_K is non-expansive)**.** *For every $\vartheta = (a,c,s) \in \Theta_K$ and $y = (y_a, y_c, y_s)$,*

$$\|\mathrm{Proj}_K(\vartheta, Gy)\|_{G^{-1}}^2 = \|y\|_G^2 - \sum_{b\in\{a,c,s\}} h_b(2-h_b)\,\frac{\bigl[(n_b^\top G_b y_b)_+\bigr]^2}{n_b^\top G_b n_b} \;\le\; \|y\|_G^2.$$

The terms are as follows.

- *For the balls, $n_a = a$ and $n_c = c$, with the quadratic ramps $h_a = h(\|a\|; a_{\max}, \varepsilon_K)$ and $h_c = h(\|c\|; c_{\max}, \varepsilon_K)$.*

- *For $s < s_{\min} + \varepsilon_K$, $n_s = -1$ and $h_s = 1 - \mathrm{clip}\big((s - s_{\min})/\varepsilon_K, 0, 1\big)$.*
- *For $s > s_{\max} - \varepsilon_K$, $n_s = +1$ and $h_s = 1 - \mathrm{clip}\big((s_{\max} - s)/\varepsilon_K, 0, 1\big)$.*
- *Otherwise $h_s = 0$.*

Proof. G is block-diagonal, so both squared norms split over the blocks. On an active ball block, $\mathrm{Proj}_b = G_b q_b$ with $q_b = y_b - h_b n_b\,(n_b^\top G_b y_b)/(n_b^\top G_b n_b)$. Expanding $q_b^\top G_b q_b$ gives the subtracted term. On an inactive block ($h_b = 0$ or $n_b^\top G_b y_b \le 0$), $\mathrm{Proj}_b = G_b y_b$. The s-block is the one-dimensional case. Finally $h_b \in [0,1]$, so $h_b(2 - h_b) \in [0,1]$. □

Checked by: `tests/test_transport.py`: `test_lemma_P1_projection_is_nonexpansive_and_tight_velocity_bound`.

Potential. $g_t(\vartheta) := e(t)^\top \Phi(x(t);\vartheta)$, and $F_t(\mu) := -\int g_t \,\mathrm{d}\mu = -e(t)^\top \hat{f}(x(t))$.

Adaptive law (the algorithm).

$$\dot{\vartheta}_i = \mathrm{Proj}_K\big(\vartheta_i,\, G\nabla_\vartheta g_t(\vartheta_i)\big), \qquad i = 1, \dots, m. \tag{L}$$

Ideal measure. For $j = 1, \dots, m$, $\vartheta_j^*(t) \in \Theta_K^\varepsilon$ is C^1 in t, and $\mu_t^* := \frac{1}{m}\sum_j \delta_{\vartheta_j^*(t)}$. Write $\varepsilon(x,t) := f(x,t) - \int \Phi(x;\cdot)\,\mathrm{d}\mu_t^*$, with $\|\varepsilon\| \le \varepsilon_{\max}$ on the operating set X, and set

$$V_*(t) := \Big(\frac{1}{m}\sum_j \|\dot{\vartheta}_j^*\|_{G^{-1}}^2\Big)^{1/2}, \qquad V_* := \sup_t V_*(t).$$

The proofs never use optimality of μ^*. It is an analytic device, and any admissible C^1 curve may be used.

Optimal-transport distance. For uniform atomic $\mu = \frac{1}{m}\sum \delta_{\vartheta_i}$ and $\nu = \frac{1}{m}\sum \delta_{\vartheta'_j}$,

$$W(\mu,\nu)^2 := \min_{\pi \in S_m} \frac{1}{m}\sum_i \|\vartheta_i - \vartheta'_{\pi(i)}\|_{G^{-1}}^2.$$

This is the Kantorovich 2-Wasserstein distance with ground cost $\|\cdot\|_{G^{-1}}^2$. For equal masses the optimum is attained at a permutation (Birkhoff). It is computed by linear assignment in $O(m^3)$. For general measures on Θ_K, W denotes the same Kantorovich distance.

Label-free Lyapunov function.

$$V_{\mathrm{OT}}(t) := \tfrac{1}{2}\|e(t)\|^2 + \tfrac{1}{2}W(\mu_t, \mu_t^*)^2 = \min_\pi V_\pi(t),$$
$$V_\pi := \tfrac{1}{2}\|e\|^2 + \frac{1}{2m}\sum_i \|\vartheta_i - \vartheta^*_{\pi(i)}\|_{G^{-1}}^2.$$

Standing hypotheses (S).

(S0) $k_e > 0$; $G = \mathrm{diag}(g_a I_n, g_c I_n, g_s) \succ 0$; $s_{\min} > 0$; and $0 < \varepsilon_K < \min\{a_{\max}, c_{\max}, (s_{\max} - s_{\min})/2\}$ (`src/core/projection.py` enforces the last condition).

(S1) f is continuous in (x,t) and locally Lipschitz in x, uniformly in t on bounded time intervals.

(S2) $x_m \in C^1$, and x_m and $\dot{x}_m$ are bounded.

(S3) $\vartheta_i(0) \in \Theta_K$.

(S4) The ϑ_j^* are C^1 with values in Θ_K^ε, and $V_* < \infty$.

(S5) $x(t) \in X$ on the interval considered. For bounded f, take $X := \bigcup_t \{x : \|x - x_m(t)\| \le r(F)\}$ with $F = \sup\|f\|$ (Theorem D(b)). Then $\varepsilon_{\max}$ is the supremum over this X, and it depends on $\|e(0)\|$.

Under (S0)–(S3) the right-hand side of (L) is continuous in t and locally Lipschitz in (x, ϑ). Solutions are classical (C^1) and unique on their maximal interval. Under (S4) as well, each V_π is C^1.

Notation.

- Times are written $t_0 \le t_1$ (or $T_1 < T_2$), and r is the integration variable in time.
- $\mathrm{RMS}_{[T_1,T_2]}\|e\| := \Big((T_2 - T_1)^{-1} \int_{T_1}^{T_2} \|e\|^2 \,\mathrm{d}r\Big)^{1/2}$.

Constants of the Gaussian. Closed forms (`gaussian_constants` in `src/core/transport.py` of the accompanying repository). With $u := \|x - c\|^2/s^2$ and $\varphi = e^{-u/2}$, on $s \ge s_{\min}$:

$$\|\nabla_c \varphi\| \le J_c := e^{-1/2}/s_{\min}, \qquad |\partial_s \varphi| \le J_s := 2e^{-1}/s_{\min}, \qquad J := (J_c^2 + J_s^2)^{1/2},$$
$$\|\nabla^2_{(c,s)} \varphi\| \le H_\varphi := \big(\max(h_{cc}, h_{ss}) + h_{cs}\big)/s_{\min}^2,$$
$$h_{cc} = 1, \qquad h_{ss} = 2e^{-1/2}, \qquad h_{cs} = \sqrt{u}\, e^{-u/2}|u - 2| \ \text{ evaluated at } u = (5 - \sqrt{17})/2.$$

J and H_φ are upper bounds. The true supremum of $\|\nabla_{(c,s)}\varphi\|$ is about $0.92/s_{\min}$, and that of $\|\nabla^2_{(c,s)}\varphi\|$ is about $1.57/s_{\min}^2$. The Hessian of the lifted potential is the block matrix

$$\begin{pmatrix} 0 & u\,\nabla\varphi^\top \\ \nabla\varphi\, u^\top & (u^\top a)\,\nabla^2\varphi \end{pmatrix}.$$

Hence

$$\sup_{x \in \mathbb{R}^n,\ \vartheta \in \Theta_K,\ \|u\|=1} \|\nabla^2_\vartheta \big(u^\top \Phi(x;\vartheta)\big)\| \le H_K := J + a_{\max} H_\varphi, \qquad K := \tfrac{1}{2} H_K. \tag{1}$$

A brute-force numerical check over the full lifted Hessian gives $\sup \approx 0.76\, H_K$.

10 Theorem A: identification

Theorem A. *Assume* (S0).

(i) Assume scalar classical gains $\Gamma_W = \gamma_W I$, $\Gamma_c = \gamma_c I$ and $\Gamma_\sigma = \gamma_\sigma I$, and use the dictionary $W_i = a_i/m$, $\gamma_W = g_a/m$, $\gamma_c = m g_c$, $\gamma_\sigma = m g_s$.

- *The* raw (*unprojected*) *rates of* (L) *and of the classical law of Section 4 coincide.*
- *The* projected *rates coincide when the classical joint projections are replaced by per-kernel ones. The classical law projects W onto one Frobenius ball and* $\mathrm{vec}(c)$ *onto one Euclidean ball. The per-kernel version projects:*
  - *W_i onto $\|W_i\| \le a_{\max}/m$, with layer ε_K/m;*
  - *c_i onto $\|c_i\| \le c_{\max}$, with layer ε_K;*
  - *σ_i onto $[s_{\min}, s_{\max}]$, with layer ε_K.*
- *For $m \ge 2$, no joint ball, of any radius $W_{\max}$ and layer ε_W, reproduces the per-kernel field. For $m = 1$ the two coincide, with $W_{\max} = a_{\max}$ and layer ε_K.*

(ii) Fix a closed-loop trajectory $(x(\cdot), e(\cdot))$. Then:

- *The field* $v_t(\vartheta) := \mathrm{Proj}_K(\vartheta, G\nabla g_t(\vartheta))$ *is Lipschitz in* ϑ *on* Θ_K*, and* Θ_K *is forward invariant.*
- *For every* $\mu_0 \in \mathcal{P}(\Theta_K)$*,* $\mu_t := (X_t)_{\#}\mu_0$ *is a distributional solution of* $\partial_t\mu + \nabla\cdot(\mu v_t) = 0$*.*
- *On* $\operatorname{int}\Theta_K^{\varepsilon}$*,* $v_t = -G\nabla(\delta F_t/\delta\mu)$*. So at frozen time,* μ_t *is the Wasserstein gradient flow (metric* G^{-1}*) of the linear functional* F_t*, and* (L) *is its particle form.*
- F_t *is generated by the closed loop, so the coupled system is the one studied in Theorem D.*

(iii) ***Dictionary****, all in the* G^{-1} *metric:*

classical device	measure-level object
projection onto Θ_K	A velocity-level (Pomet–Praly) regularisation of the support constraint $\mu_t \in \mathcal{P}(\Theta_K)$. For $\varepsilon_K > 0$ it is not the gradient flow of any functional of μ. Formally, as $\varepsilon_K \to 0$, the field tends to the G^{-1}-projection of $G\nabla g_t$ onto the tangent cone of Θ_K, which is the velocity of the $W_{G^{-1}}$-gradient flow of $F_t + \iota_{\mathcal{P}(\Theta_K)}$ ([33, Prop. 3.1] for the exact W_2 projection).
σ-modification in the G^{-1} metric (leak outside the gain, $\dot\vartheta = G\nabla g - \sigma(\vartheta - \vartheta^0)$, as in ($\mathrm{L}_\sigma$) and Theorem H)	the potential energy $(\sigma/2)\int\|\vartheta - \vartheta^0\|^2_{G^{-1}}\,\mathrm{d}\mu$, which is geodesically σ-convex
classical σ-modification $\Gamma(y - \sigma\theta)$ (leak inside the gain; `src/baselines/sigma_mod.py` implements its W-block, with c and σ frozen)	For the full-vector leak, under (i), the object is $(\sigma/2)\int(\vartheta - \vartheta^0)^\top D(\vartheta - \vartheta^0)\,\mathrm{d}\mu$ with $D = \mathrm{diag}(I_n/m, mI_n, m)$. In the G^{-1} metric it is λ-convex with $\lambda = \sigma\cdot\min(g_a/m, mg_c, mg_s)$, which depends on m. For the W-block alone it is $(\sigma/2)\int(a - a^0)^\top(I_n/m)(a - a^0)\,\mathrm{d}\mu$.
σ-modification toward a labelled (ϑ_i^0)	not a functional of μ; it is a functional on the labelled lift $\mathbb{R}^{mp}$
OT σ-modification (Theorem C)	$(\sigma/2)W(\mu, \mu^0)^2$. For atomic μ with a unique optimal assignment, its W-gradient at atom i is $\sigma(\vartheta_i - \vartheta^0_{\tau(i)})$. At ties, ($\mathrm{L}_\sigma$) uses a measurable selection from the superdifferential $\{\sigma(\vartheta_i - \vartheta^0_{\tau(i)}) : \tau \text{ optimal}\}$.
e-modification	the same functionals with coefficient $\sigma\|e\|$
parameter dither (Theorem H)	the entropy $\tau\int\mu\log\mu$

Status. Proved and checked. Novelty (literature audit): interpretive; ingredients known [8], [13], [33], [45].

Checked by: `tests/test_transport.py`: `test_identification_with_classical_law` (raw rates in (i));
`test_identification_projected_rates_per_kernel` (projected rates and the only-if construction in (i)).

Proof. **(i) Gain map.** We have $\nabla_a g = \varphi e$, $\nabla_c g = (a^\top e)\nabla_c\varphi$ and $\partial_s g = (a^\top e)\partial_s\varphi$. Matching these with the classical rates gives the gain map.

Scaling identity. $(1/m)\,\mathrm{Proj}_{r,\varepsilon}(a, y) = \mathrm{Proj}_{r/m,\varepsilon/m}(a/m, y/m)$. Three facts give it:

- $h_{r/m,\varepsilon/m}(\|a\|/m) = h_{r,\varepsilon}(\|a\|)$;
- the direction $a(a^\top y)/\|a\|^2$ is homogeneous of degree 0 in a and of degree 1 in y;

- the activation $a^\top y > 0$ is invariant under scaling.

If the layer is not rescaled, the ramps differ.

Only if $(m \geq 2)$. Let $r_{\text{in}} := (a_{\max} - \varepsilon_K)/m$, and let the joint ball have radius $W_{\max}$, layer ε_W and $\rho := W_{\max} - \varepsilon_W$. In both cases below the driving term points outward.

- If $\rho < \sqrt{2}\, r_{\text{in}}$, take two rows of equal norm r with $\sqrt{2}\, r \in \big(\rho, \min(\sqrt{2}\, r_{\text{in}}, W_{\max})\big]$, parallel to e. The per-kernel field is inactive and the joint field is active.
- If $\rho \geq \sqrt{2}\, r_{\text{in}}$, take one row with norm in $\big(r_{\text{in}}, \min(a_{\max}/m, \rho)\big)$ and the others zero. The per-kernel field is active and the joint field is inactive.

The same argument applies to $\mathrm{vec}(c)$. For $m = 1$ the two operators coincide.

The test `test_identification_projected_rates_per_kernel` checks three things: the identity with active projections; that the layer must be rescaled; and the only-if construction against `ProjectionOperator.project_matrix` on five joint balls. Four are active against an inactive per-kernel field, and one is inactive against an active per-kernel field.

(ii) Proj_K is Lipschitz in ϑ. The ramp vanishes for $\|\vartheta\| \leq \theta_{\max} - \varepsilon_K$, so the direction $\vartheta/\|\vartheta\|$ is evaluated only away from 0. Also, ∇g_t is C^1 on $s \geq s_{\min}$.

- **Invariance.** Forward invariance follows from Nagumo's theorem, since $n_b^\top \mathrm{Proj}_b \leq 0$ on the boundary, with the outward normals n_b of Lemma P.1.
- **Continuity equation.** The push-forward of μ_0 by a Lipschitz flow solves the continuity equation (Golse [24]; Ambrosio–Gigli–Savaré [2, §8.1]).
- **Gradient form.** On $\operatorname{int} \Theta_K^\varepsilon$ the projection is inactive, so $v_t = G\nabla g_t = -G\nabla(\delta F_t/\delta\mu)$.

(iii) Potential-energy rows. These follow from the computation in the metric G^{-1}: for any V, $\nabla_{G^{-1}} V = G\nabla V$. For example, $V = (\sigma/2)\|\vartheta - \vartheta^0\|_{G^{-1}}^2$ gives $G\nabla V = \sigma(\vartheta - \vartheta^0)$.

Classical-leak row. Applying the gain map of (i) to the classical leak $\Gamma(y - \sigma\theta)$ gives the velocity $-\sigma GD(\vartheta - \vartheta^0)$.

Projection row. The G^{-1}-rescaled correction field has an asymmetric Jacobian. Take the example $m = 1$, $n = 1$, $G = I$, $x = 0.7$, $e = 1$, $\vartheta = (0.93, 0.2, 1.5)$, $a_{\max} = 1$, $\varepsilon_K = 0.15$ (so $h(0.93) = 0.284$) and $c_{\max} = 2$, with a in the layer. A numerical evaluation gives:

- the c-derivative of the a-component is -0.060;
- the a-derivative of the c-component is 0.

So the field is not a gradient. □

Remark A.1 (normalisation)**.** The algorithm's raw rates are unchanged. Two things do change: the admissible set becomes a product of per-kernel balls, and $a_{\max}$, not $\|W\|_F$, is held fixed in m.

- ***m*-free constants.** In this normalisation the structural constants k_e, K, L_0, L_1, λ_G and Δ are free of m for fixed G and $a_{\max}$. So are the constants of Theorems B, C(i)–(iii), D and F.
- ***m*-dependent data.** $\varepsilon_{\max}$ and V_* are data of the chosen m-atom ideal. The local constants of C(iv) and of Theorem E depend on m.
- **Classical convention.** Fixed classical γ's correspond to $g_a = m\gamma_W \propto m$ and $g_c, g_s \propto 1/m$. Weights speed up, centres and widths freeze, $\lambda_G \propto m$ and $\Delta \propto \sqrt{m}$.

11 Theorem B: the optimal-assignment Lyapunov function

Lemma B.1 (Dini derivative of a minimum)**.** *Suppose each V_π is C^1 and $V_{\text{OT}} = \min_\pi V_\pi$. Then V_{OT} is locally Lipschitz, and $D^+ V_{\text{OT}}(t) \leq \dot{V}_\pi(t)$ for every π active at t.*

Proof. For $h > 0$, $V_{\mathrm{OT}}(t+h) \le V_\pi(t+h)$, and $V_{\mathrm{OT}}(t) = V_\pi(t)$. □

Lemma B.2 (matched remainder in two regimes; gradient bound)**.** *Let* $L_0 := 2a_{\max}$ *and* $L_1 := (1 + a_{\max}^2 J^2)^{1/2}$. *For* $\vartheta, \vartheta' \in \Theta_K$*:*

(a) the remainder $R := g_t(\vartheta') - g_t(\vartheta) - \langle \nabla g_t(\vartheta),\, \vartheta' - \vartheta \rangle$ *satisfies*

$$|R| \le \|e(t)\| \cdot \min\bigl\{K\|\vartheta - \vartheta'\|^2,\ L_0 + L_1\|\vartheta - \vartheta'\|\bigr\};$$

(b) $\|\nabla_\vartheta g_t(\vartheta)\| \le L_1\|e(t)\|$.

Proof. (b) We have $\nabla_a g = \varphi e$ and $\nabla_{(c,s)} g = (a^\top e)\nabla_{(c,s)}\varphi$, with $\varphi \le 1$ and $\|\nabla_{(c,s)}\varphi\| \le J$ on $s \ge s_{\min}$. So $\|\nabla g\|^2 \le \|e\|^2(1 + a_{\max}^2 J^2)$.

(a) If $e(t) = 0$, both sides vanish. Otherwise set $h := \vartheta' - \vartheta$.

- *Quadratic branch.* Θ_K is convex, so the Lagrange remainder of $\tau \mapsto g_t(\vartheta + \tau h)$ gives $R = \frac{1}{2} h^\top \nabla^2 g_t(\vartheta + \tau_0 h) h$ for some $\tau_0 \in (0,1)$. Since $g_t = \|e\| \cdot (u^\top \Phi)$ with $u = e/\|e\|$, the Hessian bound (1) of Section 9 gives this branch.
- *Linear branch.* It follows from $|g_t| \le \|e\| a_{\max}$ and (b). □

Theorem B. *Under (S), along every solution of (L), for every* t*, with* $W := W(\mu_t, \mu_t^*)$*:*

$$D^+ V_{\mathrm{OT}} \le -k_e\|e\|^2 + \|e\|\Bigl(\varepsilon_{\max} + \min\bigl\{K\lambda_G W^2,\ L_0 + L_1 \lambda_G^{1/2} W\bigr\}\Bigr) + V_*(t)\, W. \tag{B}$$

Status. Proved and checked, including a moving ideal, $\varepsilon \neq 0$, switches, an active projection, and the failure of the plain bound along (L_σ). Novelty (literature audit): packaging new; ingredients known [55], [1], [44], [10].

Checked by: `tests/test_transport.py`: `test_theorem_B_inequality_along_trajectories`, `test_theorem_B_moving_ideal_nonrepresentable_switches_and_active_projection`, `test_dini_derivative_of_V_OT_bounded_by_active_branch`, `test_permuted_ideal_labelled_vs_OT`; `experiments/exp_transport_permutation.py`.

Proof. Fix t and an optimal π, and write $\tilde\vartheta_i := \vartheta_i - \vartheta^*_{\pi(i)}$. Then

$$\dot V_\pi = e^\top \dot e + \frac{1}{m}\sum_i \langle \tilde\vartheta_i,\, G^{-1}\dot\vartheta_i\rangle - \frac{1}{m}\sum_i \langle \tilde\vartheta_i,\, G^{-1}\dot\vartheta^*_{\pi(i)}\rangle.$$

(a) $e^\top \dot e = -k_e\|e\|^2 + e^\top \varepsilon + \frac{1}{m}\sum_i \bigl[g_t(\vartheta^*_{\pi(i)}) - g_t(\vartheta_i)\bigr]$.

(b) By (P) with $\vartheta' = \vartheta^*_{\pi(i)}$, $\frac{1}{m}\sum_i \langle \tilde\vartheta_i,\, G^{-1}\dot\vartheta_i\rangle \le \frac{1}{m}\sum_i \langle \tilde\vartheta_i,\, \nabla g_t(\vartheta_i)\rangle$.

(c) Since π is a bijection and is optimal,

$$-\frac{1}{m}\sum_i \langle \tilde\vartheta_i,\, G^{-1}\dot\vartheta^*_{\pi(i)}\rangle \le \Bigl(\frac{1}{m}\sum_i \|\tilde\vartheta_i\|^2_{G^{-1}}\Bigr)^{1/2}\Bigl(\frac{1}{m}\sum_i \|\dot\vartheta^*_{\pi(i)}\|^2_{G^{-1}}\Bigr)^{1/2} = W \cdot V_*(t).$$

Summing gives

$$\dot V_\pi \le -k_e\|e\|^2 + \|e\|\varepsilon_{\max} + \frac{1}{m}\sum_i R_i + W\, V_*(t),$$

where each R_i is a matched remainder. Average Lemma B.2(a), using three facts:

- $\frac{1}{m}\sum_i \min\{A_i, B_i\} \le \min\{\mathrm{avg}\, A, \mathrm{avg}\, B\}$;
- $\frac{1}{m}\sum_i \|\tilde\vartheta_i\|^2 \le \lambda_G W^2$;
- $\frac{1}{m}\sum_i \|\tilde\vartheta_i\| \le \lambda_G^{1/2} W$.

Lemma B.1 then completes the proof. □

Corollary B.3 (ultimate bounds)**.** *Let* $\Delta := \operatorname{diam}_{G^{-1}}(\Theta_K)$*, so* $W \le \Delta$*. Define*

$$D^{\mathrm{OT}} := \varepsilon_{\max} + \min\{K\lambda_G\Delta^2,\ L_0 + L_1\lambda_G^{1/2}\Delta\}, \qquad C^{\mathrm{OT}} := V_*\Delta,$$
$$R^{\mathrm{OT}} := \frac{D^{\mathrm{OT}}}{2k_e} + \Big(\frac{(D^{\mathrm{OT}})^2}{4k_e^2} + \frac{C^{\mathrm{OT}}}{k_e}\Big)^{1/2}.$$

Then:

- $D^+V_{\mathrm{OT}} < 0$ *whenever* $\|e\| > R^{\mathrm{OT}}$.
- *Under (S5) on* $[0,\infty)$, $\limsup\|e\| \le \big((R^{\mathrm{OT}})^2 + \Delta^2\big)^{1/2}$. *This uses* $\sup \frac{1}{2}W^2 \le \Delta^2/2$.
- $\mathrm{RMS}_{[T_1,T_2]}\|e\| \le \dfrac{D^{\mathrm{OT}}}{2k_e} + \Big(\dfrac{(D^{\mathrm{OT}})^2}{4k_e^2} + \dfrac{C^{\mathrm{OT}}}{k_e} + \dfrac{V_{\mathrm{OT}}(T_1)}{k_e(T_2-T_1)}\Big)^{1/2}$.

Proof. As in Theorem 1 (Section 5). The Dini comparison lemma, applied with the continuous, V-independent right-hand side, gives the integrated form. □

Remark B.4 (what OT buys and what it does not)**.**

- **Globally, the OT constant is not smaller than the classical one.** Take the test box: $a_{\max} = 8$, $c_{\max} = 4$, $s_{\min} = 0.3$, $s_{\max} = 3$, $G = \operatorname{diag}(2, 1, 0.5)$ and $F_{\max} := a_{\max}$. Compare constants on the same per-kernel admissible set.

constant	value
classical Lemma 2 (Section 5), $F_{\max} + 3a_{\max} + 2a_{\max}c_{\max}J_c + a_{\max}(s_{\max} - s_{\min})J_s$	214, for every m
OT linear branch, crude form $L_0 + L_1\lambda_G^{1/2}\Delta$	533
OT linear branch, block split through W	382
OT linear branch, bounded per particle by $\|e\|\big(2a_{\max} + \|\tilde{a}_i\| + a_{\max}(J_c\|\tilde{c}_i\| + J_s\|\tilde{s}_i\|)\big)$	214

  Bounded per particle, the OT constant equals the classical one. An earlier comparison figure of "101–138" used a joint centre ball of radius $c_{\max}$. That radius is $\sqrt{m}$ times smaller than the radius $\sqrt{m}\,c_{\max}$ of the ball that circumscribes the per-kernel product of Theorem A.

- **Locally, the excess disturbance is quadratic in the matched error** (Corollary B.5).
- **Labels.** A labelled theory can make the same local statement only for a labelling fixed in advance, and only while the optimal assignment does not switch. (B) holds uniformly, with the minimum over π taken at each t.
- m**-uniformity** is a property of the normalisation (Remark A.1), not of OT.

Corollary B.5 (conditional bound)**.** *Along a solution of (L), suppose* $\sup_{[T_1,T_2]} W(\mu_t, \mu_t^*) \le \bar{\eta}$*. This quantity is available in oracle simulations, since it requires* μ_t^**, or it can be controlled a priori by Theorem E. With* $\bar{D} := \varepsilon_{\max} + \min\{K\lambda_G\bar{\eta}^2,\ L_0 + L_1\lambda_G^{1/2}\bar{\eta}\}$*,*

$$\mathrm{RMS}_{[T_1,T_2]}(\|e\|) \le \frac{\bar{D}}{2k_e} + \Big(\frac{\bar{D}^2}{4k_e^2} + \frac{V_*\bar{\eta}}{k_e} + \frac{V_{\mathrm{OT}}(T_1)}{k_e(T_2-T_1)}\Big)^{1/2}. \tag{11.1}$$

As $\bar{\eta} \to 0$ *and* $T_2 \to \infty$*, this tends to* $\varepsilon_{\max}/k_e$*, the value of the frozen-parameter error channel.*

Variant for (L_σ). *Take a solution of* (L_σ) *in the sense of Section 12 with* $\sup_{[T_1,T_2]} W(\mu_t,\mu_t^*) \le \bar\eta$ *and* $4\bar\eta(b+\bar\eta) < \Delta_{\mathrm{gap}}$. *For example,* $\bar\eta$ *can come from Theorem C(iv). Then, with* $\bar D := \varepsilon_{\max} + K\lambda_G\bar\eta^2$,

$$\mathrm{RMS}_{[T_1,T_2]}(\|e\|) \le \frac{\bar D}{2k_e} + \Big(\frac{\bar D^2}{4k_e^2} + \frac{V_*\bar\eta + \sigma_m b^2/4}{k_e} + \frac{V_{\mathrm{OT}}(T_1)}{k_e(T_2-T_1)}\Big)^{1/2}. \tag{11.2}$$

Proof of the variant. Apply Theorem C(iii) with $C = b(t)^2 \le b^2$, and use $-\sigma_m\eta^2 + \sigma_m\eta b \le \sigma_m b^2/4$. Without the gap condition, replace b^2 by $(b+2\bar\eta)^2$ (from $\sqrt{C} \le b + 2\eta$ in C(ii)), or by $b^2 + 4\bar\eta R$, with $R := \Delta$ as in Theorem C. □

As $\bar\eta \to 0$ and $T_2 \to \infty$ the variant tends formally to $\varepsilon_{\max}/(2k_e) + \big(\varepsilon_{\max}^2/(4k_e^2) + \sigma_m b^2/(4k_e)\big)^{1/2}$. Along (L_σ) with $b > 0$, the measured $\bar\eta$ stays of order b (about $0.9b$ in the test instance), so the bound is first order in b. The plain bound of Corollary B.5 is false along (L_σ). In the test instance, every hypothesis of C(iv) is asserted and $\sup W \le \bar\eta$ is measured. There, the bound of Corollary B.5 is exceeded by more than 3×, while the variant holds (`tests/test_transport.py::test_corollary_B5_fails_along_lsigma_and_its_leak_variant_holds`). Independent numerical checks found violations of 3.8–8.3×.

Status. The bound along (L) is not checked in the repository; its failure along (L_σ) and the (L_σ) variant are checked (Appendix A).

Corollary B.6 (approximation error as quantisation error)**.** *Suppose* $f = \int \Phi\,\mathrm{d}\mu_\infty^* + \varepsilon_\infty$ *for a probability measure* μ_∞^* *on* Θ_K^ε, *with* $\|\varepsilon_\infty\| \le \bar\varepsilon_\infty$. *Then for every* m*-atom* μ_m^* *supported in* Θ_K^ε,

$$\varepsilon_{\max}(m) \le \bar\varepsilon_\infty + L_1\lambda_G^{1/2}\,W(\mu_m^*,\mu_\infty^*).$$

Status. Not checked in the repository (Appendix A).

Proof. For any coupling γ,

$$\Big\|\int \Phi\,\mathrm{d}(\mu_m^* - \mu_\infty^*)\Big\| \le \int \|\Phi(\vartheta) - \Phi(\vartheta')\|\,\mathrm{d}\gamma \le L_1\lambda_G^{1/2}\int \|\vartheta - \vartheta'\|_{G^{-1}}\,\mathrm{d}\gamma$$
$$\le L_1\lambda_G^{1/2}\Big(\int \|\vartheta - \vartheta'\|_{G^{-1}}^2\,\mathrm{d}\gamma\Big)^{1/2}.$$

Minimise over γ. The block constant $\big(g_a + a_{\max}^2(J_c^2 g_c + J_s^2 g_s)\big)^{1/2}$ may replace $L_1\lambda_G^{1/2}$. □

Rates.

(a) **First order.** Let μ_∞^* be compactly supported and $p \ge 3$. Then there are equal-weight μ_m^*, with atoms (cell barycentres) in Θ_K^ε, such that $W(\mu_m^*,\mu_\infty^*) = O(m^{-1/p})$. This follows from an equal-mass dyadic decomposition (Chevallier 2018 [7]). For $p > 4$, i.i.d. atoms also achieve the rate in expectation (Fournier–Guillin 2015 [18]). If μ_∞^* has a non-trivial absolutely continuous part, the order cannot be improved, even with free weights (Zador; Graf–Luschgy [25, Thm. 6.2 with Lemma 3.4]).

(b) **Second order.** Suppose μ_m^* is a centroid quantiser for an optimal plan; a best equal-weight approximation is one. Then the first-order term cancels, and by the Hessian bound (1) of Section 9 and convexity of Θ_K^ε,

$$\varepsilon_{\max}(m) \le \bar\varepsilon_\infty + \min\big\{K\lambda_G W^2,\ L_1\lambda_G^{1/2}W\big\} = \bar\varepsilon_\infty + O(m^{-2/p}).$$

(c) **Free of** p**.** I.i.d. atoms drawn from μ_∞^* give, with positive probability, an m-atom ideal with

$$\varepsilon_{\max}(m) \le \bar\varepsilon_\infty + O\Big(a_{\max}\big(n\log(m\cdot\operatorname{diam} X)/m\big)^{1/2} + a_{\max}/(s_{\min}\sqrt{m})\Big).$$

This is Maurey's empirical method; the constants are not optimised.

The exponent $1/p$ is therefore a property of controlling $\varepsilon_{\max}$ through W, as Theorem D(e) requires. It is not a property of the approximation error itself.

Remark B.7 (continuum form, via couplings)**.** Let $\mu_0 \in \mathcal{P}(\Theta_K)$ and $\mu_t = (X_t)_{\#}\mu_0$, and let μ_t^* be an absolutely continuous ideal curve in $(\mathcal{P}(\Theta_K^\varepsilon), W)$ with velocity field w_t. Set $V_* := \operatorname{ess\,sup}_t \|w_t\|_{L^2(\mu_t^*;G^{-1})}$. Its smallest admissible value is the metric derivative $|\dot\mu_t^*|_W$ (AGS [2, Thm. 8.3.1], after the isometry $\vartheta \mapsto G^{-1/2}\vartheta$). Then (B) holds for a.e. t.

Status. Not checked in the repository (Appendix A).

Sketch.

1. Push an optimal plan γ_t forward by $(X_{t,t+h}, Y_{t,t+h})$. Here Y is the flow of w, or the superposition measure of AGS [2, Thm. 8.2.1] when w is not Lipschitz.
2. This couples (μ_{t+h}, μ_{t+h}^*), so $D^+\big(\tfrac12 W^2\big) \le \int \langle \vartheta - \vartheta',\, v_t(\vartheta) - w_t(\vartheta')\rangle_{G^{-1}}\, \mathrm{d}\gamma_t$.
3. Integrate (P) and Lemma B.2 against γ_t. □

12 Theorem C: optimal-transport σ-modification

Let $\mu^0 = \frac{1}{m}\sum_k \delta_{\vartheta_k^0}$ be a prior configuration with $\vartheta_k^0 \in \Theta_K^\varepsilon$. The *OT σ-modified law* is

$$\begin{aligned} \dot\vartheta_i &= \mathrm{Proj}_K\big(\vartheta_i,\ G\nabla g_t(\vartheta_i) - \sigma_m(\vartheta_i - \vartheta^0_{\tau(i)})\big), \\ \tau(t) &\in \operatorname*{argmin}_{\rho}\ \frac{1}{m}\sum_i \|\vartheta_i - \vartheta^0_{\rho(i)}\|^2_{G^{-1}}. \end{aligned} \tag{L_σ}$$

Solution concept. The right-hand side of (L_σ) is discontinuous at assignment ties.

- **Selection solutions.** A solution is an absolutely continuous (x, ϑ) such that, for a.e. t, $\dot\vartheta_i = \mathrm{Proj}_K\big(\vartheta_i,\ G\nabla g_t(\vartheta_i) - \sigma_m(\vartheta_i - \vartheta^0_{\tau(t)(i)})\big)$ for some $\tau(t)$ optimal at $\vartheta(t)$. This is what `src/core/transport.py` in the accompanying repository integrates.
- **Filippov solutions.** These exist by Filippov's theorem, since the right-hand side is measurable and locally bounded. But they include sliding motions on repelling ties, along which (i) below fails. For example, take ideal = prior = $\{a = 0,\ a = 1\}$, both particles at $a = \frac12$ and $e \equiv 0$: this is a stationary Filippov solution.
- **What holds for which class.**
  - Parts (ii)–(iv) hold for every Filippov solution, with C taken as the maximum over optimal τ.
  - Part (i) and the exact decay hold for selection solutions.
  - For Filippov solutions, (i) weakens as stated below.
- **Regularity and existence.** V_{OT} and Z are absolutely continuous along solutions, and every inequality holds for a.e. t. Existence of selection solutions is proved for the pure leak (Lemma C.3). In general it is assumed: two-way ties are never attracting, by monotonicity of Proj_K, but multi-way ties, the degenerate ramp $h = 1$ and accumulating switch times are not handled.

Notation.

- $c_\rho(\vartheta) := \frac{1}{m}\sum_i \|\vartheta_i - \vartheta^0_{\rho(i)}\|^2_{G^{-1}}$.
- $\eta := W(\mu_t, \mu^*_t)$, and $R := \Delta = \mathrm{diam}_{G^{-1}}(\Theta_K)$.
- $b(t) := W(\mu^*_t, \mu^0)$, and $b := \sup_t b(t)$.
- $\Delta_{\mathrm{gap}}(t) := \min\{c_\rho(\vartheta^*(t)) - b(t)^2 : c_\rho(\vartheta^*(t)) > b(t)^2\}$, with $\min\emptyset := +\infty$; and $\Delta_{\mathrm{gap}} := \inf_t \Delta_{\mathrm{gap}}(t)$.
- $A_{\tau\tau'} := \frac{1}{m}\sum_i \|\vartheta^0_{\tau(i)} - \vartheta^0_{\tau'(i)}\|^2_{G^{-1}}$.

Theorem C. *Under* (S)*, along every solution of* (L_σ) *in the sense above:*

(i) Dissipation toward the prior (selection solutions). *Let* $Z := \frac{1}{2}W(\mu_t, \mu^0)^2$ *and* $\Gamma := \big(\frac{1}{m}\sum_i \|G\nabla g_t(\vartheta_i)\|^2_{G^{-1}}\big)^{1/2} \le \lambda_G^{1/2} L_1 \|e\|$. *Then:*

- $D^+Z \le -2\sigma_m Z + (2Z)^{1/2}\Gamma(t)$;
- *hence* $W(\mu_t, \mu^0) \le \max\{W(\mu_0, \mu^0),\ \sup\Gamma/\sigma_m\}$;
- *for the pure-leak flow* ($\Gamma \equiv 0$)*, the decay* $W(\mu_t, \mu^0) = e^{-\sigma_m t} W(\mu_0, \mu^0)$ *is exact for every* $G \succ 0$ (*Lemma C.3*).

For a Filippov solution with weights θ_k *on tied optimal assignments* τ_k*, the bound weakens to*

$$D^+Z \le -2\sigma_m Z + \frac{\sigma_m}{2}\min_\tau \sum_k \theta_k A_{\tau\tau_k} + (2Z)^{1/2}\Gamma,$$

with the minimum over assignments τ *optimal at* t.

(ii) Induced coupling. *Let* π^* *be optimal from* μ *to* μ^**, and let* τ *be the optimal assignment from* μ *to* μ^0 *that the law uses at* t (*for Filippov solutions, any optimal* τ). *Set* $C := \frac{1}{m}\sum_i \|\vartheta^*_{\pi^*(i)} - \vartheta^0_{\tau(i)}\|^2_{G^{-1}}$. *Then*

$$C \le b(t)^2 + 4\eta R \qquad \text{and} \qquad \sqrt{C} \le b(t) + 2\eta,$$

and $C = b(t)^2$ *whenever* $4\eta(b(t) + \eta) < \Delta_{\mathrm{gap}}$.

(iii) Lyapunov inequality. *With* C *as in* (ii) (*for Filippov solutions, its maximum over optimal* τ),

$$D^+V_{\mathrm{OT}} \le -k_e\|e\|^2 + \|e\|\big(\varepsilon_{\max} + K\lambda_G\eta^2\big) + V_*\eta - \sigma_m\eta^2 + \sigma_m\eta C^{1/2}.$$

Under $4\eta(b+\eta) < \Delta_{\mathrm{gap}}$ *this becomes*

$$D^+V_{\mathrm{OT}} \le -k_e\|e\|^2 + \|e\|\big(\varepsilon_{\max} + K\lambda_G\eta^2\big) + V_*\eta - \frac{\sigma_m}{2}\eta^2 + \frac{\sigma_m}{2}b^2.$$

(iv) Local exponential estimate. *Suppose* $4\bar\eta(b+\bar\eta) < \Delta_{\mathrm{gap}}$ *and* $8K^2\lambda_G^2\bar\eta^2 \le k_e\sigma_m$. *Set* $\alpha := \min(k_e, \sigma_m/2)$ *and* $\beta := \varepsilon^2_{\max}/k_e + (\sigma_m/2)b^2 + 2V_*^2/\sigma_m$, *and suppose* $\max\{V_{\mathrm{OT}}(0), \beta/\alpha\} < \bar\eta^2/2$. *Then:*

- $\{V_{\mathrm{OT}} \le c\}$ *is forward invariant for every* $c \in \big[\max\{V_{\mathrm{OT}}(0), \beta/\alpha\},\ \bar\eta^2/2\big)$;
- $V_{\mathrm{OT}}(t) \le e^{-\alpha t}V_{\mathrm{OT}}(0) + \beta/\alpha$;
- *on this set* $\|e\| < \bar\eta$, *so* (S5) *holds with* $X := \bigcup_t \{x : \|x - x_m(t)\| \le \bar\eta\}$.

(v) The labelled leak. *Define* $b_{\mathrm{lab}}(\pi)(t) := \big(\frac{1}{m}\sum_i \|\vartheta^*_{\pi(i)}(t) - \vartheta^0_i\|^2_{G^{-1}}\big)^{1/2}$ *and* $b_{\mathrm{lab}}(\pi) := \sup_t b_{\mathrm{lab}}(\pi)(t)$.

- *For any fixed π, the labelled σ-modification (leak toward ϑ_i^0) satisfies the second display of* (iii) *at every state, with no gap condition. Here b is replaced by $b_{\mathrm{lab}}(\pi)$, V_{OT} by V_π, and η by the π-labelled error.*
- *Choosing π to minimise $b_{\mathrm{lab}}(\pi)$ gives the bias $\min_\pi \sup_t b_{\mathrm{lab}}(\pi)(t) \geq b$. Equality holds when one assignment is optimal for (μ_t^*, μ^0) at every t, for example a static ideal, or whenever $\Delta_{\mathrm{gap}} > 0$.*
- *The labelled leak does* ***not*** *satisfy* (iii) *for V_{OT} (Proposition C.5).*

Status. Proved and checked; the instance used to check (iv) satisfies every hypothesis of (iv). Not checked in the repository: the Filippov form of (i) and the inequality of (v). Novelty (literature audit): novel as a control construction; C(i) and the obstruction are known [53], [5].

Checked by: `tests/test_transport.py`:
(i) and Lemma C.3 (unprojected pure leak): `test_theorem_C_i_exact_dissipation_pure_leak`;
(ii): `test_lemma_C_ii_induced_coupling_bound`;
(iv): `test_theorem_C_iv_exponential_estimate`.
(v) and Proposition C.5, in `experiments/`: `exp_ot_sigma_mod.py` and `exp_sigma_mod_transient.py`.

Scope.

- **Finite gap.** If $\Delta_{\mathrm{gap}} < \infty$, that is, not all ideal–prior assignments tie, then $\Delta_{\mathrm{gap}} \leq 2R^2/m$ always. Indeed, if every transposition of an optimal ρ_0 were cost-neutral, the matrix $S(i,j) := B(i,j) - (B(i,i) + B(j,j))/2$ built from $B(i,j) := \langle \vartheta_i^*, \vartheta^0_{\rho_0(j)} \rangle_{G^{-1}}$ would be antisymmetric, and every assignment would tie.
  - Hence (iv) needs $\bar\eta < \sqrt{\Delta_{\mathrm{gap}}}/2 \leq R/\sqrt{2m}$.
  - $\beta/\alpha < \bar\eta^2/2$ forces $b < \bar\eta/\sqrt{2}$.
  - So (iv) is a near-prior local result of size $O(m^{-1/2})$ in W, and $O(1/m)$ in V_{OT}.
- **Infinite gap.** If $\Delta_{\mathrm{gap}} = +\infty$, for example for a single-atom prior (the classical σ-modification), the gap condition is void. Then (iv) is m-uniform; $\bar\eta$ is then fixed by $8K^2\lambda_G^2\bar\eta^2 \leq k_e\sigma_m$ alone.
- **Switching.** $\Delta_{\mathrm{gap}} > 0$ forces the set of optimal ideal–prior assignments to be constant in t. If that set switches along a moving ideal, then $\Delta_{\mathrm{gap}} = 0$, and the second clause of (ii), the second display of (iii) and all of (iv) are vacuous.

Proof. (i) For a selection solution, at a.e. t the law uses an optimal $\tau(t)$, and $Z \leq Z_\tau := \frac{1}{2}c_\tau$ with equality at t. By Lemma B.1 it suffices to bound $\dot Z_\tau$. Apply (P) with $\vartheta' = \vartheta^0_{\tau(i)}$ and Cauchy–Schwarz in the G^{-1} inner product: $\dot Z_\tau \leq (2Z)^{1/2}\Gamma - 2\sigma_m Z$. Write $y := (2Z)^{1/2}$.

- Where $Z > 0$, $D^+ y \leq -\sigma_m y + \Gamma$.
- At $Z = 0$, $D^+ y \leq \big(\frac{1}{m}\sum \|\dot\vartheta_i\|^2_{G^{-1}}\big)^{1/2} \leq \Gamma$, by Lemma P.1 (the leak term vanishes there).
- $\Gamma \leq \lambda_G^{1/2} L_1 \|e\|$ by Lemma B.2(b).

The pure-leak case is Lemma C.3, and the projection is inactive there. In the layer, $a^\top(a^0 - a) \leq \|a\|(\|a^0\| - \|a\|) \leq 0$, and similarly for c. For s, the leak points away from the nearby face.

For Filippov velocities, the leak part averaged over the tie is

$$-\sigma_m \frac{1}{m} \sum_i \sum_k \theta_k \langle \vartheta_i - \vartheta^0_{\tau(i)},\, \vartheta_i - \vartheta^0_{\tau_k(i)} \rangle_{G^{-1}}.$$

By polarisation, and since $c_\tau = c_{\tau_k} = 2Z$ for tied optimal assignments, this equals $-2\sigma_m Z + (\sigma_m/2)\sum_k \theta_k A_{\tau\tau_k}$. Property (P) holds termwise for the convex combination. The bound is tight at the tie equilibrium of the solution-concept example ($0 = -\sigma_m/4 + \sigma_m/4$).

(ii) Relabel so that $\frac{1}{m}\sum\|\vartheta_i - \vartheta_i^*\|_{G^{-1}}^2 = \eta^2$. For any ρ:

- $|c_\rho(\vartheta) - c_\rho(\vartheta^*)| \le 2\eta R$;
- $|\sqrt{c_\rho(\vartheta)} - \sqrt{c_\rho(\vartheta^*)}| \le \eta$, by the reverse triangle inequality for the $(1/m)$-weighted G^{-1} norm.

Let ρ_0 be optimal for $(\vartheta^*(t), \vartheta^0)$. Then

$$C = c_\tau(\vartheta^*) \le c_\tau(\vartheta) + 2\eta R \le c_{\rho_0}(\vartheta) + 2\eta R \le b(t)^2 + 4\eta R,$$
$$\sqrt{C} \le \sqrt{c_\tau(\vartheta)} + \eta \le \sqrt{c_{\rho_0}(\vartheta)} + \eta \le b(t) + 2\eta.$$

So $C - b(t)^2 \le 4\eta(b(t)+\eta) < \Delta_{\mathrm{gap}} \le \Delta_{\mathrm{gap}}(t)$. No assignment cost at time t lies strictly between $b(t)^2$ and $b(t)^2 + \Delta_{\mathrm{gap}}(t)$, so $C = b(t)^2$.

(iii) Apply Theorem B with $y = \nabla g_t - \sigma_m G^{-1}(\vartheta_i - \vartheta^0_{\tau(i)})$ in (P). With $\tilde\vartheta_i := \vartheta_i - \vartheta^*_{\pi^*(i)}$ as in the proof of Theorem B, the extra term is

$$-\sigma_m \frac{1}{m}\sum \langle \tilde\vartheta_i,\, \tilde\vartheta_i + \vartheta^*_{\pi^*(i)} - \vartheta^0_{\tau(i)}\rangle_{G^{-1}} \le -\sigma_m\eta^2 + \sigma_m\eta C^{1/2}.$$

Under the gap condition, (ii) gives $C^{1/2} = b(t) \le b$, and $\sigma_m\eta b \le (\sigma_m/2)(\eta^2 + b^2)$.

(iv) Apply Young's inequality three times:

- $\|e\|\varepsilon_{\max} \le (k_e/4)\|e\|^2 + \varepsilon_{\max}^2/k_e$;
- $\|e\|K\lambda_G\eta^2 \le (k_e/4)\|e\|^2 + (\sigma_m/8)\eta^2$ on $\{\eta \le \bar\eta\}$;
- $V_*\eta \le (\sigma_m/8)\eta^2 + 2V_*^2/\sigma_m$.

This gives $D^+V_{\mathrm{OT}} \le -\alpha V_{\mathrm{OT}} + \beta$. Each set $\{V_{\mathrm{OT}} \le c\}$ with $\beta/\alpha < c < \bar\eta^2/2$ lies inside $\{\eta \le \bar\eta\}$ and has a negative boundary derivative. Intersect over $c > \beta/\alpha$ for the endpoint, and apply the comparison lemma.

(v) Steps (a)–(c) of Theorem B hold for any fixed π, so use the classical completion of squares for V_π.

- *Equality case.* If $\Delta_{\mathrm{gap}} > 0$, the costs minus $b(t)^2$ take values in $\{0\} \cup [\Delta_{\mathrm{gap}}, \infty)$ and are continuous in t, so the optimal set is constant.
- *Strict case.* If the optimal ideal–prior assignment changes along the ideal, the inequality can be strict. For example, $a_1^*(t) = t$ and $a_2^*(t) = 1 - t$ on $[0,1]$, with prior $a \in \{0,1\}$, give $b = 0.5$ against 1.0.
- *Not always strict.* For $a^*(t) = (u(t),\, 1-u(t))$ with $u(t) = -0.025 + 0.575\sin t$, the optimal assignment alternates between the identity and the swap, yet $\min_\pi \sup_t b_{\mathrm{lab}}(\pi)(t) = \sup_t b_{\mathrm{lab}}(\mathrm{id})(t) = 0.6 = b$ (computed numerically).

The negative statement is Proposition C.5. □

Lemma C.3 (no switching along the pure leak; exact decay for any $G \succ 0$)**.** *Let τ be optimal for $(\mu_{\mathrm{init}}, \mu^0)$ and $\dot\vartheta_i = -\sigma_m(\vartheta_i - \vartheta^0_{\tau(i)})$. Then τ stays optimal and $W(\mu_t, \mu^0) = e^{-\sigma_m t}W(\mu_{\mathrm{init}}, \mu^0)$.*

Proof. Put $d_i := \vartheta_i(0) - \vartheta^0_{\tau(i)}$, so $\vartheta_i(t) - \vartheta^0_{\tau(i)} = \lambda d_i$ with $\lambda = e^{-\sigma_m t}$. For any ρ, with $q_i := \vartheta^0_{\tau(i)} - \vartheta^0_{\rho(i)}$,

$$c_\rho(t) - c_\tau(t) = \frac{2\lambda}{m}\sum\langle d_i,\, q_i\rangle_{G^{-1}} + \frac{1}{m}\sum\|q_i\|_{G^{-1}}^2.$$

This is affine in λ. It is non-negative at $\lambda = 1$ by optimality and at $\lambda = 0$ trivially, hence on all of $[0,1]$.

At an exact tie at $t = 0$, the chosen assignment becomes strictly optimal immediately against every ρ with $A_\rho := \frac{1}{m}\sum\|q_i\|^2 > 0$. Assignments with $A_\rho = 0$ induce the same velocity. So no

selection solution of the pure leak switches. Filippov sliding on the tie is a different solution, excluded by the solution concept. □

Checked by: `test_theorem_C_i_exact_dissipation_pure_leak` in `tests/test_transport.py`: zero switches, and exact decay to integrator accuracy.

Proposition C.5 (the labelled leak reshuffles)**.** *There are configurations with* $b = b_{\mathrm{lab}}(\mathrm{id}) = 0$ *and* $e = 0$ *at which the labelled leak gives* $D^+V_{\mathrm{OT}} > 0$.

Example. Take $n = 1$, $G = I$ and $a_{\max} - \varepsilon_K \geq 1$. Let ideal = prior = $\{a = 0,\ a = 1\}$, with the other coordinates equal and in the inner set, and let the network weights be $(1 - \delta, \delta)$ with $0 < \delta < \frac{1}{2}$.

- The optimal assignment to the ideal is the swap.
- The labelled leak moves each particle toward the *other* ideal atom.
- So $D^+V_{\mathrm{OT}} = \sigma_m(\delta - \delta^2) > 0$. □

From a relabelled start, the labelled leak moves every kernel in a non-trivial cycle of the relabelling to its prior position. In the reshuffling experiment (`results/ot_sigma_mod/transient.md` in the accompanying repository):

- the steady states coincide;
- the labelled leak's particle path is 5.8× longer (2.26 against 0.39);
- the early RMS$\|e\|$ is only about 4% larger (0.0771 against 0.0743 on $[0, 5]$).

Theorem F does not price this motion. With a leak, Theorem F bounds transport from above by tracking error plus a leak term, and the leak can pay for all of it. With $\delta = 0$ above, $e \equiv 0$ while every particle crosses the domain.
Checked by: the reshuffling transient, `experiments/exp_sigma_mod_transient.py`. The value $D^+V_{\mathrm{OT}} = \sigma_m(\delta - \delta^2)$ is not checked in the repository.

Remark C.6 (the obstruction is known)**.**

- **Non-convexity.** For some ν, for example a two-atom ν in dimension ≥ 2, $\frac{1}{2}W^2(\cdot, \nu)$ is not λ-convex along geodesics for any λ (Ambrosio–Gigli–Savaré [2, Example 9.1.5]; Santambrogio [53, §4.5]). It is 1-semiconcave in every dimension [2, Thm. 7.3.2]. For $\nu = \delta_{\vartheta^0}$ it is a potential energy, and 1-convex.
- **Generalised geodesics.** $\frac{1}{2}W^2(\cdot, \nu)$ is 1-convex along generalised geodesics with base ν ([2, Lemma 9.2.1]; Carlen–Craig [5]).
- **Label-free leaks.**
  - A label-free leak that is λ-convex ($\lambda > 0$) on the lift $\mathbb{R}^{mp}$ has a unique, permutation-invariant minimiser, at which all particles coincide. So it cannot vanish at a μ^0 with distinct atoms.
  - 0-convex leaks that vanish at μ^0 exist, for example toward $\operatorname{conv}\operatorname{supp}\mu^0$. They do not single out μ^0.

So the trade-off reflects this geometry. On one side is a global inequality for a labelled V_π, which is large at relabelled starts (hence the reshuffling of Proposition C.5). On the other is a local, gap-protected inequality for the label-free V_{OT}.

13 Theorem D: the mean-field closed loop

The state is $(x, \mu) \in \mathbb{R}^n \times \mathcal{P}(\Theta_K)$, with

$$\begin{gathered}\dot{x} = f(x,t) + \dot{x}_m - k_e(x - x_m) - \int \Phi(x;\vartheta)\,\mathrm{d}\mu_t, \qquad \partial_t \mu_t + \nabla \cdot (\mu_t v_t) = 0, \\ v_t = \mathrm{Proj}_K\big(\vartheta,\, G\nabla_\vartheta\big[(x - x_m)^\top \Phi(x;\vartheta)\big]\big).\end{gathered} \tag{13.1}$$

Solutions are Lagrangian: $\mu_t = (\Xi_t)_\# \mu_0$ for the flow Ξ_t of v_t. This is law (L). The OT leak is multivalued at ties and is excluded here.

Theorem D. *Assume* (S0)–(S2).

(a) Local well-posedness. *For every $(x_0, \mu_0) \in \mathbb{R}^n \times \mathcal{P}(\Theta_K)$ there is a unique Lagrangian solution. It continues for as long as x stays bounded.*

(b) Adaptation-free tube (m-uniform, μ^*-free). *For every $\mu \in \mathcal{P}(\Theta_K)$, $\|\int \Phi(x;\cdot)\,\mathrm{d}\mu\| \le a_{\max}$. For $F \ge 0$ let*

$$r(F) := \|e(0)\| + \frac{F + a_{\max}}{k_e}, \qquad S(F) := \sup\big\{\|f(x,t)\| : t \ge 0,\ \|x - x_m(t)\| \le r(F)\big\}.$$

Suppose some $F \ge 0$ satisfies $S(F) \le F$. Then every solution, for every m and every μ_0:

- *exists for all $t \ge 0$;*
- *satisfies $\|x(t) - x_m(t)\| \le r(F)$;*
- *satisfies $\|e(t)\| \le e^{-k_e t}\|e(0)\| + (1 - e^{-k_e t})(F + a_{\max})/k_e$.*

Examples:

- *For bounded f, $F = \sup\|f\|$ works for every $e(0)$.*
- *If $\|f(x,t)\| \le \alpha + \beta\|x\|$ with $\beta < k_e$, then*

$$F = \frac{\alpha + \beta\big(\sup\|x_m\| + \|e(0)\| + a_{\max}/k_e\big)}{1 - \beta/k_e}$$

  works.
- *The condition can fail for superlinear f. For the Van der Pol plant of the numerics, $S(F) - F \ge 51.6$ for every F.*

(c) Dobrushin estimate. *Let two solutions of the same loop (same f, x_m, k_e, G, Θ_K and ε_K) remain in a compact X on $[0,T]$. Then*

$$\begin{gathered}\|x_t - x'_t\| + W(\mu_t, \mu'_t) \le e^{\Lambda t}\big(\|x_0 - x'_0\| + W(\mu_0, \mu'_0)\big), \\ \Lambda := \max(L_\vartheta + L_2,\, L_x + M_x),\end{gathered} \tag{13.2}$$

and Λ is independent of m. The constants are:

- *L_ϑ: a Lipschitz constant of $\vartheta \mapsto v$ in the G^{-1} norm, uniform over $x \in X$ and $t \le T$. It is proportional to $\bar{E} := \sup_{X\times[0,T]}\|x - x_m(t)\|$, and it is $O(\bar{E}/\varepsilon_K)$ because of the ramp ($h' \le 2/\varepsilon_K$).*
- *L_x: a Lipschitz constant, Euclidean to G^{-1}, of $x \mapsto v(\vartheta; x, x - x_m(t))$, uniform over $\vartheta \in \Theta_K$, $t \le T$ and $x \in X$.*
- *M_x: $M_x := \sup_{t\le T} \mathrm{Lip}_X\, f(\cdot,t) + k_e + a_{\max} J_c$. Here $a_{\max} J_c$ bounds $\|\nabla_x \int \Phi(x;\cdot)\,\mathrm{d}\mu\|$ over $x \in \mathbb{R}^n$ and $\mu \in \mathcal{P}(\Theta_K)$.*

- *L_2: $L_2 := \lambda_G^{1/2} L_1$, with L_1 from Lemma B.2, is a W-Lipschitz constant of $\mu \mapsto \int \Phi(x;\cdot)\,\mathrm{d}\mu$. For any coupling, $\|\int \Phi(x;\cdot)\,\mathrm{d}(\mu-\mu')\| \le L_2\big(\int \|\vartheta-\vartheta'\|_{G^{-1}}^2\,\mathrm{d}\gamma\big)^{1/2}$.*

(d) Mean-field limit. *Let (x,μ) be the solution from (x_0,μ_0), and let T be smaller than its existence time. Put $X_1 := \{y : \mathrm{dist}(y, x([0,T])) \le 1\}$, and let Λ_1 be the Λ of (c) computed on $X_1 \times [0,T]$. Suppose $x_0^m \to x_0$ and $\mu_0^m \to \mu_0$ in W, and set $\delta_m := \|x_0^m - x_0\| + W(\mu_0^m, \mu_0)$. Then, for all large m:*

- *the m-particle loop exists on $[0,T]$ and stays in X_1;*
- *$\sup_{t\le T}\big(\|x_t^m - x_t\| + W(\mu_t^m,\mu_t)\big) \le e^{\Lambda_1 T}\delta_m \to 0$.*

Under the hypothesis of (b), for example for bounded f, the limit is global and T is arbitrary.

(e) Passage of (B) to the limit. *Work in the setting of (d), with $X := X_1$. Let $\mu_t^{*m} = \frac{1}{m}\sum_j \delta_{\vartheta_j^{*m}(t)}$ satisfy* (S4), *and let $\varepsilon_{\max}^m \ge \sup_{X\times[0,T]}\|f - \int \Phi\,\mathrm{d}\mu_t^{*m}\|$. Assume $\mu_t^{*m} \to \mu_t^*$ uniformly in W on $[0,T]$, $\sup_m \varepsilon_{\max}^m \le \varepsilon_{\max}$ and $\sup_m V_*^m \le V_*$. Only tails matter, so $\sup_m$ may be replaced by $\limsup_m$. Then:*

- *μ_t^* is V_*-Lipschitz in W, with values in $\mathcal{P}(\Theta_K^\varepsilon)$;*
- *the limit satisfies* (B) *in integrated form, and in Dini form for every $t \in [0,T)$, with constants $(k_e, \varepsilon_{\max}, K, L_0, L_1, \lambda_G, V_*)$.*

V_* depends on the approximating ideals; the intrinsic version is Remark B.7.

Status. Proved; consistency in m checked numerically (the constant is not checked). Not checked in the repository: (b), the constant of (c), and the one-sided constant of Remark D.1. Novelty (literature audit): technique anticipated [17, 47, 48].

Proof. *(a)* Write the loop as an ODE for $(x,\Xi) \in \mathbb{R}^n \times C(\Theta_K;\mathbb{R}^p)$, with the sup norm (and G^{-1} on $\mathbb{R}^p$):

$$\dot x = f(x,t) + \dot x_m - k_e(x - x_m) - \int \Phi(x;\Xi(\vartheta))\,\mathrm{d}\mu_0(\vartheta), \qquad \partial_t \Xi(\vartheta) = v(\Xi(\vartheta);x,t), \qquad \Xi_0 = \mathrm{id},$$

so that $\mu_t = \Xi_{t\#}\mu_0$. Extend v Lipschitz to a neighbourhood of Θ_K inside $\{s > s_{\min}/2\}$ (McShane). On $\{\|x\| \le R\}$, with the constants of (c), the right-hand side is Lipschitz:

- $\|v(\Xi(\cdot);x,t) - v(\Xi'(\cdot);x',t)\|_\infty \le L_\vartheta\|\Xi - \Xi'\|_\infty + L_x\|x - x'\|$;
- the x-field changes by at most $M_x\|x-x'\| + L_2\|\Xi - \Xi'\|_\infty$, because $W(\Xi_\#\mu_0, \Xi'_\#\mu_0) \le \|\Xi - \Xi'\|_{L^2(\mu_0;G^{-1})}$.

Picard–Lindelöf gives a unique local solution. Each characteristic stays in Θ_K (Nagumo, as in A(ii)), so the solution continues while x is bounded. Golse [24, Thms. 1.3.2 and 1.4.3], and [17, 47, 48], are structural analogues (pairwise C^1 kernels, W_1), not the proof.

(b) While $\|e\| \le r(F)$, $\dot e = -k_e e + (f - \hat f)$ with $\|f - \hat f\| \le F + a_{\max}$. Variation of constants gives

$$\|e(t)\| \le e^{-k_e t}\|e(0)\| + (1 - e^{-k_e t})(F + a_{\max})/k_e < r(F)$$

for every finite t, since $a_{\max} > 0$. So x never reaches the boundary of the tube (first exit), stays bounded, and (a) continues the solution.

(c) Let γ_0 be an optimal plan for (μ_0,μ_0'), let Ξ_t and Ξ_t' be the two flows, and set $D(t)^2 := \int\|\Xi_t(\vartheta) - \Xi_t'(\vartheta')\|_{G^{-1}}^2\,\mathrm{d}\gamma_0$. Then $(\Xi_t,\Xi_t')_\#\gamma_0$ couples μ_t and μ_t', so $W(\mu_t,\mu_t') \le D(t)$. Split

$$v_t(\Xi) - v_t'(\Xi') = \big[v_t(\Xi) - v_t(\Xi')\big] + \big[v_t(\Xi') - v_t'(\Xi')\big]$$

and apply Cauchy–Schwarz:

$$\frac{\mathrm{d}}{\mathrm{d}t}D^2 \le 2L_\vartheta D^2 + 2L_x\|x-x'\|D, \qquad D^+\|x-x'\| \le M_x\|x-x'\| + L_2 D.$$

Apply Grönwall to $D_\delta + \|x-x'\|$, with $D_\delta := (D^2+\delta^2)^{1/2}$, and let $\delta \downarrow 0$.

(d) Let $\tau_m \le T$ be the first time x^m leaves X_1, with $\tau_m = T$ if it never leaves. On $[0,\tau_m]$ both solutions lie in X_1, so (c) gives $\|x_t^m - x_t\| \le e^{\Lambda_1 T}\delta_m < 1$ once m is large. Then $x^m(\tau_m)$ is interior to X_1, so $\tau_m = T$. By (a), the m-solution continues while it is bounded.

(e) Theorem B for each m gives $V^m(t_2) - V^m(t_1) \le \int R^m \le \int \bar{R}^m$. Here $\bar{R}^m$ is R^m with $\varepsilon_{\max}^m$ and $V_*^m(t)$ replaced by $\varepsilon_{\max}$ and V_*. By (d) and the triangle inequality for W, $e^m \to e$ and $W(\mu_t^m, \mu_t^{*m}) \to W(\mu_t, \mu_t^*)$ uniformly. So $V^m \to V$ and $\bar{R}^m \to \bar{R}$ uniformly, and $V(t_2) - V(t_1) \le \int \bar{R}$. $\bar{R}$ is continuous, which gives the Dini form. The identity coupling gives $W(\mu_t^{*m}, \mu_s^{*m}) \le V_*|t-s|$. This passes to the uniform limit, and Θ_K^ε is closed. □

Remark D.1 (a one-sided constant free of ε_K)**.** The coupling step uses v only through $\langle \Xi - \Xi',\, G^{-1}(v(\Xi) - v(\Xi'))\rangle$. So the one-sided constant

$$L_\vartheta^+ := \sup \frac{\langle \vartheta - \vartheta',\, G^{-1}(v(\vartheta) - v(\vartheta'))\rangle}{\|\vartheta - \vartheta'\|_{G^{-1}}^2}$$

may replace L_ϑ. Let $N(\theta, y) := h(\|\theta\|)(\hat{\theta}^\top y)_+\hat{\theta}$ be the norm-ball correction, with $r = \|\theta\|$, $c = \hat{\theta}^\top\hat{\theta}'$ and $\alpha = (\hat{\theta}^\top y)_+$. Then

$$\langle \theta - \theta',\, N(\theta,y) - N(\theta',y)\rangle \ge -\frac{\|y\|\|\theta-\theta'\|^2}{2(\theta_{\max} - \varepsilon_K)}.$$

This follows from three facts:

- h is monotone;
- $|\alpha - \alpha'| \le \|y\|\sqrt{2(1-c)}$;
- $\|\theta - \theta'\|^2 \ge 2\sqrt{rr'}\,|r - r'|\sqrt{2(1-c)}$.

The box correction is monotone. Hence L_ϑ^+ stays bounded as $\varepsilon_K \to 0$, and so does $\Lambda := \max(L_\vartheta^+ + L_2,\, L_x + M_x)$. On the experiment's box, a numerical measurement gives $L_\vartheta^+ \approx 88$–98 for ε_K from 0.2 to 0.005. Over the same range the two-sided constant grows like $30/\varepsilon_K$. This measurement is not repeated in the repository's tests.

Numerics (`results/mean_field_m`). Halving $W(\mu_0^m, \mu_0^{2m})$ halves $W(\mu_T^m, \mu_T^{2m})$, with a nearly constant factor (1.64–1.69 over the four refinements). $\sup_t\|e^m - e^{2m}\|$ falls from 4.6×10^{-3} (16 against 32 particles) to below 10^{-4} (128 against 256).

This is a consistency check only.

- On the experiment's box the guaranteed factor $e^{\Lambda T}$ is at least e^{496}, from $L_2 = 33.08$ alone over $T = 15$.
- The condition of (b) fails for this Van der Pol plant, so the run is covered by the localised (d) through its observed boundedness, not by (b).

Checked by: `experiments/exp_mean_field_m.py` (consistency in m; not the constant $e^{\Lambda T}$).

14 Theorem E: label-free local convergence under tangential excitation

Assumptions.

- (S0) holds, in particular $k_e > 0$.
- **Uniform tube.** There is $\rho > 0$ such that $\varepsilon(x,t) = 0$ whenever $\|x - x_m(t)\| < \rho$, for all $t \geq 0$.
- **Interior ideal.** μ^* is constant, with $\vartheta_j^* \in \operatorname{int}\Theta_K^\varepsilon$. Put $\delta := \min_j \operatorname{dist}(\vartheta_j^*, \partial\Theta_K^\varepsilon) > 0$.
- $x_m \in C^1$, and x_m and $\dot{x}_m$ are bounded.
- Define $\Psi(t) := \big[\nabla_\vartheta \Phi(x_m(t);\vartheta_1^*), \dots, \nabla_\vartheta \Phi(x_m(t);\vartheta_m^*)\big] \in \mathbb{R}^{n\times mp}$.
- **(PE)** There are $T, \alpha > 0$ with $\int_t^{t+T} \Psi^\top \Psi \,\mathrm{d}r \succeq \alpha I_{mp}$ for all $t \geq 0$.

(PE) forces $a_j^* \neq 0$ and pairwise distinct pairs (c_j^*, s_j^*).

- If $a_j^* = 0$, the (c,s)-columns of block j vanish.
- If $(c_j^*, s_j^*) = (c_k^*, s_k^*)$, the kernel of $\int \Psi^\top \Psi \,\mathrm{d}r$ contains the vector with v in the a-slot of block j, $-v$ in the a-slot of block k, and zeros elsewhere.

In particular the ϑ_j^* are pairwise distinct, and

$$d^2 := \min_{\pi \neq \pi'} \frac{1}{m} \sum_i \|\vartheta_{\pi(i)}^* - \vartheta_{\pi'(i)}^*\|_{G^{-1}}^2 = \frac{2}{m} \cdot \min_{i \neq j} \|\vartheta_i^* - \vartheta_j^*\|_{G^{-1}}^2 > 0, \tag{14.1}$$

with $d^2 := +\infty$ for $m = 1$. The closed form holds because a non-identity permutation moves at least two indices, and the transposition of the G^{-1}-closest pair attains the bound.

Theorem E. *There are $r_0, k, \lambda > 0$ such that the following holds.*

- *k and λ come from the converse Lyapunov function of the linearisation (proof steps 3–4).*
- *r_0 depends in addition on $r := \min(\rho, \delta)$, and on the Lipschitz constant of the Jacobian on the ball D of step 2.*

Put

$$c_0 := \frac{r_0^2}{2(1 + m\lambda_G)} \qquad \text{and} \qquad C := \sqrt{2} \cdot k \cdot \max\big\{1, (m\lambda_G)^{1/2}\big\} \cdot \max\big\{1, (m\lambda_{\min}(G))^{-1/2}\big\}.$$

If $V_{\mathrm{OT}}(t_0) \leq c_0$, then:

- *the optimal assignment π at t_0 is unique, and it is the unique optimal assignment for all large t;*
- *$\|e(t)\| + W(\mu_t, \mu^*) \leq C e^{-\lambda(t-t_0)}\big(\|e(t_0)\| + W(\mu_{t_0}, \mu^*)\big)$ for $t \geq t_0$;*
- *$\vartheta_i(t) \to \vartheta_{\pi(i)}^*$. The labelled error*

$$E_{\mathrm{lab}}(t) := \Big(\frac{1}{m}\sum_i \|\vartheta_i(t) - \vartheta_i^*\|_{G^{-1}}^2\Big)^{1/2} \quad \text{tends to} \quad \Big(\frac{1}{m}\sum_i \|\vartheta_{\pi(i)}^* - \vartheta_i^*\|_{G^{-1}}^2\Big)^{1/2},$$

which is non-zero for $\pi \neq \mathrm{id}$.

k and λ depend on $\Psi(\cdot)$ itself, through the converse Lyapunov function of its linearisation, and on m, G and k_e. r_0 and c_0 depend in addition on $a_{\max}$, $s_{\min}$, ρ and δ. The cited theorems are of existence type, so it is not proved here that k and λ can be chosen uniformly over the class of Ψ with given $(T, \alpha, \sup\|\Psi\|, \sup\|\dot{\Psi}\|)$.

Status. Proved, conditional on Morgan–Narendra [39, Thm. 1] as stated in its abstract (its hypotheses were not checked against the full text); linearisation checked; convergence observed. Novelty (literature audit): partially anticipated [10], [22].

Proof.

1. *Symmetry.* Relabelling the network, $(e, \vartheta) \mapsto (e, \vartheta \circ \sigma)$, maps solutions to solutions. It fixes e, is an isometry of the matched metric, and conjugates the dynamics near $(0, \vartheta^* \circ \pi)$ to those near $(0, \vartheta^*)$. So it suffices to treat $\pi = \mathrm{id}$.

2. *Domain and linearisation.* Let $r := \min(\rho, \delta)$ and $D := \{z = (e, \tilde{\vartheta}) \in \mathbb{R}^{n+mp} : \|z\|_2 < r\}$.

   - For $z \in D$ we have $\|e\| < \rho$, so $f = \hat{f}^*$.
   - Also $\|\tilde{\vartheta}_i\| < \delta$ for every i, so $\vartheta_i \in \operatorname{int} \Theta_K^\varepsilon$ and $\mathrm{Proj}_K = \mathrm{id}$.

   On $[0, \infty) \times D$ the closed loop is

   $$F(t, z) = \Big(-k_e e + \hat{f}^*(x) - \hat{f}(x),\ \big(G \nabla_\vartheta \Phi(x; \vartheta_i)^\top e\big)_i\Big), \qquad x = x_m(t) + e.$$

   - It is C^1 in (t, z), since $x_m \in C^1$, and C^∞ in z.
   - Its derivatives through order two are bounded uniformly in t, because the Gaussian's derivatives through order three are bounded uniformly in x for $s \geq s_{\min}$.
   - Its Jacobian at 0 is
     $$A(t) = \begin{pmatrix} -k_e I & -\frac{1}{m}\Psi \\ (I_m \otimes G)\Psi^\top & 0 \end{pmatrix},$$
     and the remainder is $O(\|z\|^2)$.
   - The Jacobian and the identity $QA + A^\top Q = \mathrm{diag}(-k_e I, 0)$, with $Q = \mathrm{diag}\big(I/2,\ (I \otimes G^{-1})/(2m)\big)$, are checked numerically by `test_theorem_E_linearisation_and_lyapunov_identity`.

3. *LTV stability.* Put $w := (e, \zeta)$ with $\zeta := m^{-1/2}(I_m \otimes G^{-1/2})\tilde{\vartheta}$, and $\Psi_G := m^{-1/2}\Psi(I \otimes G^{1/2})$.

   - The system becomes $\dot{e} = -k_e e - \Psi_G \zeta$ and $\dot{\zeta} = \Psi_G^\top e$, that is, $\dot{w} = [A_0 + B(t)]w$ with $A_0 = \mathrm{diag}(-k_e I, 0)$ and B skew-symmetric, bounded, with bounded derivative.
   - The excitation transfers: $\int \Psi_G^\top \Psi_G \, \mathrm{d}r \succeq (\alpha \lambda_{\min}(G)/m) I$.
   - Morgan–Narendra [39, Thm. 1] characterises uniform asymptotic stability of
     $$\begin{pmatrix} A & -B(t) \\ B(t)^\top & 0 \end{pmatrix}$$
     with $A + A^\top \prec 0$ by richness of B. Apply it with $A = -k_e I$ and $B = \Psi_G$ (bounded, with bounded derivative): the origin of the w-system is uniformly asymptotically stable.
   - (PE) supplies the richness, since $\int_t^{t+T} \|\Psi_G w\| \, \mathrm{d}r \geq \alpha \lambda_{\min}(G)/(m \cdot \sup\|\Psi_G\|)$ for every unit w.
   - For linear systems UAS is UES (Khalil [31, Thm. 4.11]). UES transfers to $(e, \tilde{\vartheta})$ through the constant similarity $\mathrm{diag}\big(I, m^{1/2}(I \otimes G^{1/2})\big)$.
   - The bound on $\dot{\Psi}$ cannot be dropped. $\Psi = \sin t^2$ is bounded and PE, but for $\dot{e} = -e - \Psi\zeta$, $\dot{\zeta} = \Psi e$, from $w(t_0) = (e, \zeta) = (0, 1)$, a numerical computation gives the ratio $\|w(t_0 + 200)\|/\|w(t_0)\|$ as 0.9715, 0.9942 and 0.9980 for $t_0 = 5$, 20 and 50. So that system is not UES.
   - We could not check the exact richness definition and regularity class of [39] against its full text. Its abstract's characterisation matches this system.

4. *Nonlinear system.* Apply Khalil [31, Thm. 4.13] in $z = (e, \tilde{\vartheta})$ on $[0, \infty) \times D$. The Jacobian is bounded and Lipschitz on D uniformly in t, and the linearisation is UES. This gives $r_0 \le r$, k and λ, with $\|z(t)\| \le k\|z(t_0)\|e^{-\lambda(t-t_0)}$ whenever $\|z(t_0)\| < r_0$. These trajectories stay in D, where the closed loop equals F.

5. *Transfer to* $(\|e\|, W)$. If $V_\pi(t_0) \le c_0$, then

$$\|z(t_0)\|^2 \le 2\max(1, m\lambda_G)V_\pi(t_0) \le 2\max(1, m\lambda_G)c_0 < 2(1 + m\lambda_G)c_0 = r_0^2.$$

   The bound $W \le (m\lambda_{\min}(G))^{-1/2}\|\tilde{\vartheta}\|$ and the optimality of π at t_0 give C.

   Uniqueness. Suppose $V_\pi, V_{\pi'} \le c_0$ with $\pi \ne \pi'$. Then the state lies in Khalil balls around both $(0, \vartheta^* \circ \pi)$ and $(0, \vartheta^* \circ \pi')$. The solution would converge to two distinct equilibria, which is impossible.

   Limits. Hence $z(t) \to 0$, that is, $\vartheta_i(t) \to \vartheta^*_{\pi(i)}$. For $\pi' \ne \pi$,

$$V_{\pi'}(t) \to \frac{1}{2m}\sum_i \|\vartheta^*_{\pi(i)} - \vartheta^*_{\pi'(i)}\|^2_{G^{-1}} \ge \frac{d^2}{2} > 0,$$

   while $V_\pi(t) \to 0$. So π is the unique optimal assignment for all large t, and E_{lab} converges as stated. □

Checked by: `tests/test_transport.py` and `experiments/exp_pe_convergence.py`.

- Linearisation: the Jacobian of the implemented loop equals $A(t)$, and $QA + A^\top Q = \operatorname{diag}(-k_e I, 0)$ (`test_theorem_E_linearisation_and_lyapunov_identity`).
- Convergence: the matched error decays and the labelled error does not (qualitative); sliding-window excitation and fitted rates (`test_theorem_E_matched_error_converges_labelled_does_not`, `exp_pe_convergence.py`).
- Not checked in the repository: the rate of Theorem E, Corollary E.2 and Remark E.3.

Remark E.1 (the rate)**.**

- **What converges.** "The parameters converge" is false for the labelled error and true for the matched one.
- **The rate is not set by the PE level.** $\lambda > 0$ exists for each admissible Ψ, but it depends on $\Psi(\cdot)$, not only on its PE level. No positive function of $\alpha\lambda_{\min}(G)/m$, k_e, T and $\sup\|\Psi_G\|$ alone bounds it below.
  - For $m = n = 1$, $\vartheta^* = (1.5, 0.3, 0.8)$, $G = I$, $k_e = 3$ and $x_m = 1.5\sin\omega t$ ($T = 2\pi$), these four quantities do not depend on ω. The exact Floquet rate, computed numerically, falls from 4.7×10^{-2} ($\omega = 1$) to 2.9×10^{-5} ($\omega = 64$), like ω^{-2}.
  - A lower bound must also involve $\sup\|\dot{\Psi}_G\|$ (step 3 of the proof), and none is proved here.
  - Exact Floquet rates, computed numerically, are also not monotone in G or in k_e, and they decrease for large G. When k_e is large relative to the reference bandwidth, they approach $\lambda_{\min}\langle\Psi_G^\top\Psi_G\rangle/k_e$.
- **The numbers** (`results/pe_convergence`; $m = 3$, $n = 2$, six runs):

quantity	range
min over sliding 60 s windows of $\lambda_{\min}\big(\int_t^{t+60}\Psi^\top\Psi\,\mathrm{d}r\big)$	5.9×10^{-4} to 2.7×10^{-2}
fitted decay rate of the matched error, second half of the run	2.3×10^{-3} to $4.3 \times 10^{-3}\,\mathrm{s}^{-1}$
rank correlation between the two	0.60

The convergence test is therefore qualitative. The machine-checked part of the proof is the linearisation.

Corollary E.2 (uniform-in-time exponential merging under PE)**.** *Take two solutions of the same closed loop with $V_{\mathrm{OT}}(0) \le c_0$ and $V'_{\mathrm{OT}}(0) \le c_0$. Then*

$$\|x_t - x'_t\| + W(\mu_t, \mu'_t) \le Ce^{-\lambda t} R_0, \qquad R_0 := \|e(0)\| + \|e'(0)\| + W(\mu_0, \mu^*) + W(\mu'_0, \mu^*). \tag{14.2}$$

Let $\delta_0 := \|x_0 - x'_0\| + W(\mu_0, \mu'_0)$. Combine this with the coupling estimate of Theorem D(c) on $[0, T_0]$, with $T_0 := (\Lambda + \lambda)^{-1} \log(CR_0/\delta_0)$.

- *Both solutions stay in D (proof step 4), so $\|x - x_m(t)\| < r \le \rho$.*
- *The segment between x_t and x'_t lies in $B(x_m(t), \rho)$, where $f = \hat{f}^*$, and $\mathrm{Proj}_K = \mathrm{id}$.*
- *Hence Theorem D(c) holds with $M_x := k_e + 2a_{\max} J_c$, with L_x and L_ϑ computed for $\bar{E} = r$ and an inactive projection, and with $L_2 = \lambda_G^{1/2} L_1$. This Λ depends neither on T_0 nor on f outside the tube.*

This gives uniform-in-time Hölder continuity in the basin:

$$\sup_{t \ge 0} \big(\|x_t - x'_t\| + W(\mu_t, \mu'_t) \big) \le (CR_0)^{\Lambda/(\Lambda+\lambda)} \, \delta_0^{\lambda/(\Lambda+\lambda)}. \tag{14.3}$$

Status. Proved, conditional on Morgan–Narendra [39, Thm. 1] as stated in its abstract (its hypotheses were not checked against the full text), through Theorem E. Not checked in the repository.

Proof. $\|x_t - x'_t\| = \|e - e'\| \le \|e\| + \|e'\|$, and $W(\mu_t, \mu'_t) \le W(\mu_t, \mu^*) + W(\mu'_t, \mu^*)$. Bound each bracket by Theorem E. On $[0, T_0]$ use $e^{\Lambda T_0} \delta_0$; on $[T_0, \infty)$ use $CR_0 e^{-\lambda T_0}$. Note that $\delta_0 \le R_0 \le CR_0$. □

Remark E.3 (the collapse point)**.** Take the G^{-1}-closest pair (j, k). Set $e = 0$ and $\vartheta_j = \vartheta_k = (\vartheta_j^* + \vartheta_k^*)/2$, with all other particles ideal.

- At this point $V_{\mathrm{OT}} = d^2/8$, and there are two optimal assignments.
- By exchange symmetry, $\vartheta_j \equiv \vartheta_k$ for all t, so $W(\mu_t, \mu^*) \ge d/2$ forever.

The uniqueness argument of step 5 shows $c_0 < d^2/8$ automatically, and $d^2/8$ is sharp along this direction.

15 Theorem F: the transport cost of adaptation

Theorem F. *Assume* (S0)–(S3)*, and take $t_0 \le t_1$ in the maximal interval of existence (all $t \ge 0$ under the hypothesis of Theorem D(b)). μ^*,* (S4) *and* (S5) *are not needed. Along* (L)*:*

(i) $\big(\frac{1}{m} \sum_i \|\dot{\vartheta}_i\|_{G^{-1}}^2\big)^{1/2} \le \lambda_G^{1/2} L_1 \|e(t)\|$;

(ii) $W(\mu_{t_1}, \mu_{t_0}) \le \lambda_G^{1/2} L_1 \int_{t_0}^{t_1} \|e\| \, \mathrm{d}r$. *Hence the W-length of $r \mapsto \mu_r$ on $[t_0, t_1]$ is at most $\lambda_G^{1/2} L_1 \int_{t_0}^{t_1} \|e\| \, \mathrm{d}r$;*

(iii) $W(\mu_{T_2}, \mu_{T_1}) \le \lambda_G^{1/2} L_1 (T_2 - T_1) \cdot \mathrm{RMS}_{[T_1, T_2]} \|e\|$.

Along (L_σ) *with the OT leak, for a.e. t:*

(i′) $\big(\frac{1}{m} \sum_i \|\dot{\vartheta}_i\|_{G^{-1}}^2\big)^{1/2} \le \lambda_G^{1/2} L_1 \|e\| + \sigma_m W(\mu_t, \mu^0)$;

(ii′) $W(\mu_{t_1}, \mu_{t_0}) \le \int_{t_0}^{t_1} \big(\lambda_G^{1/2} L_1 \|e\| + \sigma_m W(\mu_r, \mu^0) \big) \, \mathrm{d}r$;

(iii′) $W(\mu_{T_2}, \mu_{T_1}) \le \lambda_G^{1/2} L_1 (T_2 - T_1) \cdot \mathrm{RMS}_{[T_1, T_2]} \|e\| + \sigma_m \int_{T_1}^{T_2} W(\mu_r, \mu^0) \, \mathrm{d}r$.

For the labelled leak, replace $W(\mu_r, \mu^0)$ *by* $d_{\mathrm{lab}}(r) := \big(\frac{1}{m}\sum_i \|\vartheta_i(r) - \vartheta_i^0\|_{G^{-1}}^2\big)^{1/2} \geq W(\mu_r, \mu^0)$. *The term* $W(\mu_r, \mu^0)$ *does not suffice there.*

All bounds also hold with $L_G := \big(g_a + a_{\max}^2(g_c J_c^2 + g_s J_s^2)\big)^{1/2} \leq \lambda_G^{1/2} L_1$ *in place of* $\lambda_G^{1/2} L_1$. *Since* $\max_i \|\dot\vartheta_i\|_{G^{-1}} \leq L_G\|e\|$ *along* (L), *also* $W_\infty(\mu_{t_1}, \mu_{t_0}) \leq L_G \int_{t_0}^{t_1} \|e\|\,\mathrm{d}r$.

Status. Proved and checked, including the leak clause, a tight leak instance with $e \equiv 0$ and a tight velocity instance. Novelty (literature audit): elementary.

Proof. (i) By Lemma P.1 and Lemma B.2(b),

$$\|\dot\vartheta_i\|_{G^{-1}} \leq \|\nabla g_t(\vartheta_i)\|_G \leq \lambda_G^{1/2}\|\nabla g_t(\vartheta_i)\| \leq \lambda_G^{1/2} L_1 \|e(t)\|.$$

Square and average over i.

(ii) Use the identity coupling $(\vartheta_i(t_0), \vartheta_i(t_1))$ and Minkowski's inequality in $L^2\big(\frac{1}{m}\sum_i; G^{-1}\big)$. For general measures, use the Lagrangian coupling $(X_{t_0}, X_{t_1})_\#\mu_0$, or Ambrosio–Gigli–Savaré [2, Thm. 8.3.1] after the isometry $\vartheta \mapsto G^{-1/2}\vartheta$. Summing over a partition gives the length statement.

(iii) Cauchy–Schwarz.

(i′) Apply Lemma P.1 with $y = \nabla g_t - \sigma_m G^{-1}(\vartheta_i - \vartheta^0_{\tau(i)})$. Then Minkowski's inequality gives $\|y\|_G \leq \|\nabla g_t\|_G + \sigma_m\|\vartheta_i - \vartheta^0_{\tau(i)}\|_{G^{-1}}$. Average over i with τ optimal. At ties, Filippov velocities are convex combinations of fields with τ optimal at the current point. Each such field satisfies (i′), and the norm is convex.

(ii′) and (iii′) follow as (ii) and (iii).

For L_G: $\|\nabla g\|_G^2 = g_a\|\varphi e\|^2 + (a^\top e)^2\big(g_c\|\nabla_c\varphi\|^2 + g_s(\partial_s\varphi)^2\big) \leq \|e\|^2 L_G^2$. □

Reading. Along (L), moving the kernel measure by W costs at least $W/(\lambda_G^{1/2}L_1)$ of integrated tracking error. This is an exchange rate between learning and tracking.

Along a leaky law the proved statement is

$$\int_{t_0}^{t_1} \big(\lambda_G^{1/2} L_1\|e\| + \sigma_m d_r\big)\,\mathrm{d}r \geq W(\mu_{t_1}, \mu_{t_0}),$$

with $d_r = W(\mu_r, \mu^0)$ for (L_σ) and $d_r = d_{\mathrm{lab}}(r)$ for the labelled leak. This gives no lower bound on $\int\|e\|$, because the leak can pay for the whole motion. Under (L_σ) there are runs with $e \equiv 0$ and $W(\mu_{10}, \mu_0) \approx 1.05$ (found numerically).

Numerical checks. The following are checked at every sample (with the particle path length $\geq$ the Wasserstein length):

- (i) and Lemma P.1, including a tight instance with ratio ≥ 0.99;
- (ii) along (L);
- (ii′) along (L_σ), on a grid of pairs, including an instance with $e \equiv 0$ where it is tight (ratio ≥ 0.99) and the (L) reading fails.

Checked by: `tests/test_transport.py`:
`test_lemma_P1_projection_is_nonexpansive_and_tight_velocity_bound` ((i) and Lemma P.1);
`test_theorem_F_wasserstein_length_bounded_by_integrated_error` ((ii) along (L));
`test_theorem_F_leak_clause_along_ot_sigma_mod` and
`test_theorem_F_leak_clause_is_tight_when_the_leak_pays_for_all_motion` ((ii′) along (L_σ)).

16 Proposition G: the slow geometry

This section concerns joint descent, and variable projection (VarPro) in the induced metric.

Setting.

- $f = f(x)$ is time-invariant and C^2.
- x_m is T_0-periodic, with $\omega_{\mathrm{ref}} := 2\pi/T_0$ and occupation measure $\rho := \frac{1}{T_0}\int_0^{T_0} \delta_{x_m(t)}\,\mathrm{d}t$.
- Law (L) acts in the interior, with gains $(g_a, \delta\gamma_c, \delta\gamma_s)$. Put $\Gamma := \mathrm{diag}(\gamma_c I_n, \gamma_s)$, acting per kernel, and $G_\xi := \delta\Gamma$. Write the geometry as $\xi := (c, s)$.
- $\bar{L}(a, c, s) := \frac{1}{2}\int \|f - \frac{1}{m}\sum_i a_i \varphi(\cdot\,; c_i, s_i)\|^2\,\mathrm{d}\rho$, and $M(c, s) := \frac{1}{m^2}\int \varphi\varphi^\top\,\mathrm{d}\rho \otimes I_n$ is the Hessian of $\bar{L}$ in a.
- $\mathcal{K}$ is the set of geometries ξ on which (H1) holds, and $\mathcal{K}' \Subset \mathcal{K}$ is compact. (The set $\mathcal{K}$ is not the functional $\mathcal{G}(\mu)$ of Section 17.)
- $a^*(\xi) := \mathrm{argmin}_a\, \bar{L}$, and $L^*(\xi) := \bar{L}(a^*(\xi), \xi)$ is the profile (variable-projection) loss.
- Three dimensionless parameters:
  - the rate ratio $R := g_a m_0 \big/ \big(\delta \cdot \sup_{\mathcal{K}'} \|\Gamma^{1/2}\nabla^2 L^* \Gamma^{1/2}\|\big)$;
  - the slope $S := (\delta/g_a) \cdot \sup_{\mathcal{K}'} \|D_\xi a^* \cdot \Gamma^{1/2}\|^2$;
  - the averaging parameter $\varepsilon_{\mathrm{avg}} := (m/k_e)\,\lambda_{\max}(G^{1/2}\nabla^2\bar{L}\,G^{1/2})/\omega_{\mathrm{eff}}$, the second ratio in (H2a′). The result tables in `results/variable_projection` report its inverse, with ω_{ref} in place of ω_{eff}.

Hypotheses.

(H0) On the interval considered, the closed-loop and reduced geometries stay in $\mathcal{K}'$. The parameters stay in $\mathrm{int}\,\Theta_K^\varepsilon$, and $\sup_{\mathcal{K}'} \max_i \|a_i^*\| < a_{\max} - \varepsilon_K$.

(H1) $M \succeq m_0 I$ on $\mathcal{K}$.

(H2a′) *Two scales.* $k_e \gg \omega_{\mathrm{eff}}$ and $\varepsilon_{\mathrm{avg}} \ll 1$, where

$$\omega_{\mathrm{eff}} := \max\Big\{\omega_{\mathrm{ref}},\ \frac{\sup_t\|\dot{x}_m\|}{s_{\min}},\ \sup_X\|\partial_x(f - \hat{f})\|,\ \frac{\sup_t\|\frac{\mathrm{d}}{\mathrm{d}t}r(x_m(t))\|}{\sup\|r\|}\Big\}.$$

The last entry is the bandwidth of the residual along the reference. Also $\sup\|r\|/k_e \ll s_{\min}$.

(H2b) *Rate separation.* $R \gg 1$: the slowest weight rate $g_a m_0 m/k_e$ is much larger than the geometry rate $(m\delta/k_e)\|\Gamma^{1/2}\nabla^2 L^*\Gamma^{1/2}\|$.

(H2c) *Flat slow manifold.* $S \ll 1$. Since $Da^* = -M^{-1}\partial^2_{a,\xi}\bar{L}$, a checkable sufficient condition is $\delta \cdot \max(\gamma_c, \gamma_s) \cdot \|\partial^2_{a,\xi}\bar{L}\|^2 \ll g_a m_0^2$.

Three remarks on the hypotheses.

- **(H2b) is about $\lambda_{\min}$.** It is a condition on $\lambda_{\min}$ of the feature Gram matrix. Since $0 < \varphi \le 1$, $m_0 \le 1/m^2$. So the slowest weight rate is at most $g_a/(k_e m)$, and overlapping kernels make (H2b) fail.
- **The ratios are gain choices.** R and S contain neither k_e nor ω_{ref}. The separation between weights and geometry is therefore set by the gains.
- **(H2b) does not imply (H2c).** The product

$$R \cdot S = m_0 \cdot \sup\|Da^*\Gamma^{1/2}\|^2 \big/ \sup\|\Gamma^{1/2}\nabla^2 L^*\Gamma^{1/2}\|$$

does not depend on g_a, δ or k_e. It is fixed by the problem and by the shape of Γ: in the slope configuration below, a numerical check gives values from 166 to 1149 over $\gamma_s/\gamma_c \in [10^{-4}, 10^4]$. It is large whenever refitting the weights largely compensates a change of geometry.

Proposition G (formal reduction)**.**

(a) Two scales, under (H0) and (H2a′). *To leading order in* ω_{eff}/k_e *and* ε_{avg}*, the period-averaged closed loop is the joint gradient flow*

$$\dot{\vartheta}_i = -\frac{m}{k_e}\, G\, \nabla_{\vartheta_i} \bar{L}(\vartheta_1, \dots, \vartheta_m). \tag{16.1}$$

Its measure form is $-\frac{1}{k_e} G \nabla (\delta \bar{L}/\delta \mu)$*.*

- *This is the drift of* $(\mathfrak{A})$ *at* $\sigma = \tau = 0$*, with* Φ *in place of* $\tilde{\Phi}$ *and* (a, c, s) *in place of* (α, γ, κ)*, by the same averaging computation as Theorem H(i).*
- *It is not literally a special case of H(i): the saturated coordinates there induce a different, state-dependent metric.*
- $\bar{L}$ *is non-increasing along the averaged flow. Along the closed loop,* $\bar{L}$ *at period-averaged parameters decreases while the averaged dissipation* $\frac{m}{k_e}\|\nabla \bar{L}\|_G^2$ *dominates the* $O(\omega_{\text{eff}}/k_e)$ *corrections of Proposition H.7 and Remark H.7a. Near equilibrium it can increase by* $O((\omega_{\text{eff}}/k_e)^2)$*, as the weights settle at their shifted equilibrium (in a numerical check:* $+8 \cdot 10^{-7}$ *over 35 periods at* $k_e = 20$*, and* $3.7 \cdot 10^{-8}$*, about 22 times less, at* $k_e = 80$*).*

(b′) Three scales, for the averaged system (a), under (H0), (H1), (H2a′) and (H2b). *After a boundary layer, the weights follow the slow manifold:*

$$\begin{aligned} a - a^*(\xi) &= -\frac{k_e}{m g_a} M^{-1} Da^* \cdot \dot{\xi} + \text{h.o.t.} \\ &= \frac{\delta}{g_a} M^{-1} Da^* \Big(\Gamma^{-1} + \frac{\delta}{g_a}(Da^*)^\top Da^*\Big)^{-1} \nabla L^*(\xi) + \text{h.o.t.} \end{aligned} \tag{16.2}$$

This reduces to $\frac{\delta}{g_a} M^{-1} Da^*\, \Gamma\, \nabla L^*(\xi)$ *when* $S \ll 1$*. The geometry follows, to leading order in* $1/R$ *and for every* S*,*

$$\dot{\xi} = -\frac{m\delta}{k_e}\Big(\Gamma^{-1} + \frac{\delta}{g_a}(Da^*)^\top Da^*\Big)^{-1} \nabla L^*(\xi). \tag{16.3}$$

- *This is the gradient flow of* $(m/k_e)L^*$ *in the metric that the* G^{-1} *metric induces on the graph* $\{(a^*(\xi), \xi)\}$*.*
- *Its metric tensor is* $G_\xi^{-1} + g_a^{-1}(Da^*)^\top Da^*$*, and it couples all kernels through* Da^**.*
- *In the closed loop, the weight equilibrium is further shifted by* $O(\omega_{\text{eff}}/k_e)$ *(Proposition H.7). This shift can exceed the offset displayed above: in a numerical check on the* `satisfying` *run it is* $3\text{–}6 \cdot 10^{-3}$*, against an offset of* $4 \cdot 10^{-5}\text{–}3 \cdot 10^{-4}$*.*

(b) Variable projection, under (H2c) in addition. *Let* $(\bar{c}, \bar{s})$ *solve* $\bar{\xi}' = -\frac{m\delta}{k_e}\Gamma \nabla L^*(\bar{\xi})$*, that is,* $\bar{c}_i' = -\frac{m\delta\gamma_c}{k_e}\nabla_{c_i} L^*$ *and* $\bar{s}_i' = -\frac{m\delta\gamma_s}{k_e}\partial_{s_i} L^*$*, with* $(\bar{c}, \bar{s})(0) = (c, s)(0)$*. Then*

$$\sup_{t \le \tau_1 k_e/(m\delta)} \|(c,s)(t) - (\bar{c},\bar{s})(t)\| \to 0 \quad \text{as } (\omega_{\text{eff}}/k_e,\ \varepsilon_{\text{avg}},\ S,\ 1/R) \to 0, \text{ with } \tau_1 \text{ fixed.} \tag{16.4}$$

- *Formally the error is* $O(\omega_{\text{eff}}/k_e + \varepsilon_{\text{avg}} + S + 1/R)$*.*
- L^* *is non-increasing along* $\bar{\xi}$*, and the weights track* a^**.*
- *This flow is the* $S \to 0$ *limit of (b′).*

Status. Formal (a derivation without error control); checked directly, including a counterexample to plain VarPro at $S = O(1)$ (see the numerics below). Novelty (literature audit): a fixed-m, closed-loop analogue of the two-timescale/VarPro training of [38], [4].

No mean-field form. For fixed m, on configurations of m distinct geometries in $\mathcal{K}$, the flow (b) is the gradient flow of L^*/k_e in the metric $(1/m)\sum_i \|\dot{\xi}_i\|^2_{G_\xi^{-1}}$. Locally, that metric is the Wasserstein metric restricted to uniform m-atom measures.

- L^* is the unregularised profile loss. It is the $\lambda \to 0$ limit of $\lambda\mathcal{L}^\lambda$ of [4] (time rescaled by λ). It is not the $\lambda \to 0$ functional $\mathcal{L}^0$ of [4], which is a min-norm interpolation functional whose flow is an ultra-fast diffusion.
- As a functional of the geometry marginal, L^* depends only on the support, because the free weights absorb the masses. So it is not W-continuous, and the statement has no mean-field form. This is consistent with $m_0 \le 1/m^2$.

Derivation (formal). Each step is a standard theorem on its own time scale. Their composition on the $O(k_e/(m\delta))$ horizon is formal.

1. *Error channel (exact identity).* For $w := e - r(x,\vartheta)/k_e$, $\dot{w} = -k_e w - \dot{r}/k_e$. So $\|w(t)\| \le e^{-k_e t}\|w(0)\| + \sup\|\dot{r}\|/k_e^2$, and
$$e = r(x_m)/k_e + O\big(\sup\|\dot{r}\|/k_e^2 + \sup\|\partial_x r\| \cdot \sup\|r\|/k_e^2\big).$$
This is $O(\omega_{\text{eff}} \cdot \sup\|r\|/k_e^2)$, with the bandwidth term in ω_{eff}.
2. *Period averaging of the joint system* (Khalil [31, Thm. 10.4]; Sastry–Bodson [54, Ch. 4]). This gives (a), on $[0, b/\varepsilon_{\text{avg}}]$.
3. *Weights against geometry.* Use Tikhonov in slow time (Kokotović–Khalil–O'Reilly [32]) with the uniformly exponentially stable averaged weights on $\mathcal{K}'$ (H1).
   - On the slow manifold $a = h(\xi)$, $\dot{a} = Dh\,\dot{\xi} = -\frac{mg_a}{k_e} M(h - a^*)$, so $h - a^* = -\frac{k_e}{mg_a} M^{-1} Dh\,\dot{\xi}$. This must be evaluated with the (b′) velocity, not the VarPro velocity.
   - Since $\bar{L}$ is quadratic in a and $\partial^2_{\xi a}\bar{L} = -(Da^*)^\top M$, we get $\nabla_\xi \bar{L}(h,\xi) = \nabla L^* - (Da^*)^\top M(h - a^*) + O((\delta/g_a)^2)$.
   - Substituting into $\dot{\xi} = -\frac{m\delta}{k_e}\Gamma\nabla_\xi \bar{L}(h,\xi)$, with $Dh = Da^* + O(1/R)$, gives (b′). The resummed form solves the invariance equation exactly once $Dh - Da^*$ and the quadratic term are dropped, so the remaining corrections are in $1/R$ (they involve $D^2 a^*$ and $\partial_\xi M$), not in S.
4. *Envelope theorem.* $\nabla_\xi \bar{L}(a^*(\xi),\xi) = \nabla L^*(\xi)$. When $S \ll 1$, the metric correction in (b′) drops out, which gives (b).

The composition requires the averaged system to remain valid up to $t \sim \tau_1 k_e/(m\delta)$, which is much longer than $b/\varepsilon_{\text{avg}}$. Step 2 does not certify this. A proof would combine:

- a near-identity averaging transform;
- uniform exponential stability of the averaged weights on $\mathcal{K}'$;
- Tikhonov in slow time with an $O(\varepsilon_{\text{avg}})$ perturbation;
- Grönwall in ξ (Sastry–Bodson [54, §4.4]; Sanders–Verhulst–Murdock [51, Ch. 5]).

□

Numerics. The results are in `results/variable_projection` of the accompanying repository; Table 12 summarises the comparison.

- The first two rows use centred differences of period means.
- The slope rows use the displacement between the first and last period means. They start on the slow manifold of (b′), so no weight boundary layer enters the comparison.

- (b′) was evaluated only in the slope rows.
- k_e/ω_{eff} is 2.0 in the first two rows with the box's $s_{\min}$, and 2.6–4.8 with the actual widths, so (H2a′) holds only marginally there. In the `satisfying` run the residual bandwidth, about 7.7, then dominates.
- k_e/ω_{eff} is about 16 in the slope rows.

configuration	(H2b) ratio	S	G(a): closed loop against the joint flow (a)	G(b) and the induced-metric flow (b′)
`violating` ($m = 4$, overlapping)	0.10–0.23	5.4–25	joint cosine 0.9994 / 0.9985, size ratio 1.00 / 0.94	(b) fails: cosine −0.12 / 0.39, and L^* *increases* from $3.8 \cdot 10^{-3}$ to $5.9 \cdot 10^{-3}$
`satisfying` ($m = 2$)	79–87	0.006	joint cosine 0.985 / 0.962, size 0.96 / 0.94; geometry block 0.999 / 0.998; the weight block fails (see below)	(b) holds: cosine 0.999, ratio 0.97; L^* decreases from $1.20 \cdot 10^{-2}$ to $4.11 \cdot 10^{-3}$
`slope`, $\delta = 0.47$ ($m = 1$)	81–178	2.0–2.8	cosine 0.9992, size 1.10	(b) fails: ratio 0.42, cosine 0.93, and the centre moves the opposite way; (b′): cosine 0.9986, ratio 1.12
`slope`, $\delta = 0.47$, all gains $\div 4$	81–178	2.0–2.8	cosine 1.0000, size 1.05	(b) fails: ratio 0.40, cosine 0.92; (b′): cosine 0.9999, ratio 1.065
`slope`, $\delta = 0.047$	654–1779	0.20–0.32	cosine 1.0000, size 1.04	(b): ratio 0.85, cosine 0.9986; (b′): ratio 1.05, cosine 1.0000
`slope`, $\delta = 0.0047$	13162–17790	0.020–0.023	cosine 1.0000, size 1.04	(b): ratio 1.02; (b′): ratio 1.04; both cosines 1.0000

Table 12: Proposition G: the closed loop against the averaged predictions. In the first two rows, an entry 'p / q' gives the medians over the first quarter (p) and over the last quarter (q) of the run.

- **Joint descent.** In all runs, $\bar{L}$ at the period-mean parameters is non-increasing, sampled once per period.
- **(H2b) is not enough.** The `slope` rows satisfy (H2b) by a wide margin, yet (b) fails until S is small. Meanwhile (b′) tracks the closed loop at every S. So (H2c) is the hypothesis missing from an earlier formulation of Proposition G(b), and (b′) is the correct leading-order slow flow. The configuration is a counterexample to that earlier formulation, made reproducible in `experiments/exp_slow_manifold_slope.py`.
- **Error decomposition** (`slope`, $\delta = 0.47$, started on the slow manifold):
  - $\|(\text{a})\|/\|(\text{b}')\| = 1.015$. This is the intrinsic $O(1/R)$ error of (b′).
  - $\|\text{closed}\|/\|(\text{a})\| = 1.10$. It falls to 1.05 when all gains are divided by 4, which leaves R and S unchanged and divides $\varepsilon_{\text{avg}} \approx 0.08$ by 4. So this part is the averaging error.
  - The remaining $\approx 4.4\%$ is the $O(1/k_e)$ error-channel factor at $k_e = 20$. It is the same at $\delta = 0.047$ and 0.0047.
- **The size ratio in the `satisfying` run.** The centred ratio of 0.97 is the error-channel factor times a +1.4% averaging correction from the weights' oscillation within each period, the analogue of Remark H.7a.

  - The frozen-parameter ratio is 0.951–0.956 along the run, and 0.996 at $k_e = 80$.
  - The fastest weight rate over ω_{ref} is about 0.14.
  - In a further numerical check, with the weights running at frozen geometry the ratio is 0.969–0.970; with $g_a/4$ it is 0.956–0.960.
  - The slope correction is below 0.01% here, because ∇L^* is nearly orthogonal to the range of $(Da^*)^\top$. An earlier estimate of 0.94–0.95 contained a 2.5% forward-difference bias.

- **The weight block.** In the `satisfying` run the weights sit at $a^* + O(1/k_e)$, within 0.1% of a^*.
  - Their actual drift is tiny: 0.001–0.008 of the geometry speed in the G^{-1} norm.
  - The prediction $-(m/k_e)g_a\nabla_a\bar{L}$ at the period-mean weights is 0.17–0.30 of the geometry speed, 40–200 times the actual drift. It is the restoring velocity of the $O(1/k_e)$ offset, which (a) does not contain.
  - So the weight block fails (cosine -0.21 / 0.50, carrying 92–95% of the residual), while the geometry block matches.

Checked by: `experiments/exp_variable_projection.py`, which writes `table_violating.md` and `table_satisfying.md`; and `experiments/exp_slow_manifold_slope.py`, which writes `table_slope.md` (all in `results/variable_projection`).

Status.

- **What the loop supplies.** The error-channel and averaging scales, and the data distribution (the occupation measure of the reference), come from the closed loop.
- **What the designer supplies.** The separation between weights and geometry is a gain choice, as in [38], [4]: $\delta \ll g_a \cdot \min\big(m_0/\|\Gamma^{1/2}\nabla^2 L^*\Gamma^{1/2}\|,\ 1/\|Da^*\Gamma^{1/2}\|^2\big)$. Equivalently, $\|G_\xi^{1/2}\nabla^2 L^* G_\xi^{1/2}\| \ll g_a m_0$ and $\|Da^* G_\xi^{1/2}\|^2 \ll g_a$.
- **What is new here.**
  - The learner's data distribution is its own excitation, and at finite k_e it moves with the parameters (Proposition H.7(c)).
  - The slow flow is variable projection in the *induced* metric, valid for every S. The slope S measures the departure from plain VarPro.

17 Theorem H: Langevin adaptive control and global convergence to the regularised optimum

Why something new is needed. Theorems B–G are local, or describe the dynamics. A global statement about kernel placement needs either the Chizat–Bach theory or entropy.

- **Boundary conditions.** The Chizat–Bach theory for noiseless Wasserstein flows (partially 1-homogeneous case, [8]) requires boundary conditions: either Neumann conditions on a bounded convex domain, or C^1-uniform behaviour at infinity. Gaussian kernels with free centres and widths satisfy neither.
- **Separation.** Its Theorem 3.5 also needs the support of μ_0 to separate $\{-r_0\} \times \Theta$ from $\{r_0\} \times \Theta$. A Dirac start violates this, and that is an obstruction distinct from the boundary conditions.

- **The collapsed configuration is invariant.** It remains a moving Dirac mass and never separates without noise, by permutation symmetry and uniqueness of solutions. The test `test_noiseless_symmetric_initialisation_never_separates_but_dither_does` in the accompanying repository checks this on the real closed loop with $m = 6$. The rows stay bit-identical at every snapshot, the common particle moves, and dither separates the rows within 0.1 s.

Entropy removes the obstruction: Theorem H(ii) covers Dirac starts after any positive time.

Design. The particles are unconstrained, $\vartheta = (\alpha, \gamma, \kappa) \in \mathbb{R}^p$, and the features are saturated coordinate by coordinate:

$$a = a_{\max} \tanh(\alpha/a_{\max}), \qquad c = c_{\max} \tanh(\gamma/c_{\max}), \qquad s = s_{\min} + (s_{\max} - s_{\min}) \cdot \mathrm{sig}(\kappa).$$

Here $a_{\max}$ bounds each coordinate of a, unlike Section 9, where it bounds $\|a\|$. So $\tilde{\Phi}(x; \vartheta) = a\varphi(x; c, s)$ satisfies $\|\tilde{\Phi}\| \le K_\Phi := a_{\max}\sqrt{n}$, and every ϑ-derivative of $\tilde{\Phi}$ is bounded uniformly in x. The law is

$$\mathrm{d}\vartheta_i = \big[\, G\nabla_\vartheta g_t(\vartheta_i) - \sigma(\vartheta_i - \vartheta^0) \,\big]\,\mathrm{d}t + \sqrt{2\tau}\, G^{1/2}\,\mathrm{d}B_i, \qquad g_t(\vartheta) := e(t)^\top \tilde{\Phi}(x(t); \vartheta). \tag{L_τ}$$

It has three parts: the classical gradient law, σ-modification written in the G^{-1} metric, and parameter dither of temperature τ.

Averaged mean-field system. Assume $f = f(x)$ is continuous and time-invariant, as in Proposition G (the functions `occupation_loss` and `first_variation` of the accompanying code assume this). Let ρ be the occupation measure of a T_0-periodic reference, and $F_\rho := \sup_{\mathrm{supp}\,\rho} \|f\|$. Define

$$r_\mu := f - \int \tilde{\Phi}(\cdot\,; \vartheta')\,\mathrm{d}\mu(\vartheta'), \qquad \bar{L}(\mu) := \frac{1}{2}\int \|r_\mu\|^2\,\mathrm{d}\rho,$$
$$V_L[\mu](\vartheta) := -\int r_\mu(x)^\top \tilde{\Phi}(x; \vartheta)\,\mathrm{d}\rho(x), \qquad V[\mu] := \frac{1}{k_e} V_L[\mu] + \frac{\sigma}{2}\|\vartheta - \vartheta^0\|^2_{G^{-1}},$$

and the averaged system

$$\partial_t \mu = \nabla \cdot \big(\mu G \nabla V[\mu]\big) + \tau \nabla \cdot (G \nabla \mu). \tag{$\mathfrak{A}$}$$

Remark H.0 (from the closed loop to ($\mathfrak{A}$): a formal reduction with a measured error)**.**

- **An algebraic identity.** Substitute $e := r_\mu(x_m)/k_e$ and $x := x_m$ in the drift of (L_τ), and average over one reference period at frozen μ. The result is exactly $-G\nabla V[\mu]$. The test `test_averaged_drift_is_the_mean_field_langevin_drift` checks this identity to 10^{-7}. It does not test averaging.
- **The formal reduction.** ($\mathfrak{A}$) is the leading-order description of the mean-field closed loop under two conditions:
  - *Tikhonov:* $k_e \gg \max\big(\omega_{\mathrm{ref}},\ \mathrm{Lip}(f) + K_\Phi/s_{\min},\ (F_\rho + K_\Phi)/s_{\min}\big)$, so that the shift $x - x_m = O(1/k_e)$ is small on the kernel scale.
  - *Period averaging:* $T_0 \cdot \sup_\vartheta \|(1/k_e) G \nabla V_L[\mu](\vartheta)\| \ll \ell$, $\sigma T_0 \ll 1$ and $\tau \lambda_G T_0 \ll \ell^2$, where ℓ is the ϑ-scale of $\tilde{\Phi}$. The left-hand side is at most $(T_0/k_e)\lambda_G(F_\rho + K_\Phi) \cdot \sup\|\nabla_\vartheta \tilde{\Phi}\|$. The confinement drift does not depend on time and needs no averaging.
- **The first corrections.**
  - At frozen μ (and frozen ϑ) they are $O(1/k_e)$ relative to the drift, do not depend on the gains, and are given exactly by Proposition H.7.

– Along the loop, the within-period oscillation of the parameters adds the second-order averaging term b_{osc} of Remark H.7a. Its size relative to the frozen terms is $O(|G|T_0)$, so it does depend on the gains.
– In the formal limit below, both are $O(k_e^{-2})$.

Measured relative errors of the frozen-μ averaged drift against $(\mathfrak{A})$, with b_{osc} excluded:

instance	$k_e = 10$	20	40	80	$k_e \times$ error
$n = 1$	4.2%	2.0%	0.91%	0.48%	≈ 0.4
$n = 2$, ellipse	43%	21%	10%	5.1%	≈ 4.2

Tests: `test_averaging_error_of_the_formal_reduction_decreases_like_one_over_ke` (instance $n = 1$) and `test_averaged_drift_error_decays_like_one_over_ke_for_an_orbit_enclosing_area` (instance $n = 2$; $k_e = 10, 20, 40, 80$).

– $k_e = 10$ is outside the averaging regime for the second instance.
– Its larger error does not come from the enclosed area. Proposition H.7 attributes it to the orientation-even state-shift term: it predicts $k_e \times$ error ≈ 4.05, of which U_{odd} alone would give 1.5, and a numerical check on the reversed orbit gives 4.0.
– A numerical check shows that at the gains of these instances b_{osc} is not small. k_e times its relative size is 0.92 ($n = 1$, $G = \mathrm{diag}(4, 1, 1)$) and 4.4 ($n = 2$, $G = I$), against 0.42 and 4.05 for the frozen terms.

- **The exact formal limit** is $k_e \to \infty$ with G, $k_e\sigma$ and $k_e\tau$ fixed, in slow time t/k_e (Conjecture H.6).
- **Not proved here.** The mean-field limit of (L_τ) itself is not proved; Theorem D covers the deterministic projected law (L).

Status. Formal reduction, with a measured error.

Theorem H. *Assume the setting above, and $\sigma > 0$, $\tau > 0$ where stated.*

(i) Identification. *$V[\mu]$ is the first variation of $\mathcal{G}(\mu) := \frac{1}{k_e}\bar{L}(\mu) + \frac{\sigma}{2}\int \|\vartheta - \vartheta^0\|^2_{G^{-1}}\,\mathrm{d}\mu$. Hence $(\mathfrak{A})$ is the mean-field Langevin dynamics, in the G^{-1} metric, of the free energy $F_\tau := \mathcal{G} + \tau H$, with $H(\mu) = \int \mu \log \mu$. Equivalently, with $\gamma_\tau := \mathcal{N}(\vartheta^0, (\tau/\sigma)G)$,*

$$F_\tau = \frac{1}{k_e}\bar{L} + \tau H(\cdot \,|\, \gamma_\tau) + \mathrm{const.}$$

F_τ is well defined on all of $\mathcal{P}(\mathbb{R}^p)$, and finite exactly when $H(\mu \,|\, \gamma_\tau) < \infty$.

(ii) Global exponential convergence ($\sigma, \tau > 0$). *F_τ has a unique minimiser, $\mu_\tau^* \propto e^{-V[\mu_\tau^*]/\tau}$. For every $\mu_0 \in \mathcal{P}_2(\mathbb{R}^p)$ with $F_\tau(\mu_0) < \infty$,*

$$F_\tau(\mu_t) - F_\tau(\mu_\tau^*) \le e^{-2\tau\rho_\tau t}\big(F_\tau(\mu_0) - F_\tau(\mu_\tau^*)\big),$$
$$W_{G^{-1}}(\mu_t, \mu_\tau^*)^2 \le \frac{2}{\tau\rho_\tau}\, e^{-2\tau\rho_\tau t}\big(F_\tau(\mu_0) - F_\tau(\mu_\tau^*)\big).$$

Here ρ_τ is a log-Sobolev constant, for the whitened measures, of every proximal Gibbs measure $\hat{\mu} \propto e^{-V[\mu]/\tau}$. In ϑ-coordinates this means Fisher information in the G metric. It satisfies

$$\rho_\tau \ge \frac{\sigma}{\tau} \cdot \exp\Big(-\frac{2K_\Phi(F_\rho + K_\Phi)}{k_e\tau}\Big).$$

Suppose μ_0 is only in $\mathcal{P}_2(\mathbb{R}^p)$, for example a Dirac mass or an empirical measure, so that $H(\mu_0) = +\infty$. Then $F_\tau(\mu_{t_0}) < \infty$ for every $t_0 > 0$, and the estimates hold with (μ_0, t) replaced by $(\mu_{t_0}, t - t_0)$.

(iii) Loss and tracking. *Let* $\bar{e}_t := r_{\mu_t}/k_e$. *Then* $\langle\|\bar{e}_t\|^2\rangle_\rho = 2\bar{L}(\mu_t)/k_e^2$, *and*

$$\begin{aligned}|\bar{L}(\mu_t) - \bar{L}(\mu_\tau^*)| &\le (F_\rho + K_\Phi)L_\Phi \cdot W_{G^{-1}}(\mu_t, \mu_\tau^*)\\ &\le (F_\rho + K_\Phi)L_\Phi\Big(\frac{2}{\tau\rho_\tau}\Big)^{1/2} e^{-\tau\rho_\tau t}\big(F_\tau(\mu_0) - F_\tau(\mu_\tau^*)\big)^{1/2},\end{aligned}$$

where $L_\Phi := \sup_{x,\vartheta,\|u\|=1}\|G^{1/2}\nabla_\vartheta(u^\top\tilde{\Phi})\|$. *The regularisation bias satisfies*

$$\bar{L}(\mu_\tau^*) \le \inf_\nu\big\{\bar{L}(\nu) + k_e\tau H(\nu\,|\,\gamma_\tau)\big\}.$$

The link $e \approx \bar{e}$ *between* $(\mathfrak{A})$ *and the closed loop is formal (Remark H.0).*

(iv) Annealing. *Let* $\mu_0 \in \mathcal{P}_2$ *with* $F_{\tau_{\text{init}}}(\mu_0) < \infty$. *Let* (τ_t) *be smooth and decreasing from* τ_{init}, *with* $\tau_t = \alpha_a/\log t$ *for* $t \ge t_1 > 1$, *where* $\alpha_a > \alpha^* := 2K_\Phi(F_\rho + K_\Phi)/k_e$. *Then* $\mathcal{G}(\mu_t) - \inf\mathcal{G} \le C' \cdot \log\log t/\log t$, *where* $\inf\mathcal{G}$ *is the* σ*-regularised optimum. This is Theorem 4.1 of [9]. Its hypotheses hold:*

- $\mathcal{G} \ge 0$*;*
- *Assumptions 1–3 hold for every* $\tau > 0$*;*
- *for* $\tau \in (0, \tau_{\text{init}}]$, *(ii) gives* $\rho_\tau \ge C_0 e^{-\alpha^*/\tau}$ *with* $C_0 = \sigma/\tau_{\text{init}}$, *which is all its proof uses.*

Status. Proved for $(\mathfrak{A})$; the reduction to $(\mathfrak{A})$ is formal (Remark H.0); real loop: Conjecture H.6. Novelty (literature audit): a bridge to [9], [41]; new in adaptive control.

Checked by: `tests/test_langevin.py`:
`test_first_variation_bound_used_in_holley_stroock` (the oscillation bound in (ii));
`test_lsi_lower_bound_formula` (formula transcription only);
`experiments/exp_langevin_global.py`, `exp_langevin_tradeoff.py` and `exp_gibbs_reference.py` (global behaviour, speed–bias trade-off, reference values).

Proof. (i) We have

$$\frac{\mathrm{d}}{\mathrm{d}\varepsilon}\bar{L}\big((1-\varepsilon)\mu + \varepsilon\nu\big)\Big|_{\varepsilon=0} = -\int r_\mu^\top\Big(\int\tilde{\Phi}\,\mathrm{d}(\nu-\mu)\Big)\,\mathrm{d}\rho = \int V_L\,\mathrm{d}(\nu-\mu).$$

This is exact, because $\bar{L}$ is quadratic in μ. The confinement is linear in μ. In the G^{-1} metric, $\nabla\cdot\big(\mu G\nabla(V+\tau\log\mu)\big) = \nabla\cdot(\mu G\nabla V) + \tau\nabla\cdot(G\nabla\mu)$. Since $\log\gamma_\tau = -(\sigma/(2\tau))\|\vartheta-\vartheta^0\|^2_{G^{-1}} + \text{const}$, we get $\tau H(\mu\,|\,\gamma_\tau) = \tau H(\mu) + (\sigma/2)\int\|\vartheta-\vartheta^0\|^2_{G^{-1}}\,\mathrm{d}\mu + \text{const}$.

(ii) Whiten with $\zeta := G^{-1/2}\vartheta$. The metric becomes Euclidean, the confinement becomes $(\sigma/2)\|\zeta-\zeta^0\|^2$, $W_{G^{-1}}$ becomes W_2, and H shifts by $\frac{1}{2}\log\det G$. $(\mathfrak{A})$ becomes equation (12) of [9]. We verify Assumptions 1–3 of [9].

- *Assumption 1.* $\nabla V[\mu](\zeta)$ is Lipschitz in (ζ,μ). The constants do not depend on μ, because $\|r_\mu\| \le F_\rho + K_\Phi$ on $\operatorname{supp}\rho$ for every $\mu \in \mathcal{P}(\mathbb{R}^p)$.
  - In ζ, this holds because $\nabla^2\tilde{\Phi}$ is bounded (saturation, and $s \ge s_{\min}$).
  - In μ, it holds because $\int\tilde{\Phi}\,\mathrm{d}\mu$ is W_1-Lipschitz, $W_1 \le W_2$, and $\nabla\tilde{\Phi}$ is bounded.
- *Assumption 2.* $\mathcal{G}$ is convex: a linear map followed by a convex quadratic, plus a linear term. F_τ has a minimiser by confinement and lower semicontinuity (proof of [9, Prop. 5.1]).
- *Assumption 3.* $e^{-V[\mu]/\tau} = e^{-(\sigma/(2\tau))\|\zeta-\zeta^0\|^2}\cdot e^{-V_L[\mu]/(k_e\tau)}$.
  - The Gaussian factor satisfies LSI(σ/τ) (Bakry–Émery [3]).

– $|V_L[\mu]| \le \int \|r_\mu\| \|\tilde\Phi\| \,\mathrm{d}\rho \le (F_\rho + K_\Phi) K_\Phi$, so the perturbation has oscillation at most $2K_\Phi(F_\rho + K_\Phi)/(k_e\tau)$. Here F_ρ may be replaced by $\int \|f\| \,\mathrm{d}\rho$.

– Holley–Stroock [27] gives ρ_τ.

Theorem 3.2 and Corollary 3.3 of [9] (the latter with Otto–Villani [42]) give the two inequalities. Uniqueness follows from strict convexity of H.

$\mathcal{P}_2$ *starts.* Work in ζ-coordinates and fix the solution μ_t. Compare the particle SDE with the Ornstein–Uhlenbeck process $\mathrm{d}Y = -\sigma(Y - \zeta^0)\,\mathrm{d}t + \sqrt{2\tau}\,\mathrm{d}B$ started from the same law.

- The two drifts differ by $b_t = -(1/k_e) G^{1/2} \nabla V_L[\mu_t]$, with $\|b_t\| \le B := (1/k_e)\lambda_G^{1/2}(F_\rho + K_\Phi) \cdot \sup\|\nabla_\vartheta \tilde\Phi\|$.
- Girsanov and data processing give $H(\mu_{t_0} \,|\, \mathrm{OU}_{t_0}) \le B^2 t_0/(4\tau)$.
- OU_{t_0} has a density at most $\big(2\pi(\tau/\sigma)(1 - e^{-2\sigma t_0})\big)^{-p/2}$, so $H(\mu_{t_0}) < \infty$.
- $\mu_{t_0} \in \mathcal{P}_2$, because the drift grows linearly.

Hence $F_\tau(\mu_{t_0}) < \infty$.

(iii) $\bar L(\mu) - \bar L(\nu) = \frac{1}{2}\int (r_\mu - r_\nu)^\top (r_\mu + r_\nu)\,\mathrm{d}\rho$. Here $\|r_\mu + r_\nu\| \le 2(F_\rho + K_\Phi)$, and $\|r_\mu - r_\nu\| = \|\int \tilde\Phi \,\mathrm{d}(\nu - \mu)\| \le L_\Phi \cdot W_{1,G^{-1}} \le L_\Phi \cdot W_{G^{-1}}$. Conclude with (ii). For the bias: $(1/k_e)\bar L(\mu_\tau^*) + \tau H(\mu_\tau^* \,|\, \gamma_\tau) \le (1/k_e)\bar L(\nu) + \tau H(\nu \,|\, \gamma_\tau)$ for every ν, and $H(\mu_\tau^* \,|\, \gamma_\tau) \ge 0$.

(iv) This is Theorem 4.1 of [9] with the stated hypotheses. □

Remark H.1 (dither adds no output noise in the mean-field limit; its cost is the entropic bias)**.** In the limit, $\hat f = \int \tilde\Phi \,\mathrm{d}\mu$ is deterministic by the law of large numbers. The only effect of dither is the entropic bias, which Theorem H(iii) quantifies.

In the experiment ($\sigma = 0.02$, $\tau = 0.004$, $G = \mathrm{diag}(4, 1, 1)$, $k_e = 10$), a grid computation gives $\bar L(\mu_\tau^*) = 0.11118$. It is grid-converged (`exp_gibbs_reference.py` → `results/langevin_global/gibbs_reference.md`). At $(\sigma, \tau) = (0.002, 5 \cdot 10^{-4})$ the same computation gives 0.0085. The unregularised offline optimum over 32 kernels is about $7.6 \cdot 10^{-6}$ (best of 5 L-BFGS restarts, an upper bound), and $\bar L(0) = 0.311$.

At finite m, the fluctuation of $\hat f$ is $O(K_\Phi/\sqrt{m})$ at each fixed t (finite-horizon propagation of chaos). It is uniform in time only under Remark H.5.

Remark H.2 (the control knobs, and what the numbers say)**.**

- **The knobs.** σ is the strong-convexity constant of the confinement, τ is the temperature, and $a_{\max}$ sets K_Φ.
- **The role of k_e.** Since $k_e F_\tau = \bar L + (k_e\sigma/2)\int \|\vartheta - \vartheta^0\|_{G^{-1}}^2 \,\mathrm{d}\mu + k_e\tau H$, μ_τ^* and ρ_τ depend on (σ, τ, k_e) only through $(k_e\sigma, k_e\tau)$. Raising k_e at fixed (σ, τ) has three effects:
  – it improves ρ_τ and the guaranteed rate $2\sigma e^{-\alpha^*/\tau}$, since $\alpha^* \propto 1/k_e$;
  – it moves μ_τ^* toward the prior $\mathcal{N}(\vartheta^0, (\tau/\sigma)G)$, which increases the regularisation bias;
  – it shrinks the loss-driven drift like $1/k_e$.
- **The guaranteed rate** is exponentially small in $1/\tau$, a known feature of Holley–Stroock bounds [27]:
  – $2\tau\rho_\tau \ge e^{-8401}\,\mathrm{s}^{-1}$ at $(\sigma, \tau) = (0.002, 5 \cdot 10^{-4})$;
  – $2\tau\rho_\tau \ge e^{-1053}\,\mathrm{s}^{-1}$ at $(0.02, 0.004)$.

  The experiments do not test this rate.
- **The temperature sweep** (`results/langevin_tradeoff`; $m = 32$, $\sigma = 0.002$, $k_e = 10$, $T = 1500\,\mathrm{s}$, 2 seeds):

τ	escape time of the collapsed start (s), per seed	loss floor of the spread start
0	never, never	$7.0 \cdot 10^{-3}$
$5 \cdot 10^{-4}$	never, 1050	$1.1 \cdot 10^{-2}$
$1 \cdot 10^{-3}$	1480, 1005	$1.5 \cdot 10^{-2}$
$2 \cdot 10^{-3}$	1115, 800	$2.3 \cdot 10^{-2}$
$4 \cdot 10^{-3}$	970, 430	$3.6 \cdot 10^{-2}$
$8 \cdot 10^{-3}$	390, 295	$6.5 \cdot 10^{-2}$
annealed, hot start	990, 530	$3.2 \cdot 10^{-2}$

- **How escape is measured.** Escape is the first sample at which the collapsed run's loss falls to at most twice the spread run's final loss, at the same τ and seed. Two caveats follow:
  - the threshold grows with τ, from 0.022 to 0.13, and the escape time includes convergence after escape;
  - the $\tau = 5 \cdot 10^{-4}$ point is censored and is excluded from the fit.
- **The fit.** The seed-averaged escape time fits $t_{\text{esc}} \approx -1693 + 427 \cdot \log(1/\tau)$ (4.5% relative residual) better than an Arrhenius law (25%). On the eight per-seed values the two laws fit equally poorly (27% and 33%; `langevin_tradeoff/table.md`).
- **No saddle explanation.** The noiseless collapsed end point is a saddle of the averaged landscape, but, as a numerical check shows, a weak one. Its unstable rate is about $5 \cdot 10^{-5}\,\text{s}^{-1}$ for $(\mathfrak{A})$, and about $2 \cdot 10^{-4}\,\text{s}^{-1}$ for the exactly averaged loop at $k_e = 10$. A saddle-escape law would therefore need $\tau \lesssim 10^{-5}$, and would have a $\log(1/\tau)$ slope of 10^3–10^4 s.
- **What the data show.** At the tested temperatures escape is governed by diffusion, by the approach transient and by the τ-dependent threshold. The data do not identify the escape mechanism. An earlier "saddle, not barrier" reading of these data is withdrawn.
- **The annealed runs** use α_a between about $2 \cdot 10^{-3}$ and $2 \cdot 10^{-2}$, far below $\alpha^* = 4.2$. They are not instances of H(iv).

Remark H.3 (why saturation and not projection)**.** The mean-field Langevin results are on $\mathbb{R}^d$. Projecting the drift with the Pomet–Praly operator [50], while keeping additive noise, leaves the noise unconfined. Reflected (Skorokhod) dynamics on the convex body Θ_K would confine it.

- Such dynamics satisfy a log-Sobolev inequality with constant of order $1/\operatorname{diam}^2$, even for $\sigma = 0$.
- The argument is a diameter-based log-Sobolev inequality for the uniform measure on a convex body, followed by Holley–Stroock [27]. The diameter-based inequality comes from $\mathrm{CD}(0, \infty)$ with a convex Neumann boundary plus a diameter bound (F.-Y. Wang 1997 [57]), or from Frieze–Kannan (1999) [19] for log-concave measures on convex bodies. Together these give $\mathrm{LSI}\big(c \cdot \operatorname{diam}^{-2} \cdot e^{-2K_\Phi(F_\rho + K_\Phi)/(k_e \tau)}\big)$.
- Bakry–Émery [3] alone gives nothing at $\sigma = 0$.
- That route connects to [45] and is left open.

Remark H.4 (the role of σ-modification)**.** In the classical theory, σ-modification is a robustness device with a bias. In the saturated design on $\mathbb{R}^p$ it supplies the confinement that makes $e^{-V/\tau}$ integrable and log-Sobolev, so the global result rests on it. Its bias is the regularisation term of the variational problem that μ_τ^* solves. Under reflection on Θ_K, compactness would play that role instead.

Remark H.5 (finite m; conditional)**.** The m-particle averaged system is

$$\mathrm{d}\vartheta_i = -G\nabla V[\mu_t^m](\vartheta_i)\,\mathrm{d}t + \sqrt{2\tau}\,G^{1/2}\,\mathrm{d}B_i, \qquad \mu_t^m := \frac{1}{m}\sum_i \delta_{\vartheta_i}.$$

Uniform-in-time propagation of chaos would follow from [6], under its hypotheses.

- *Available:* convexity of $\mathcal{G}$; a uniform LSI, from (ii); and a bounded, smooth second variation $(1/k_e)\int \tilde{\Phi}(x;\vartheta)^\top \tilde{\Phi}(x;\vartheta')\,\mathrm{d}\rho$.
- *Still to check:* the remaining regularity conditions of [6], verbatim (open question 4 of `LITERATURE_TRANSPORT.md` in the accompanying repository).

Status. Conditional on the hypotheses of [6]; their remaining regularity conditions are still to be checked.

Conjecture H.6 (the real closed loop, in the scaling where $(\mathfrak{A})$ is the limit)**.** *Let $f = f(x)$ satisfy the tube condition of Theorem D(b), with $a_{\max}$ replaced by K_Φ. Fix $G \succ 0$ and $\sigma_0, \tau_0 > 0$, and set $\sigma := \sigma_0/k_e$ and $\tau := \tau_0/k_e$.*

- *Then μ_τ^*, the minimiser of $\bar{L} + (\sigma_0/2)\int \|\vartheta - \vartheta^0\|^2_{G^{-1}}\,\mathrm{d}\mu + \tau_0 H$, does not depend on k_e.*
- *In slow time $s = t/k_e$, $(\mathfrak{A})$ is the formal limit of the closed loop.*

Conjecture: *for the mean-field limit of the closed loop under (L_τ), for every $\mu_0 \in \mathcal{P}_2$ and every $e(0)$, there are $S(\mu_0)$ and $r(k_e) \to 0$ such that $W_{G^{-1}}(\mu_t, \mu_\tau^*) \le r(k_e)$ for all $t \ge k_e \cdot S$.*

Status. Conjectural.

At fixed k_e the analogous statement is false. The limit of small gains is the exactly averaged system, with the true periodic error e_μ and $x = x_m + e_\mu$.

- Its stationary state is offset from μ_τ^*: $\bar{L} = 0.11415$ against $\bar{L}(\mu_\tau^*) = 0.11118$ at $k_e = 10$ (`results/langevin_global/gibbs_reference.md`).
- In the scaling of Conjecture H.6, the $L^2(\rho)$ distance between the two learned functions is $6.8 \cdot 10^{-3}$, $1.4 \cdot 10^{-3}$ and $9.7 \cdot 10^{-4}$ at $k_e = 10$, 20 and 40. It is small and decreasing, but a clean $1/k_e$ law is not resolved in this range.
- By Proposition H.7(e) it is not the $W_{G^{-1}}$-gradient flow of any free energy. The cyclic sums of its antisymmetric second variation are non-zero in the checked instances, including the network of the experiments.

18 Finite gain: semi-gradient and non-reciprocal structure

This section continues Section 17 at finite feedback gain k_e. The notation is that of Section 9 and Section 17: the saturated features $\tilde{\Phi}$, the residual r_μ, the potential $V_L[\mu]$, the averaged system $(\mathfrak{A})$ and the law (L_τ). We write $\tilde{\Phi}_\vartheta := \tilde{\Phi}(\cdot;\vartheta)$.

18.1 Frozen-measure structure of the exactly averaged loop

Proposition H.7 (frozen-measure structure of the exactly averaged closed loop)**.** *Assume the following:*

- *$f \in C^2(\mathbb{R}^n)$ is autonomous;*
- *$x_m \in C^2$ is T_0-periodic, with oriented orbit $\mathcal{O}$;*
- *the features are the saturated features of the design (Section 17);*
- *$\mu \in \mathcal{P}(\mathbb{R}^p)$ is frozen.*

Then for every $R > 0$ there is $k_0(R)$, independent of μ, such that for $k_e \geq k_0(R)$ the following hold.

(a) Periodic error. *The frozen-μ error equation $\dot{e} = r_\mu(x_m + e) - k_e e$ has exactly one T_0-periodic solution e_μ with $\sup_t \|e_\mu\| \leq R$. It attracts exponentially every solution with $\|e(0)\| \leq R$, and*

$$e_\mu = \frac{r_\mu(x_m)}{k_e} + \frac{\nabla r_\mu(x_m)\, r_\mu(x_m) - \frac{\mathrm{d}}{\mathrm{d}t} r_\mu(x_m(t))}{k_e^2} + O(k_e^{-3}), \tag{18.1}$$

uniformly in t and μ.

- *Here ∇r_μ is the Jacobian $(\partial r_k / \partial x_l)$, and $\partial_x \tilde{\Phi} = (\partial \tilde{\Phi}_k / \partial x_l)$.*
- *If f is one-sided Lipschitz, $\langle f(x) - f(y), x - y \rangle \leq L_f \|x - y\|^2$, then for $k_e > L_f + a_{\max} \sqrt{n}\, e^{-1/2} / s_{\min}$ the periodic solution is unique and attracts every solution.*
- *Without such a condition uniqueness is only local. For $f(x) = x^2$ there is a second, repelling periodic solution near $e \approx k_e$, and solutions above it blow up.*

(b) Expansion. *The period average of the drift of* (L_τ) *along e_μ is $G \nabla_\vartheta P_\mu(\vartheta) - \sigma(\vartheta - \vartheta^0)$, where $P_\mu(\vartheta) := \langle e_\mu^\top \tilde{\Phi}(x_m + e_\mu; \vartheta) \rangle_t$ satisfies*

$$P_\mu(\vartheta) = -\frac{1}{k_e} V_L[\mu](\vartheta) + \frac{1}{k_e^2} \big(U_{\mathrm{even}}[\mu](\vartheta) + U_{\mathrm{odd}}[\mu](\vartheta) \big) + O(k_e^{-3}), \tag{18.2}$$

$$U_{\mathrm{even}}[\mu](\vartheta) := \int (r_\mu \cdot \nabla_x) \big(r_\mu^\top \tilde{\Phi}(\cdot; \vartheta) \big)\, \mathrm{d}\rho, \qquad U_{\mathrm{odd}}[\mu](\vartheta) := -\frac{1}{T_0} \oint_{\mathcal{O}} \tilde{\Phi}(\cdot; \vartheta)^\top \, \mathrm{d}r_\mu, \tag{18.3}$$

uniformly in ϑ and μ, together with ϑ-gradients.

(c) Semi-gradient form.

$$-\frac{1}{k_e} V_L[\mu] + \frac{1}{k_e^2} U_{\mathrm{even}}[\mu] = \frac{1}{k_e} \int r_\mu^\top \tilde{\Phi}(\cdot; \vartheta)\, \mathrm{d}\rho_\mu^{\mathrm{cl}} + O(k_e^{-3}). \tag{18.4}$$

Here ρ_μ^{cl} is the occupation measure of the closed-loop state $x_m + e_\mu$.

- *So, apart from the lag term U_{odd}, the averaged law follows the gradient of the closed-loop loss $\frac{1}{2} \int \|r_\nu\|^2 \, \mathrm{d}\rho_\mu^{\mathrm{cl}}$ at $\nu = \mu$, with the closed-loop state distribution ρ_μ^{cl} held fixed.*
- *This is a semi-gradient: the update freezes part of its own parameter dependence. In TD learning the frozen part is the bootstrapped target, and the data distribution does not depend on the parameters [56, 58].*
- *Here the frozen part is the data distribution ρ_μ^{cl}. This is the structure of performative prediction [46]: steps taken with the distribution held at its current value converge to performatively stable points, not to optima.*
- ***The performative defect carries the even non-reciprocity.*** *With $J^{\mathrm{cl}}(\mu) := \frac{1}{2} \int \|r_\mu\|^2 \, \mathrm{d}\rho_\mu^{\mathrm{cl}}$,*

$$\begin{aligned} &-\frac{1}{k_e} V_L[\mu] + \frac{1}{k_e^2} U_{\mathrm{even}}[\mu] \\ &\qquad = -\frac{1}{k_e} \frac{\delta J^{\mathrm{cl}}}{\delta \mu} - \frac{1}{k_e^2} \int (\tilde{\Phi}_\vartheta \cdot \nabla) \big(\tfrac{1}{2} \|r_\mu\|^2 \big)\, \mathrm{d}\rho + O(k_e^{-3}). \end{aligned} \tag{18.5}$$

The first term is a true first variation, hence reciprocal. The second is the defect: it has the same order as U_{even}, and it carries all of A_{even} in (e).

(d) Circulation kernel. $U_{\mathrm{odd}}[\mu](\vartheta) = -\frac{1}{T_0} \oint_{\mathcal{O}} \tilde{\Phi}(\cdot; \vartheta)^\top \, \mathrm{d}f + \int K(\vartheta, \vartheta')\, \mathrm{d}\mu(\vartheta')$, *with*

$$K(\vartheta, \vartheta') := \frac{1}{T_0} \oint_{\mathcal{O}} \tilde{\Phi}(\cdot; \vartheta)^\top \, \mathrm{d}\tilde{\Phi}(\cdot; \vartheta') = \Big\langle \tilde{\Phi}(x_m; \vartheta)^\top \frac{\mathrm{d}}{\mathrm{d}t} \tilde{\Phi}(x_m; \vartheta') \Big\rangle_t. \tag{18.6}$$

- *K is antisymmetric.*
- *By Stokes, $K(\vartheta,\vartheta') = \frac{1}{T_0}\int_\Sigma \sum_k \mathrm{d}\tilde\Phi_k(\cdot;\vartheta)\wedge \mathrm{d}\tilde\Phi_k(\cdot;\vartheta')$ for any 2-chain Σ with $\partial\Sigma = \mathcal{O}$.*
- *For $n = 2$ and a simple, positively oriented orbit bounding D,*

$$K(\vartheta,\vartheta') = \frac{1}{T_0}\sum_k \int_D \det\big[\nabla_x\tilde\Phi_k(\cdot;\vartheta), \nabla_x\tilde\Phi_k(\cdot;\vartheta')\big]\,\mathrm{d}A.$$

- *K is an integral over the enclosed region, not the area itself. For Gaussian features, $\sum_k \det[\nabla\tilde\Phi_k(\vartheta), \nabla\tilde\Phi_k(\vartheta')] = (a^\top a')\cdot\det[\nabla\varphi, \nabla\varphi']$, so $K = 0$ whenever $a^\top a' = 0$ or $c = c'$, whatever the area.*
- *Hence $U_{\mathrm{odd}} \equiv 0$ for $n = 1$.*
- *Reversing the orbit preserves ρ, flips U_{odd}, and leaves U_{even} unchanged.*

(e) Non-reciprocity. *Write $S(\vartheta,\vartheta') := \delta P_\mu(\vartheta)/\delta\mu(\vartheta')$ for the derivative in the direction $\delta_{\vartheta'}$, and call $S(\vartheta,\vartheta') - S(\vartheta',\vartheta)$ its antisymmetric part (without the factor $\frac{1}{2}$). Adding a function of ϑ alone, or a constant $c(\mu)$, to P_μ changes the antisymmetric part by a coboundary, which the criterion below allows.*

- *At order $1/k_e$, S is symmetric: it equals $-\frac{1}{k_e}\int \tilde\Phi(\cdot;\vartheta)^\top\tilde\Phi(\cdot;\vartheta')\,\mathrm{d}\rho$.*
- *At order $1/k_e^2$ it is not. The second variation of the k_e^{-2} coefficient $U_{\mathrm{even}} + U_{\mathrm{odd}}$ has antisymmetric part $A_{\mathrm{even}} + 2K$, with*

$$A_{\mathrm{even}}(\vartheta,\vartheta') := -\int\Big[(\tilde\Phi_{\vartheta'}\cdot\nabla)(r_\mu^\top\tilde\Phi_\vartheta) - (\tilde\Phi_\vartheta\cdot\nabla)(r_\mu^\top\tilde\Phi_{\vartheta'})\Big]\,\mathrm{d}\rho. \tag{18.7}$$

- *P_μ, up to $O(k_e^{-3})$, can be a first variation of a functional of μ only if $A_{\mathrm{even}} + 2K$ is a coboundary $a(\vartheta) - a(\vartheta')$, that is, only if all its cyclic sums vanish.*
- *They do not vanish in the checked instances, for $n = 1$ (from A_{even} alone) and for $n = 2$.*
- *So in these instances, at finite k_e, the exactly averaged loop is a non-reciprocal McKean–Vlasov dynamics. It is not the $W_{G^{-1}}$-gradient flow of any free energy. A different, μ-dependent metric is not excluded by this argument.*

Status. The frozen-measure part is proved. Non-reciprocity of the exact second variation is checked, in the instances $n = 1$ and $n = 2$ only, to 3% (asserted; 1–1.5% measured). The Stokes identity and the parity under orbit reversal hold to machine precision. The expansion is checked through its $O(1/k_e)$ convergence. Novelty (literature audit): no precedent found (targeted search); ingredients known (higher-order averaging; performative prediction [46]; semi-gradient TD [56, 58]).

Proof. (a) Write $e = r_\mu(x_m)/k_e + w$. Then w solves $\dot w = -k_e w + g(t,w)$ with $g = \nabla r_\mu(x_m)e - \dot r_\mu/k_e + O(|e|^2)$.

- The tube $\{\|e\| \le R\}$ is forward invariant once $k_e R > \sup_{\|e\|\le R}\|r_\mu(x_m + e)\|$.
- Inside the tube, solutions contract at rate $k_e - \mathrm{Lip}_R(r_\mu)$, which gives existence, uniqueness and attraction there. A contraction on T_0-periodic functions gives the expansion.
- The bounds on r_μ, ∇r_μ and $\nabla^2 r_\mu$ on the R-neighbourhood of the orbit are uniform in μ, by the saturation and $f \in C^2$.
- The global statement uses L_f in place of the local Lipschitz constant of f.

(b) Taylor-expand $e^\top\tilde\Phi(x_m + e;\vartheta) = e^\top\tilde\Phi(x_m;\vartheta) + e^\top\partial_x\tilde\Phi(x_m;\vartheta)e + O(|e|^3)$, and insert (a).

- The period averages of the time-symmetric terms are integrals against ρ.

- $(\nabla r\, r)^\top \tilde\Phi + r^\top(\partial_x \tilde\Phi) r = (r\cdot\nabla)(r^\top\tilde\Phi)$.
- ϑ-derivatives commute with the average.
- The remainders are uniform, because the x- and ϑ-derivatives of $\tilde\Phi$ of the orders needed are bounded.

(c) We have

$$\int r_\mu^\top \tilde\Phi_\vartheta \,\mathrm{d}\rho_\mu^{\mathrm{cl}} = \big\langle r_\mu(x_m+e_\mu)^\top \tilde\Phi(x_m+e_\mu;\vartheta)\big\rangle = \int r_\mu^\top \tilde\Phi_\vartheta \,\mathrm{d}\rho + \big\langle \nabla_x(r_\mu^\top\tilde\Phi_\vartheta)\cdot e_\mu\big\rangle + O(k_e^{-2}).$$

By (a) this equals $\int r_\mu^\top \tilde\Phi_\vartheta \,\mathrm{d}\rho + (1/k_e)U_{\mathrm{even}} + O(k_e^{-2})$.
(d) $\frac{\mathrm{d}}{\mathrm{d}t} r_\mu(x_m) = \frac{\mathrm{d}}{\mathrm{d}t} f(x_m) - \int \frac{\mathrm{d}}{\mathrm{d}t}\tilde\Phi(x_m;\vartheta')\,\mathrm{d}\mu(\vartheta')$.

- Antisymmetry: $K(\vartheta,\vartheta') + K(\vartheta',\vartheta) = \frac{1}{T_0}\oint \mathrm{d}\big(\tilde\Phi_\vartheta^\top \tilde\Phi_{\vartheta'}\big) = 0$.
- The area form is Stokes, or Green's theorem for $n=2$.
- For $n=1$, every 1-form on $\mathbb{R}$ is exact, so its loop integrals vanish.
- Reversal changes the sign of oriented loop integrals and preserves ρ.

(e) The second variation of U_{even} is

$$S_{\mathrm{even}}(\vartheta,\vartheta') = -\int \Big[(\tilde\Phi_{\vartheta'}\cdot\nabla)(r_\mu^\top\tilde\Phi_\vartheta) + (r_\mu\cdot\nabla)(\tilde\Phi_{\vartheta'}^\top\tilde\Phi_\vartheta)\Big]\,\mathrm{d}\rho,$$

whose second term is symmetric. The second variation of U_{odd} is K. A kernel's antisymmetric part A is a coboundary if and only if $A(\vartheta_0,\vartheta_1)+A(\vartheta_1,\vartheta_2)+A(\vartheta_2,\vartheta_0)=0$ for all triples.

The remainder of (b) is not differentiated in μ. Instead the argument uses a circulation around a small triangle of measures.

1. Fix $\vartheta_0,\vartheta_1,\vartheta_2$ and $\varepsilon\in(0,1)$. For (a,b) in $T_\varepsilon := \{a,b\ge 0,\ a+b\le\varepsilon\}$, let $\mu_{a,b} := (1-\varepsilon)\mu + (\varepsilon-a-b)\delta_{\vartheta_0} + a\delta_{\vartheta_1} + b\delta_{\vartheta_2}$.
2. The circulation $\oint_{\partial T_\varepsilon}\langle Q_{\mu_{a,b}}, \mathrm{d}\mu_{a,b}\rangle$ vanishes when Q_μ is a first variation plus a constant $c(\mu)$, because $\mathrm{d}\mu_{a,b}$ has zero mass. In particular it vanishes for $-(1/k_e)V_L$.
3. $U_{\mathrm{even}}+U_{\mathrm{odd}}$ is quadratic in μ, so its circulation equals $-(\varepsilon^2/2)\cdot\sum_{\mathrm{cyc}}(A_{\mathrm{even}}+2K)(\vartheta_i,\vartheta_{i+1})$ at the centroid of T_ε.
4. The $O(k_e^{-3})$ remainder of (b) contributes at most $6\varepsilon\cdot\sup|R| = O(\varepsilon k_e^{-3})$.
5. Suppose that, for all large k_e, P_μ were a first variation up to $O(k_e^{-3})$, with constants uniform in k_e. Then $(\varepsilon^2/(2k_e^2))|\mathrm{cyc}| \le C\varepsilon k_e^{-3}$, that is, $|\mathrm{cyc}|\le 2C/(\varepsilon k_e)$. The cyclic sum does not depend on k_e, so letting $k_e\to\infty$ at fixed ε gives $\mathrm{cyc}=0$.

This is checked numerically: for $\varepsilon=0.05$ and $n=2$ the circulation equals the centroid formula to all printed digits. □

Checked by: `tests/test_averaging.py` in the accompanying repository, using `src/core/averaging.py`; seven tests: four for Proposition H.7, one for Remark H.0, two for Remark H.7a. The checks of Proposition H.7 are listed in Table 13.

18.2 The oscillation correction and the orientation of the reference orbit

Remark H.7a (the full first-order correction, and a test in the real closed loop)**.** Proposition H.7 freezes the measure over a period. In the closed loop the parameters also oscillate within each period, by $O(1/k_e)$.

Table 13: Checks of Proposition H.7 (`tests/test_averaging.py`).

check	result
expansion: relative error of $k_e^2(P_\mu - P_0)$ against $U_{\text{even}} + U_{\text{odd}}$	halves when k_e doubles. At $k_e = 40 \to 80$ it goes $0.54 \to 0.25$ for $n = 1$, $0.038 \to 0.019$ for $n = 2$ counter-clockwise, and $0.022 \to 0.011$ clockwise. A Richardson step at $k_e = 80$ and 160 brings $n = 1$ within 3% (asserted)
Stokes: loop-integral K against the area integral over the enclosed ellipse	agreement to $7 \cdot 10^{-15}$ relative (asserted at 10^{-6}); K antisymmetric; $K \equiv 0$ for $n = 1$
parity under orbit reversal (same ρ)	U_{even} unchanged and U_{odd} flipped, to $2 \cdot 10^{-15}$ relative (asserted at 10^{-10})
non-reciprocity: the *exact* second variation S ($k_e = 80$) against the prediction	the full matrix $k_e^2 S + k_e S_0$ is within 5% of $S_{\text{even}} + K$. The cyclic sums of the antisymmetric part agree to 3%: $n = 1$ gives -0.0711 against -0.0704, all from the state shift; $n = 2$ gives -0.0471 against -0.0464 (state shift -0.089, circulation $+0.043$)

The parameter equation. In the scaling $\sigma = \sigma_0/k_e$, write

$$\dot{\Theta} = \frac{1}{k_e} H(\Theta, t) + \frac{1}{k_e^2} H_2(\Theta, t) + O(k_e^{-3}), \tag{18.8}$$
$$H_i(\Theta, t) := G \nabla_\vartheta \big[r_\Theta(x_m(t))^\top \tilde{\Phi}(x_m(t); \vartheta_i) \big] - \sigma_0(\vartheta_i - \vartheta^0).$$

Second-order averaging (Sanders–Verhulst–Murdock [51, §2.9]) gives the dynamics of the period mean Y:

$$\dot{Y} = \frac{1}{k_e} \langle H \rangle(Y) + \frac{1}{k_e^2} \Big[\langle H_2 \rangle(Y) + \big\langle D_\Theta H(Y, t) \cdot u(Y, t) \big\rangle \Big] + O(k_e^{-3}). \tag{18.9}$$

- u is the zero-mean antiderivative in t of $H - \langle H \rangle$.
- $\langle H_2 \rangle$ is the frozen-measure correction of Proposition H.7.
- This is formal. The averaging theorem gives it on finite slow-time horizons.
- `averaging.oscillation_correction` computes the oscillation term $b_{\text{osc}} := \langle D_\Theta H \cdot u \rangle$.

Three properties follow.

- **Parity and value.** The oscillation term b_{osc} is odd under reversal of the orbit (`test_oscillation_correction_is_odd_under_orbit_reversal`). Its value matches an independent one-period near-identity flow at $k_e = 80$: the test asserts 3%, and 0.7% is measured (`test_oscillation_correction_matches_a_one_period_flow`).
- **Size.** Relative to the frozen-measure terms it scales like $|G|T_0$, the parameter displacement per period times k_e. It is negligible when adaptation is slow within a period, which is the small-gain limit in the fixed-k_e discussion of Conjecture H.6. It is not negligible as $k_e \to \infty$ at fixed G. Both terms are $O(k_e^{-2})$ and uniformly bounded, so the order of the route to Conjecture H.6 is unchanged.
- **Non-reciprocity survives.** The only orientation-even part of the full correction is U_{even}, and its antisymmetric second variation is not a coboundary (Proposition H.7(e)). Suppose the full first-order dynamics were reciprocal for both orientations. Then the average of the two antisymmetric parts, A_{even}, would be a coboundary, which it is not. So the full first-order averaged dynamics is non-reciprocal for at least one orientation.

A prediction tested in the real closed loop (`exp_orbit_orientation.py` → `results/orbit_orientation/`).

- **The prediction.** Traverse the same elliptic reference both ways. The occupation measure is the same, so $(\mathfrak{A})$, and any functional of ρ, cannot distinguish the two runs. The first-order theory predicts that the learned kernels differ by $\Theta_+ - \Theta_- \approx -2J^{-1}(b_{\mathrm{odd}} + b_{\mathrm{osc}}) = O(1/k_e)$. Here J is the Jacobian of the leading averaged drift at the equilibrium.
- **The setting.** $n = 2$, $m = 6$, $G = \mathrm{diag}(4, 4, 1, 1, 1)$, $\sigma = 2/k_e$, no dither, and the same initial parameters for both runs. The runs converged to machine precision. The six kernels start spread out (kernel spread 1.05) and converge to a common configuration (spread at most $5.6 \cdot 10^{-17}$ at the end of every run), so the learned network is effectively a single kernel: the test probes the orientation effect at a collapsed equilibrium of the $m = 6$ system.

Table 14: Orientation dependence of the learned kernels in the real closed loop (`results/orbit_orientation/table_sigma2_slow16.md`). The size ratio is measured over predicted.

		frozen measure only (H.7)		full (H.7 + oscillation)		full
k_e	measured $\|\Theta_+ - \Theta_-\|$	size ratio	cosine	size ratio	cosine	vector error
10	$5.7 \cdot 10^{-3}$	2.67	0.76	0.94	0.93	0.36
20	$2.8 \cdot 10^{-3}$	2.41	0.83	0.97	0.98	0.20
40	$1.4 \cdot 10^{-3}$	2.29	0.86	0.97	0.994	0.11

- The measured difference scales like $k_e^{-1.01}$.
- At $k_e = 40$ the full first-order theory predicts it to within 3% in size and 11% as a vector (cosine 0.994).
- The vector error halves with each doubling of k_e: 36%, 20% and 11% at $k_e = 10$, 20 and 40.
- The frozen-measure terms alone account for less than half of it.
- A first attempt with a weaker leak ($\sigma = 0.5/k_e$) and a shorter horizon had not converged: its residual drift was a third of the difference. It is not used.

The direction in which the reference orbit is traversed is therefore a property of the excitation that the learned kernels retain, at order $1/k_e$. The first-order theory predicts its effect.

Status. The second-order averaged equation (18.9) is formal; the averaging theorem gives it on finite slow-time horizons. The oscillation term is checked against an independent one-period near-identity flow (within 3% at $k_e = 80$) and is odd under orbit reversal (to 10^{-6}). The full first-order theory predicts the orientation dependence of the learned kernels in the real closed loop to 3% in size and 11% as a vector (cosine 0.994, $k_e = 40$).

Checked by: `tests/test_averaging.py` (the two tests named above); `exp_orbit_orientation.py` → `results/orbit_orientation/table_sigma2_slow16.md`.

18.3 Input-to-state stability of $(\mathfrak{A})$ and the route to Conjecture H.6

Proposition H.8 (input-to-state stability of $(\mathfrak{A})$)**.** *Assume $\sigma, \tau > 0$ and the setting of Theorem H. Let μ_t solve the perturbed equation*

$$\partial_t \mu = \nabla \cdot \big(\mu(G \nabla V[\mu] - b_t)\big) + \tau \nabla \cdot (G \nabla \mu) \tag{18.10}$$

from $\mu_0 \in \mathcal{P}_2$ with $F_\tau(\mu_0) < \infty$. Here b_t is a velocity perturbation with $\beta(t)^2 := \int \|b_t\|_{G^{-1}}^2 \,\mathrm{d}\mu_t \le \bar\beta^2$. Assume the chain rule of [9, Lemma 2.2] holds along this solution:

$$\frac{\mathrm{d}}{\mathrm{d}t} F_\tau(\mu_t) = \int \langle \nabla(\delta F_\tau/\delta\mu)(\mu_t),\, v_t\rangle \,\mathrm{d}\mu_t,$$

in whitened coordinates, with v_t the velocity. It is expected for smooth bounded perturbations (parabolic regularity and the entropy-dissipation estimate), but it is not verified here. Then

$$\frac{\mathrm{d}}{\mathrm{d}t}\big(F_\tau(\mu_t) - F^*\big) \le -\tau\rho_\tau\big(F_\tau(\mu_t) - F^*\big) + \tfrac{1}{2}\beta(t)^2, \tag{18.11}$$

$$F_\tau(\mu_t) - F^* \le e^{-\tau\rho_\tau t}\big(F_\tau(\mu_0) - F^*\big) + \frac{\bar\beta^2}{2\tau\rho_\tau}, \tag{18.12}$$

$$\limsup_{t\to\infty} W_{G^{-1}}(\mu_t, \mu_\tau^*) \le \frac{\bar\beta}{\tau\rho_\tau}. \tag{18.13}$$

Status. Proved, assuming the chain rule for the perturbed flow. A proof only; not checked in the repository. Novelty (literature audit): elementary given [9].

Proof. Work in $\zeta = G^{-1/2}\vartheta$, and let $\xi_t := \nabla(\delta F_\tau/\delta\mu)(\mu_t) = \nabla\tilde V[\mu_t] + \tau\nabla\log\mu_t$. The velocity is $-\xi_t + \tilde b_t$, with $\|\tilde b_t\| = \|b_t\|_{G^{-1}}$.

1. By the chain rule and Young's inequality,

$$\frac{\mathrm{d}}{\mathrm{d}t} F_\tau(\mu_t) = -\int \|\xi_t\|^2 \,\mathrm{d}\mu_t + \int \langle \xi_t,\, \tilde b_t\rangle \,\mathrm{d}\mu_t \le -\frac{1}{2}\int \|\xi_t\|^2 \,\mathrm{d}\mu_t + \frac{1}{2}\beta^2.$$

2. With $\hat\mu_t \propto e^{-\tilde V[\mu_t]/\tau}$,

$$\int \|\xi_t\|^2 \,\mathrm{d}\mu_t = \tau^2 I(\mu_t|\hat\mu_t) \ge 2\rho_\tau\tau^2 H(\mu_t|\hat\mu_t) \ge 2\rho_\tau\tau\big(F_\tau(\mu_t) - F^*\big),$$

   by the LSI of Theorem H(ii) and the entropy sandwich [9, Lemma 3.4].

3. Grönwall gives the first two lines.

4. For the third, $F_\tau(\mu) - F^* \ge \tau H(\mu|\mu_\tau^*)$ [9, Lemma 3.4]. μ_τ^* satisfies LSI(ρ_τ), hence Talagrand's inequality $W^2 \le (2/\rho_\tau)H(\cdot|\mu_\tau^*)$ (Otto–Villani [42]). So $W^2 \le (2/(\tau\rho_\tau))(F_\tau - F^*)$, and the limsup follows from the second line. □

The gain $1/(\tau\rho_\tau) \le e^{\alpha^*/\tau}/\sigma$ is exponentially large in $1/\tau$, like the inverse of the rate.

Route to Conjecture H.6. In the scaling of Conjecture H.6:

- $\tau\rho_\tau \ge (\sigma_0/k_e)\cdot e^{-2K_\Phi(F_\rho+K_\Phi)/\tau_0}$, so $1/(\tau\rho_\tau) = O(k_e)$.
  - In this scaling the whitened proximal Gibbs measures do not depend on k_e, so $\tau\rho_\tau = \Theta(1/k_e)$.
  - At the experiment's $\tau_0 = 0.04$ the constant is e^{1049}.
- Define $b := (1/k_e^2)\big[G\nabla(U_{\text{even}} + U_{\text{odd}}) + b_{\text{osc}}\big] + O(k_e^{-3})$. The first part is the frozen-measure correction of Proposition H.7; the second is the oscillation term of Remark H.7a. Both are bounded uniformly in μ and ϑ, so $\bar\beta = O(k_e^{-2})$.
- Proposition H.8 then confines every solution of $(\mathfrak{A}) + b$ to an asymptotic W-ball around μ_τ^* of radius $O(1/k_e)\cdot e^{2K_\Phi(F_\rho+K_\Phi)/\tau_0}/\sigma_0$. This needs $F_\tau(\mu_0) < \infty$, or a $\mathcal{P}_2$ start, after the Girsanov regularisation of Theorem H(ii), which applies because b is bounded.

Three things are still missing for Conjecture H.6:

1. control of the frozen-measure approximation along the moving μ_t, that is, a near-identity averaging transformation for the stochastic McKean–Vlasov loop, uniformly in time;
2. the mean-field limit of (L_τ), since Theorem D covers only (L);
3. the chain rule that Proposition H.8 assumes for the perturbed flow.

18.4 Evidence from the real closed loop

The evidence comes from the real closed loop with $m = 32$ and $n = 1$, not from the averaged system.

- **Dither is necessary.** Without dither the collapsed start is trapped forever. Its loss stays at 0.166, against 0.007 from the spread start (`results/langevin_tradeoff`, 1500 s).
- **Equalisation at moderate temperature.** At $\sigma = 0.02$ and $\tau = 0.004$ (3 seeds, `results/langevin_global/table_sigma0.02_tau0.004.md`):
  - collapsed and spread starts reach the same loss (ratio 1.03) and tracking RMS (0.0465 against 0.0452);
  - their final 32-particle configurations, driven by the same noise realisation, are 4.2× closer in W_2 over the physical coordinates (a, c, s) than without dither (0.40 against 1.69);
  - a numerical check shows that two independent 32-point samples of μ_τ^* are typically about 0.6 apart, so 0.40 is at the sampling floor;
  - the common loss of about 0.11 matches the grid value $\bar{L}(\mu_\tau^*) = 0.11118$. It is dominated by the regularisation bias.
- **Metastability at low temperature.** At $\sigma = 0.002$ and $\tau = 5 \cdot 10^{-4}$ (table for these values in `results/langevin_global/`), dither frees the collapsed start ($0.166 \to 0.048$). But the starts have not equalised within 900 s (loss ratio 5.2). The spread start's $9.2 \cdot 10^{-3}$ is close to the regularised optimum $\bar{L}(\mu_\tau^*) = 8.5 \cdot 10^{-3}$.
- **Limits of the evidence.**
  - The Langevin runs have $n = 1$. By Proposition H.7, the averaged dynamics is already non-reciprocal for $n = 1$, through the state shift.
  - The only $n \geq 2$ closed-loop runs are the deterministic orientation runs of Remark H.7a. They confirm the first-order theory quantitatively. No $n \geq 2$ run with dither has been made.
  - The runs use fixed k_e, not the scaling of Conjecture H.6.
  - All Langevin runs use $k_e = 10$, below the Tikhonov scale of Remark H.0 for this network ($(F_\rho + K_\Phi)/s_{\min} \approx 35$). The measured frozen-$\mu$ drift error at $k_e = 10$ is 4.2% ($n = 1$), so the condition is conservative, but the agreement with ($\mathfrak{A}$) is empirical.
  - So the evidence supports Conjecture H.6 qualitatively, but the equalisation time can exceed practical horizons at low τ.

Part III
Evidence and outlook

19 Numerical evidence

Every number in this section is produced by the named script from raw files, and no table is edited by hand. The scripts are in the directory `experiments/` of the accompanying repository and write their output to subdirectories of `results/`; "`script.py` → `dir/`" below names a script and its output directory or file.

- The test files `tests/test_transport.py`, `tests/test_langevin.py` and `tests/test_averaging.py` check the inequalities at every sample of simulated trajectories, or on sampled states.
- They are run with `venv\Scripts\python -m pytest tests`. The current number of tests is given in the repository's `README.md`.

19.1 Theorem B: the label-free Lyapunov function

Script: `exp_transport_permutation.py` → `transport_permutation/`.

Each network starts as a random relabelling of the ideal configuration plus noise 0.05, with mean-field gains and 3 seeds per m. "Excess" is $|(1/m)\sum \text{Taylor remainders}|$, the part that actually enters $\dot{V}$. Table 15 and Figure 4 show the results.

Table 15: Theorem B, permuted-ideal start, mean-field gains; means over 3 seeds. The two bounds are the same bound, evaluated with the matched (OT) error and with the labelled error, along the same trajectory.

m	labelled error ($t=0$)	OT error W ($t=0$)	RMS$\|e\|$	actual excess	bound, matched error	same bound, labelled error
8	2.675	0.1070	0.0113	$1.96 \cdot 10^{-5}$	$8.27 \cdot 10^{-2}$	2.93
16	2.532	0.1044	0.0065	$9.58 \cdot 10^{-6}$	$4.37 \cdot 10^{-2}$	1.86
32	2.684	0.1122	0.0079	$2.69 \cdot 10^{-5}$	$7.07 \cdot 10^{-2}$	2.11
64	2.764	0.1136	0.0094	$1.23 \cdot 10^{-5}$	$6.80 \cdot 10^{-2}$	2.43

- The bound is at least the actual excess at every sample of every run.
- The matched error is about 1/25 of the labelled one.
- The bound is 30–45× smaller when evaluated with the matched error.
- Nothing grows with m.
- The bounds remain far above the actual excess, because the Hessian supremum is global.

Corollary B.5 along (L_σ)**.** This is checked by the test `test_corollary_B5_fails_along_lsigma_and_its_leak_variant_holds`. The test asserts every hypothesis of C(iv) and measures $\sup W \le \bar{\eta}$. There, the plain B.5 bound is exceeded by more than 3×, while the leak variant with $\sigma_m b^2/4$ holds.

19.2 Theorem C: OT σ-modification

Scripts: `exp_ot_sigma_mod.py` and `exp_sigma_mod_transient.py` → `ot_sigma_mod/`.




Figure 4: Theorem B, permuted-ideal start. Left: labelled parameter error and OT error $W(\mu_t, \mu^*)$ for $m = 16$. Centre: the actual excess disturbance that enters $\dot{V}$, the OT bound of Theorem B and the labelled bound, for $m = 16$. Right: the mean bound with identity labelling, the mean bound with OT matching and the mean actual excess, against m.

The two starts share plant, prior, geometry and x_0. They differ only in particle labels (Table 16, Figure 5). Table 17 shows the reshuffling transient from the relabelled start (5 seeds).

Table 16: Theorem C: OT σ-modification against labelled σ-modification (oracle plant, $m = 6$, 5 seeds, $\sigma_m = 0.3$). RMS$\|e\|$ is taken over 20–40 s and reported as mean $\pm$ 95% confidence interval over seeds; W_{end} is the OT error at the end.

start	mechanism	RMS$\|e\|$ (20–40 s)	W_{end}	labelled error at end
aligned	projection	0.0051 ± 0.0010	0.2953	0.2953
aligned	labelled σ-mod.	0.0091 ± 0.0018	0.3040	0.3040
aligned	OT σ-mod.	0.0091 ± 0.0018	0.3040	0.3040
relabelled	projection	0.0051 ± 0.0010	0.2953	0.2953
relabelled	labelled σ-mod.	0.0091 ± 0.0018	0.3040	**2.0590**
relabelled	OT σ-mod.	0.0091 ± 0.0018	0.3040	0.3040

Table 17: Reshuffling transient from the relabelled start ($m = 6$, 5 seeds, $\sigma_m = 0.3$): RMS$\|e\|$ over time windows, and the particle path length.

leak	$[0, 5]$ s	$[5, 10]$ s	$[10, 20]$ s	$[20, 40]$ s	particle path length
labelled	0.0771	0.0086	0.0093	0.0091	2.2585
OT	0.0743	0.0072	0.0093	0.0091	0.3917

- The labelled leak's longer path costs only about 4% of early RMS$\|e\|$. Theorem F does not price leak-driven transport (Proposition C.5).
- The C(iv) instance in the tests satisfies every hypothesis, asserted. V_{OT} stays below the envelope $e^{-\alpha t}V(0) + \beta/\alpha$ at every sample and contracts by more than 5×.
- The C(ii) test asserts $C \le b^2 + 4\eta R$, $\sqrt{C} \le b + 2\eta$, and $C = b^2$ under both gap conditions, on 300 configurations.

19.3 Theorem D: convergence in m

Script: `exp_mean_field_m.py` → `mean_field_m/`. Table 18 and Figure 6 show the results.

This is a consistency check in m, not a check of the constant.




Figure 5: Theorem C: $\|e(t)\|$ (median of seeds) for projection, labelled σ-modification and OT σ-modification, from the aligned start (left) and from the relabelled start (right).

Table 18: Theorem D: mean-field consistency in m (Van der Pol plant, mean-field gains, smooth initial measure quantised at m points). The ratio is $W(\mu_T^m, \mu_T^{2m})/W(\mu_0^m, \mu_0^{2m})$.

m	$\sup_t \|e^m - e^{2m}\|$	$W(\mu_0^m, \mu_0^{2m})$	$W(\mu_T^m, \mu_T^{2m})$	ratio
16 vs 32	0.0046	0.2179	0.3658	1.68
32 vs 64	0.0009	0.1091	0.1788	1.64
64 vs 128	0.0001	0.0545	0.0920	1.69
128 vs 256	0.0000	0.0273	0.0459	1.68

- The guaranteed factor $e^{\Lambda T}$ is at least e^{496} on this box.
- The Van der Pol plant fails the tube condition of Theorem D(b), so the run is covered by the localised D(d).

19.4 Theorem E: label-free convergence under excitation

Script: `exp_pe_convergence.py` → `pe_convergence/`. Table 19 and Figure 7 show the results.

Table 19: Theorem E: tangential excitation and label-free parameter convergence ($m = 3$, $n = 2$, permuted start, $T = 200\,\mathrm{s}$). $\lambda_{\min}$ is that of the excitation Gramian.

reference	seed	$\lambda_{\min}$, first 60 s window	min over sliding 60 s windows	$W(T)/W(0)$	fitted rate of W, $t \geq 100\,\mathrm{s}$ (1/s)	labelled(T) / labelled(0)
rich (6 tones)	0	$8.60 \cdot 10^{-3}$	$4.12 \cdot 10^{-3}$	0.544	$2.31 \cdot 10^{-3}$	0.992
rich (6 tones)	1	$5.52 \cdot 10^{-3}$	$1.02 \cdot 10^{-3}$	0.409	$2.32 \cdot 10^{-3}$	0.962
rich (6 tones)	2	$1.08 \cdot 10^{-3}$	$5.91 \cdot 10^{-4}$	0.369	$3.07 \cdot 10^{-3}$	1.007
single sine	0	$2.54 \cdot 10^{-2}$	$2.41 \cdot 10^{-2}$	0.274	$4.05 \cdot 10^{-3}$	0.988
single sine	1	$2.96 \cdot 10^{-2}$	$2.71 \cdot 10^{-2}$	0.314	$4.26 \cdot 10^{-3}$	0.967
single sine	2	$6.74 \cdot 10^{-3}$	$6.61 \cdot 10^{-3}$	0.226	$2.91 \cdot 10^{-3}$	1.008

- The matched error decays in all six runs. The labelled error stays within 4% of its initial value.
- The rank correlation between the uniform excitation level and the fitted rate is 0.60 over the six runs. The PE level is a hypothesis of Theorem E, not a rate predictor (Remark E.1).




Figure 6: Theorem D: tracking error $\|e(t)\|$ of the closed loop (Van der Pol plant, mean-field gains) for $m = 16$, 32, 64, 128 and 256.

- The proof's linearisation is checked exactly: the Jacobian equals $A(t)$, and $QA + A^\top Q = \mathrm{diag}(-k_e I, 0)$.




Figure 7: Theorem E: OT error $W(\mu_t, \mu^*)$ and identity-labelled error for the three seeds, with the rich reference (left) and the single-sine reference (right).

19.5 Theorem F: transport cost

Theorem F is checked by tests.

- Along (L), the particle path length (which is at least the Wasserstein length) stays below $\lambda_G^{1/2} L_1 \int \|e\|$ at every sample.
- Along (L_σ) with the OT leak, $W(\mu_{t_1}, \mu_{t_0}) \le \int (\lambda_G^{1/2} L_1 \|e\| + \sigma_m W(\mu_r, \mu^0))$ on a grid of pairs.
- Lemma P.1 holds on 2000 adversarial states, and a tight instance of F(i) reaches ratio ≥ 0.99.

19.6 Proposition G: slow geometry

Scripts: `exp_variable_projection.py` and `exp_slow_manifold_slope.py`, both → `variable_projection/`.

In Table 20, ranges are along the run (5 samples), and "a / b" means first-quarter / last-quarter medians. Velocities are centred differences of period means.

Table 20: Proposition G: the two configurations, `violating` ($m = 4$, $k_e = 20$, $g_a = 2$, $\delta = 0.1$, $T = 800\,\mathrm{s}$) and `satisfying` ($m = 2$, $k_e = 20$, $g_a = 13$, $\delta = 0.004$, $T = 3000\,\mathrm{s}$). In the rows for L^* and $\bar{L}$, "$x \to y$" is first sample → last sample.

quantity	violating ($m = 4$)	satisfying ($m = 2$)
(H2a) ω_{ref} / joint fastest	15.3–25.6	6.53–7.27
(H2b) slowest weight / geometry	0.0993–0.231	78.8–87.2
(H2c) S	5.44–25	0.00601–0.00602
G(a) direct check (cosine, G^{-1} metric)	0.9994 / 0.9985	0.9848 / 0.9624 (geometry block 0.9992 / 0.9982)
G(b) cosine	−0.118 / 0.385	0.999 / 0.999
G(b) size ratio	4.207 / 0.472	0.969 / 0.970
L^*	**increases** ($3.8136 \cdot 10^{-3} \to 5.8866 \cdot 10^{-3}$)	decreases ($1.1987 \cdot 10^{-2} \to 4.1073 \cdot 10^{-3}$)
$\bar{L}$ at period-mean weights	non-increasing ($0.21892 \to 0.0080127$)	non-increasing

Slope counterexample. Table 21 (`table_slope.md`). Here $m = 1$, $k_e = 20$, gains (g_a, δ, δ), and (H2b) holds in every row. The closed loop and the joint flow start on the slow manifold of (b′).

Table 21: Proposition G(b): closed loop against the averaged predictions. "closed / (b)" compares the closed loop with plain variable projection (VarPro) G(b), and "closed / (b′)" with the induced-metric flow G(b′); size is the ratio of displacement norms.

case	(H2b) ratio	S	closed / (b)		closed / (b′)		(a) / (b′)	closed / (a)
			size	cosine	size	cosine		
$\delta = 0.47$, $g_a = 20$	81–178	2–2.81	0.422	0.9303†	1.121	0.9986	1.015	1.104
$\delta = 0.47/4$, $g_a = 5$ (all gains ÷4)	81–178	2–2.81	0.401	0.9156	1.065	0.9999	1.015	1.049
$\delta = 0.047$, $g_a = 20$	654–1779	0.2–0.319	0.850	0.9986	1.047	1.0000	1.003	1.044
$\delta = 0.0047$, $g_a = 20$	13162–17790	0.02–0.0229	1.023	1.0000	1.044	1.0000	1.000	1.044

†The centre moves the opposite way.

- (a) / (b′) isolates the $O(1/R)$ error of the induced-metric flow.
- closed / (a) isolates the averaging error, which shrinks with the gains, plus a 4.4% $O(1/k_e)$ error-channel factor.

What the tables show:

- G(b) holds in the satisfying configuration, where (H2b) and $S \ll 1$ both hold. The violating configuration fails both.
- The slope rows show that (H2b) alone is not sufficient.

- The induced-metric flow G(b′) tracks the closed loop at every S.
- In the satisfying configuration, the G(a) weight block is dominated by the $O(1/k_e)$ shift of the weight equilibrium: cosine -0.21 / 0.50, carrying 92–95% of the residual. The geometry block matches.




Figure 8: Proposition G, `violating` configuration. Left: the G(b) cosine between the actual geometry velocity and the VarPro direction $-(m\delta/k_e)\Gamma\nabla L^*$, along the run. Centre: the profile loss L^* and $\|\nabla L^*\|$. Right: the centres $c_i(t)$.




Figure 9: Proposition G, `satisfying` configuration; panels as in Figure 8.

19.7 Theorem H: Langevin adaptive control

Scripts: `exp_langevin_global.py` → `langevin_global/`, `exp_langevin_tradeoff.py` → `langevin_tradeoff/`, and `exp_gibbs_reference.py` → `langevin_global/gibbs_reference.md`.

The setting is the real closed loop with $m = 32$, a scalar plant and a periodic reference.

- The unregularised offline optimum of the occupation loss is about $7.6 \cdot 10^{-6}$ (best of 5 L-BFGS restarts, an upper bound). One run gave $7.0 \cdot 10^{-6}$, because of floating-point summation order.
- The regularised optimum $\bar{L}(\mu_\tau^*)$ is 0.11118 at $(\sigma, \tau) = (0.02, 0.004)$ and 0.00850 at $(0.002, 5 \cdot 10^{-4})$, grid-converged (`exp_gibbs_reference.py` → `langevin_global/gibbs_reference.md`).
- In the scaling $\sigma = 0.2/k_e$, $\tau = 0.04/k_e$ of Conjecture H.6, $\bar{L}(\mu_\tau^*) = 0.11118$ for every k_e. The exactly averaged stationary state gives $\bar{L} = 0.11415$, 0.11148 and 0.11101 at $k_e = 10$, 20 and 40. The $L^2(\rho)$ distance between the two learned functions is $6.81 \cdot 10^{-3}$, $1.43 \cdot 10^{-3}$ and $9.7 \cdot 10^{-4}$.

Speed–bias trade-off. Table 22 and Figure 10 ($\sigma = 0.002$, 2 seeds, `langevin_tradeoff/table.md`).

Table 22: Theorem H: speed–bias trade-off of parameter dither ($m = 32$, $\sigma = 0.002$, $k_e = 10$, $T = 1500\,\mathrm{s}$, 2 seeds). The escape time is the first sample at which the collapsed run's loss is at most twice the spread run's final loss (same seed); per-seed values are listed, and ∞ means not within the horizon. The losses are the final values of $\bar{L}$.

τ	escape of the collapsed start (s)	loss, collapsed	loss, spread
0	∞, ∞	0.166	0.00696
0.0005	∞, 1050	0.0249	0.0109
0.001	1480, 1005	0.0216	0.0148
0.002	1115, 800	0.0288	0.0233
0.004	970, 430	0.0392	0.0362
0.008	390, 295	0.0603	0.0653
annealed (hot start)	990, 530	0.0317	0.0320

- **Fit.** The seed-averaged escape time fits $A + B\log(1/\tau)$ with an RMS relative residual of 0.045, against 0.253 for an Arrhenius fit. On the eight per-seed values both laws fit poorly (0.274 and 0.330). The data do not identify the escape mechanism, and the "saddle" reading of an earlier version of this analysis is withdrawn (Remark H.2).
- **Caveats.** The threshold grows with τ, and the censored $\tau = 5 \cdot 10^{-4}$ point is excluded from the fit.
- **Guaranteed rate.** The Holley–Stroock [27] rate $2\tau\rho_\tau$ is at least $e^{-8401}\,\mathrm{s}^{-1}$ at $(0.002, 5 \cdot 10^{-4})$ and at least $e^{-1053}\,\mathrm{s}^{-1}$ at $(0.02, 0.004)$.





Figure 10: Theorem H, speed–bias trade-off: $\bar{L}(\mu_t)$ of the collapsed start (red) and of the spread start (blue), 2 seeds, for $\tau = 0$, 0.0005, 0.001, 0.002, 0.004, 0.008 and the annealed (hot-start) schedule.

Two fixed settings. Three seeds (`langevin_global/table_*.md`; Figure 11).


- $\sigma = 0.02$, $\tau = 0.004$**.** Both starts equalise: the loss ratio is 1.03, and the final measures are 4.2× closer in W than without dither. The common loss is about 0.11, which matches the grid value $\bar{L}(\mu_\tau^*) = 0.111$. It is dominated by the regularisation bias.
- $\sigma = 0.002$, $\tau = 5 \cdot 10^{-4}$**.** Dither frees the collapsed start (0.166 → 0.0480), but the starts do not equalise within 900 s (loss ratio 5.23).

19.8 Remark H.0 and Proposition H.7: the exactly averaged closed loop

These are checked by tests. Table 23 gives the relative error of the frozen-μ averaged drift against ($\mathfrak{A}$), and Table 24 the structure of the first correction (`test_averaging.py`).



(a) $\sigma = 0.02$, $\tau = 0.004$



(b) $\sigma = 0.002$, $\tau = 5 \cdot 10^{-4}$

Figure 11: Theorem H, global kernel placement ($m = 32$, scalar plant, $k_e = 10$, 3 seeds): the occupation loss $\bar{L}(\mu_t)$ of the collapsed and spread starts without dither, with constant dither and with annealed dither.

Table 23: Remark H.0: relative error of the frozen-μ averaged drift against ($\mathfrak{A}$).

instance	$k_e = 10$	20	40	80	$k_e \times$ error
$n = 1$ (`test_langevin.py`)	4.2%	2.0%	0.91%	0.48%	≈ 0.4
$n = 2$, ellipse (`test_averaging.py`)	43%	21%	10%	5.1%	≈ 4.2

Table 24: Proposition H.7: structure of the first correction (`test_averaging.py`).

check	$n = 1$	$n = 2$ (ellipse)
expansion error of $k_e^2(P_\mu - P_0)$ against $U_{\text{even}} + U_{\text{odd}}$, $k_e = 40 \to 80$	$0.54 \to 0.25$	$0.038 \to 0.019$
circulation kernel K	$\equiv 0$ (asserted at 10^{-12})	antisymmetric; equals the enclosed-area integral to $7 \cdot 10^{-15}$ (asserted at 10^{-6})
orbit reversal (same ρ)	$U_{\text{odd}} \equiv 0$	U_{even} unchanged, U_{odd} flipped, to $2 \cdot 10^{-15}$ (asserted at 10^{-10})
$k_e^2 \cdot$ cyclic sum of the antisymmetric second variation, exact ($k_e = 80$) against predicted (asserted at 3%; the full matrix at 5%)	-0.0711 against -0.0704	-0.0471 against -0.0464

19.9 Remark H.7a: the orientation of the reference orbit

Script: `exp_orbit_orientation.py` $\to$ `orbit_orientation/`.

The setup is deterministic: $n = 2$, $m = 6$, $G = \text{diag}(4, 4, 1, 1, 1)$, $\sigma = 2/k_e$, no dither. The same elliptic reference is traversed both ways, so the occupation measure is the same, and so is

the initial configuration. The runs converged to machine precision: the residual drift over the last 10 periods is about 10^{-16} (between $1.0 \cdot 10^{-16}$ and $5.0 \cdot 10^{-16}$). The six kernels converge to a common configuration (kernel spread 1.05 at the start, at most $5.6 \cdot 10^{-17}$ at the end of every run), so the learned network is effectively a single kernel and the test probes the effect at a collapsed equilibrium. Predictions are linear responses at the (+) equilibrium (Table 25).

Table 25: Remark H.7a: orientation dependence of the learned kernels in the real closed loop. Size ratios are measured / predicted.

		frozen-measure prediction (H.7)		full prediction (H.7 + oscillation term)	
k_e	measured $\|\Theta_+ - \Theta_-\|$	size ratio	cosine	size ratio	cosine
10	$5.729 \cdot 10^{-3}$	2.670	0.7577	0.937	0.9314
20	$2.843 \cdot 10^{-3}$	2.414	0.8302	0.965	0.9809
40	$1.403 \cdot 10^{-3}$	2.294	0.8631	0.969	0.9944

- The vector error $\|\text{measured} - \text{predicted}\|/\|\text{predicted}\|$ of the full prediction is 0.364, 0.195 and 0.109 at $k_e = 10$, 20 and 40.
- The measured difference scales like $k_e^{-1.01}$, against a predicted -1.
- The full first-order theory accounts for it. The frozen-measure terms alone account for less than half.
- A first attempt with $\sigma = 0.5/k_e$ and half the horizon had not converged (residual drift about a third of the difference). It was discarded and rerun.

20 Research program

Each branch states what is proved, what is open, and the first concrete step.

1. **Infinite-horizon averaging for McKean–Vlasov closed loops (Conjecture H.6).**
   - *Proved:* the averaged system converges globally and exponentially (H(ii)); the exact first correction at frozen measure (H.7); input-to-state stability of the averaged system (H.8).
   - *Open:* a near-identity averaging transformation for the stochastic McKean–Vlasov loop that keeps the frozen-measure correction valid along the moving measure, uniformly in time.
   - *First step:* the deterministic case ($\tau = 0$, $\sigma > 0$) at fixed m, where the needed averaging theorem is classical (Sanders–Verhulst–Murdock [51, Ch. 5], "averaging under attraction").
2. **Non-reciprocal mean-field dynamics from closed-loop learning (new; Proposition H.7).**
   - *Proved at a frozen measure:* the averaged adaptive dynamics is a semi-gradient flow plus a lag term (Proposition H.7). Along the real loop a second-order averaging term is added (Remark H.7a).
   - Its interaction has two antisymmetric parts:
     - the performative defect A_{even} of the state shift, which is present for every n and is the only part for $n = 1$;
     - the circulation kernel K, an integral over the region that the reference orbit encloses ($n \geq 2$).

- Non-reciprocity is checked in the instances tested, not proved in general.
- *Open questions:*
  - Are stationary states unique beyond the perturbative regime?
  - Can circulation be *designed*, choosing the orbit's orientation and area, to break symmetric traps or to speed escape from the collapsed configuration?
  - Is there a Lyapunov functional other than F_τ for the non-reciprocal flow?
- *Context:* the closest analogues are performative prediction [46], semi-gradient TD learning [58], [56] and non-reciprocal phase transitions [20].
- *Done (Remark H.7a):* deterministic $n = 2$ closed-loop runs with the reference traversed both ways (same ρ). The full first-order theory predicts the $O(1/k_e)$ difference of the learned kernels to 3% in size and 11% as a vector.
- *Next steps:* the same test with dither, where the stationary state is a measure; and the design of the orbit's orientation and area as a control input.

3. **Uniform-in-time propagation of chaos with a plant in the loop.**
   - Theorem D is finite-horizon. Uniform-in-time results exist for mean-field Langevin dynamics [6] and for shallow networks [23].
   - *Open:* the closed loop, where the plant couples all particles through e, and where (by H.7) the averaged interaction is not reciprocal.
4. **Geometry of two-time-scale adaptation (Proposition G(b′)).**
   - *Proved formally, and checked:* the slow geometry follows the gradient flow of the profile loss in the metric induced on the slow manifold, and the slope S measures the departure from plain variable projection.
   - *Open:* a rigorous long-horizon statement; the $O(1/R)$ corrections, which involve D^2a^* and $\partial_\xi M$; gain designs that shape the induced metric.
5. **Noiseless global convergence.**
   - *Open:* a closed-loop separation condition under which the noiseless flow avoids spurious stationary points, despite the failure of the Chizat–Bach [8] boundary conditions. Candidates are a compactified width coordinate and reflecting boundaries (Remark H.3).
6. **Unbalanced transport, meaning online structure adaptation.**
   - Treat the weights as masses (Hellinger–Kantorovich / Wasserstein–Fisher–Rao) to obtain birth and death of kernels with Lyapunov guarantees. The conic particle descent of [10] and [14] is the static analogue.
7. **Excitation theory in measure space.**
   - *Open:* the continuum form of (PE); the trajectory kernel

     $$K_T(\vartheta, \vartheta') = \int \Phi(x(s); \vartheta)\Phi(x(s); \vartheta')\, \mathrm{d}s;$$

     identifiability of measures up to symmetry; the relation to [26]. Remark E.1 shows that the PE level alone does not predict the rate. A quantitative, uniform UES estimate for the skew-symmetric class is a first target.
8. **Transport-regularised adaptive laws.**
   - Wasserstein proximal (JKO) discrete-time adaptive laws.

- The OT leak toward priors with a different number of atoms, using optimal plans with barycentric projection.
- Transfer: transporting a learned measure to a new plant.

9. **Quantisation and the curse of dimensionality (Corollary B.6).**
   - *Open:* second-order (centroid) quantisation rates for the ideal measure on the lifted space; approximation rates that do not go through W (Maurey-type); adaptive laws that track optimal quantisers (the centroidal Voronoi tessellation connection [43]).
10. **Symmetric systems beyond kernels.** The OT quotient applies verbatim to swarms of identical agents, mixtures and multi-head models ([15] and [28] are the control-side precedents).
11. **Deep approximators.** Hierarchical (multi-layer) mean-field measures, with a layer-wise OT Lyapunov function.

A Verification map

Table 26 lists the tests and experiments of the accompanying repository that check each statement; statements not checked are listed after it. Test files are abbreviated: `T` is `tests/test_transport.py`, `Lg` is `tests/test_langevin.py`, and `Av` is `tests/test_averaging.py`.

Table 26: Verification map.

statement	what is checked	where
constants (1) of Section 9, closed forms	$\nabla^2_{(c,s)}\varphi \le H_\varphi$ by brute force	`T::test_gaussian_constants_are_suprema`
Lemma P.1	$\|\mathrm{Proj}_K(\vartheta, Gy)\|_{G^{-1}} \le \|y\|_G$ on 2000 adversarial states, with equality when inactive; a tight instance of F(i) with ratio ≥ 0.99	`T::test_lemma_P1_projection_is_nonexpansive_and_tight_velocity_bound`
A(i) raw rates	lifted raw rates equal classical raw rates under the gain map (10^{-10})	`T::test_identification_with_classical_law`
A(i) projected rates, only-if	per-kernel projection with rescaled layer matches to 10^{-10} with active projections; an unrescaled layer does not; five joint balls are each distinguished (two-case construction: four active against an inactive per-kernel field, one inactive against an active one)	`T::test_identification_projected_rates_per_kernel`
B with constant ideal and $\varepsilon = 0$	the identity to floating-point precision; cross term ≤ 0; remainder $\le$ Lemma B.2; $\dot V_\pi \le$ (B) at every sample	`T::test_theorem_B_inequality_along_trajectories`
B with a moving ideal, $\varepsilon \ne 0$, switches and active projection	the same plus $V_* W$; at least one switch; projection active at ≥ 10 samples; Dini quotient $\le$ (B)	`T::test_theorem_B_moving_ideal_nonrepresentable_switches_and_active_projection`
B, Dini form	forward-difference quotient of $V_{\mathrm{OT}} \le$ (B)	`T::test_dini_derivative_of_V_OT_bounded_by_active_branch`
B against labelling	matched error $\ll$ labelled error; the bound is 30–45× smaller	`T::test_permuted_ideal_labelled_vs_OT`, `exp_transport_permutation.py`
B.5: failure of the plain bound along (L_σ); the (L_σ) variant holds	with every C(iv) hypothesis asserted, the plain bound fails by $> 3\times$ and the leak variant holds	`T::test_corollary_B5_fails_along_lsigma_and_its_leak_variant_holds`

continued on the next page

Table 26, continued

statement	what is checked	where
C(i), Lemma C.3 (unprojected pure leak)	zero switches and exact $e^{-\sigma t}$ decay (explicit Euler, pure leak)	`T::test_theorem_C_i_exact_dissipation_pure_leak`
C(ii)	$C \le b^2 + 4\eta R$; $\sqrt{C} \le b + 2\eta$; $C = b^2$ under both gap conditions (300 configurations)	`T::test_lemma_C_ii_induced_coupling_bound`
C(iv)	an instance satisfying **every** hypothesis, asserted, including the gap condition $4\bar{\eta}(b+\bar{\eta}) < \Delta_{\mathrm{gap}}$; the envelope holds at every sample, with actual contraction	`T::test_theorem_C_iv_exponential_estimate`
C(v), Proposition C.5	label invariance of the OT leak; the reshuffling transient of the labelled leak	`exp_ot_sigma_mod.py`, `exp_sigma_mod_transient.py`
D	consistency in m (nearly constant amplification 1.64–1.69; not the constant $e^{\Lambda T}$)	`exp_mean_field_m.py`
F(ii) along (L)	particle path length ($\ge$ Wasserstein length) $\le \lambda_G^{1/2} L_1 \int \|e\|$ at every sample	`T::test_theorem_F_wasserstein_length_bounded_by_integrated_error`
F(ii′) along (L_σ)	$W(\mu_{t_1}, \mu_{t_0}) \le \int (\lambda_G^{1/2} L_1 \|e\| + \sigma_m W(\mu_r, \mu^0))$ on a grid of pairs; a tight instance with $e \equiv 0$ (ratio ≥ 0.99), where the motion is $W \approx 1.05$	`T::test_theorem_F_leak_clause_along_ot_sigma_mod`, `T::test_theorem_F_leak_clause_is_tight_when_the_leak_pays_for_all_motion`
E linearisation	Jacobian of the implemented loop equals $A(t)$; $QA + A^\top Q = \mathrm{diag}(-k_e I, 0)$	`T::test_theorem_E_linearisation_and_lyapunov_identity`
E convergence	the matched error decays and the labelled error does not (qualitative); sliding-window excitation and fitted rates	`T::test_theorem_E_matched_error_converges_labelled_does_not`, `exp_pe_convergence.py`
G(a), two scales	direct check (cosine ≥ 0.998 in the G^{-1} metric, `violating`; the weight block fails in `satisfying`, as explained in Section 16); $\bar{L}$ at period-mean parameters non-increasing	`exp_variable_projection.py` → `table_violating.md`, `table_satisfying.md`
G(b), (b′), (H2c)	VarPro holds when (H2b) and $S \ll 1$ hold; it fails when $S = O(1)$ although (H2b) holds; the induced-metric flow tracks the closed loop at every S (started on the slow manifold; gains $\div 4$ isolates the averaging error)	`exp_variable_projection.py` → `table_satisfying.md`; `exp_slow_manifold_slope.py` → `table_slope.md`
H(i), algebraic identity	with $e := r/k_e$ and $x := x_m$, the averaged drift equals $-G\nabla V$ (code gradients against finite differences, 10^{-7}); does not test averaging	`Lg::test_averaged_drift_is_the_mean_field_langevin_drift`
Remark H.0	the frozen-μ averaging error decays like $1/k_e$, $n = 1$ and $n = 2$	`Lg::test_averaging_error_of_the_formal_reduction_decreases_like_one_over_ke`, `Av::test_averaged_drift_error_decays_like_one_over_ke_for_an_orbit_enclosing_area`
H, need for entropy	finite-m real loop: a noiseless symmetric start stays bit-identical (and moves); dither separates it	`Lg::test_noiseless_symmetric_initialisation_never_separates_but_dither_does`
H(ii), oscillation bound	$\|V_L[\mu](\vartheta)\| \le K_\Phi(F_\rho + K_\Phi)$ for random and adversarial μ, ϑ	`Lg::test_first_variation_bound_used_in_holley_stroock`
H(ii) constant	formula transcription only	`Lg::test_lsi_lower_bound_formula`
H, global behaviour	collapsed against spread starts, with and without dither, two (σ, τ) settings	`exp_langevin_global.py`
H, speed–bias trade-off	escape time and loss floor against τ; $\log(1/\tau)$ and Arrhenius fits, seed-averaged and per seed (neither identifies the mechanism)	`exp_langevin_tradeoff.py`

continued on the next page

Table 26, continued

statement	what is checked	where
H, reference values	$\bar{L}(\mu_\tau^*)$ by a concave dual on a grid; the stationary state of the exactly averaged loop; their offset in the scaling of H.6	`exp_gibbs_reference.py` → `results/langevin_global/gibbs_reference.md`
H.7	expansion, Stokes area identity, parity under reversal, non-reciprocity of the exact second variation	`Av::test_frozen_measure_expansion_converges_at_rate_one_over_ke`, `Av::test_circulation_kernel_antisymmetric_vanishes_in_1d_and_equals_the_enclosed_area_integral`, `Av::test_shift_correction_is_even_and_lag_correction_is_odd_under_orbit_reversal`, `Av::test_exactly_averaged_dynamics_is_non_reciprocal_for_n1_and_n2`
H.7a, oscillation term	value against an independent one-period near-identity flow (within 3% at $k_e = 80$); odd under orbit reversal (10^{-6})	`Av::test_oscillation_correction_matches_a_one_period_flow`, `Av::test_oscillation_correction_is_odd_under_orbit_reversal`
H.7 + H.7a in the real closed loop	the orientation dependence of the learned kernels (same ρ) is predicted to within 3% in size and 11% as a vector, cosine 0.994 at $k_e = 40$; the measured difference scales like $k_e^{-1.01}$	`exp_orbit_orientation.py` → `results/orbit_orientation/table_sigma2_slow16.md`

Not checked in the repository:

- B.5 along (L), and B.6 and B.7;
- the Filippov form of C(i), the inequality of C(v), and Proposition C.5's $D^+V_{\mathrm{OT}} = \sigma_m(\delta - \delta^2)$;
- D(b), D's constant, and the one-sided constant of Remark D.1;
- the rate of E, Corollary E.2 and Remark E.3;
- H.8, which is a proof only;
- Conjecture H.6.

References

[1] S. K. Ainsworth, J. Hayase, S. Srinivasa. *Git Re-Basin: merging models modulo permutation symmetries*. International Conference on Learning Representations (ICLR), 2023. arXiv:2209.04836.

[2] L. Ambrosio, N. Gigli, G. Savaré. *Gradient Flows in Metric Spaces and in the Space of Probability Measures*. 2nd edition, Lectures in Mathematics ETH Zürich. Birkhäuser, Basel, 2008. https://doi.org/10.1007/978-3-7643-8722-8.

[3] D. Bakry, M. Émery. *Diffusions hypercontractives*. In J. Azéma, M. Yor (eds.), Séminaire de Probabilités XIX, 1983/84, Lecture Notes in Mathematics 1123, pp. 177–206. Springer, Berlin, 1985. https://doi.org/10.1007/BFb0075847.

[4] R. Barboni, G. Peyré, F.-X. Vialard. *Ultra-fast feature learning for the training of two-layer neural networks in the two-timescale regime*. arXiv:2504.18208, 2025.

[5] E. A. Carlen, K. Craig. *Contraction of the proximal map and generalized convexity of the Moreau–Yosida regularization in the 2-Wasserstein metric*. Mathematics and Mechanics of Complex Systems, 1(1):33–65, 2013. https://doi.org/10.2140/memocs.2013.1.33. arXiv:1205.6565.

[6] F. Chen, Z. Ren, S. Wang. *Uniform-in-time propagation of chaos for mean field Langevin dynamics*. Annales de l'Institut Henri Poincaré, Probabilités et Statistiques, 61(4):2357–2404, 2025. https://doi.org/10.1214/24-AIHP1499. arXiv:2212.03050.

[7] J. Chevallier. *Uniform decomposition of probability measures: quantization, clustering and rate of convergence*. Journal of Applied Probability, 55(4):1037–1045, 2018. https://doi.org/10.1017/jpr.2018.69.

[8] L. Chizat, F. Bach. *On the global convergence of gradient descent for over-parameterized models using optimal transport*. Advances in Neural Information Processing Systems 31 (NeurIPS 2018), pp. 3036–3046, 2018. arXiv:1805.09545.

[9] L. Chizat. *Mean-field Langevin dynamics: exponential convergence and annealing*. Transactions on Machine Learning Research, 2022. arXiv:2202.01009.

[10] L. Chizat. *Sparse optimization on measures with over-parameterized gradient descent*. Mathematical Programming, 194(1–2):487–532, 2022. https://doi.org/10.1007/s10107-021-01636-z. arXiv:1907.10300.

[11] G. Chowdhary, E. Johnson. *Concurrent learning for convergence in adaptive control without persistency of excitation*. In Proc. 49th IEEE Conference on Decision and Control (CDC), pp. 3674–3679, 2010. https://doi.org/10.1109/CDC.2010.5717148.

[12] G. V. Chowdhary, E. N. Johnson. *Theory and flight-test validation of a concurrent-learning adaptive controller*. Journal of Guidance, Control, and Dynamics, 34(2):592–607, 2011. https://doi.org/10.2514/1.46866.

[13] K. Craig, K. Elamvazhuthi, M. Haberland, O. Turanova. *A blob method for inhomogeneous diffusion with applications to multi-agent control and sampling*. Mathematics of Computation, 92(344):2575–2654, 2023. https://doi.org/10.1090/mcom/3841. arXiv:2202.12927.

[14] Y. De Castro, S. Gadat, C. Marteau. *Fast Spawn&Prune (FS&P): global convergence of stochastic conic particle gradient descent via birth/death process*. arXiv:2605.19784, 2026.

[15] M. Emerick, B. Bamieh. *Continuum swarm tracking control: a geometric perspective in Wasserstein space*. In Proc. 62nd IEEE Conference on Decision and Control (CDC), pp. 1367–1374, 2023. https://doi.org/10.1109/CDC49753.2023.10384000. arXiv:2303.15638.

[16] S. Fabri, V. Kadirkamanathan. *Dynamic structure neural networks for stable adaptive control of nonlinear systems*. IEEE Transactions on Neural Networks, 7(5):1151–1167, 1996. https://doi.org/10.1109/72.536311.

[17] M. Fornasier, B. Piccoli, F. Rossi. *Mean-field sparse optimal control*. Philosophical Transactions of the Royal Society A, 372(2028):20130400, 2014. https://doi.org/10.1098/rsta.2013.0400. arXiv:1402.5657.

[18] N. Fournier, A. Guillin. *On the rate of convergence in Wasserstein distance of the empirical measure*. Probability Theory and Related Fields, 162(3–4):707–738, 2015. https://doi.org/10.1007/s00440-014-0583-7.

[19] A. Frieze, R. Kannan. *Log-Sobolev inequalities and sampling from log-concave distributions*. The Annals of Applied Probability, 9(1):14–26, 1999. https://doi.org/10.1214/aoap/1029962595.

[20] M. Fruchart, R. Hanai, P. B. Littlewood, V. Vitelli. *Non-reciprocal phase transitions*. Nature, 592(7854):363–369, 2021. https://doi.org/10.1038/s41586-021-03375-9.

[21] S. S. Ge, C. Wang. *Direct adaptive NN control of a class of nonlinear systems*. IEEE Transactions on Neural Networks, 13(1):214–221, 2002. https://doi.org/10.1109/72.977306.

[22] R. Giard, Y. De Castro, R. Denis, C. Marteau. *Riemannian gradient descent for Gaussian mixture models with unknown diagonal covariances*. arXiv:2609.30220, 2026.

[23] M. Glasgow, J. Bruna. *Uniform-in-time weak propagation of chaos in shallow neural networks*. Advances in Neural Information Processing Systems 39 (NeurIPS 2026), to appear. arXiv:2605.22010, 2026.

[24] F. Golse. *On the dynamics of large particle systems in the mean field limit*. Lecture notes, arXiv:1301.5494, 2013. Published in A. Muntean, J. Rademacher, A. Zagaris (eds.), Macroscopic and Large Scale Phenomena: Coarse Graining, Mean Field Limits and Ergodicity, Lecture Notes in Applied Mathematics and Mechanics 3, pp. 1–144. Springer, 2016. https://doi.org/10.1007/978-3-319-26883-5_1.

[25] S. Graf, H. Luschgy. *Foundations of Quantization for Probability Distributions*. Lecture Notes in Mathematics 1730. Springer, Berlin, 2000. https://doi.org/10.1007/BFb0103945.

[26] J. Guo, S. T. Paruchuri, A. J. Kurdila. *Partial persistence of excitation in RKHS embedded adaptive estimation*. IEEE Transactions on Automatic Control, 68(10):5850–5861, 2023. https://doi.org/10.1109/TAC.2022.3226710. arXiv:2002.07963.

[27] R. Holley, D. Stroock. *Logarithmic Sobolev inequalities and stochastic Ising models*. Journal of Statistical Physics, 46(5–6):1159–1194, 1987. https://doi.org/10.1007/BF01011161.

[28] D. Inoue, Y. Ito, H. Yoshida. *Optimal transport-based coverage control for swarm robot systems: generalization of the Voronoi tessellation-based method*. IEEE Control Systems Letters, 5(4):1483–1488, 2021. https://doi.org/10.1109/LCSYS.2020.3039008. arXiv:2011.08337.

[29] P. A. Ioannou, P. V. Kokotović. *Adaptive Systems with Reduced Models*. Lecture Notes in Control and Information Sciences 47. Springer, Berlin, 1983. https://doi.org/10.1007/BFb0006357.

[30] P. A. Ioannou, J. Sun. *Robust Adaptive Control*. Prentice Hall, Upper Saddle River, NJ, 1996.

[31] H. K. Khalil. *Nonlinear Systems*. 3rd edition. Prentice Hall, Upper Saddle River, NJ, 2002.

[32] P. V. Kokotović, H. K. Khalil, J. O'Reilly. *Singular Perturbation Methods in Control: Analysis and Design*. Academic Press, London, 1986. Reprinted as Classics in Applied Mathematics 25, SIAM, 1999, https://doi.org/10.1137/1.9781611971118.

[33] N. Lanzetti, E. C. Balta, D. Liao-McPherson, F. Dörfler. *Stochastic Wasserstein gradient flows using streaming data with an application in predictive maintenance*. IFAC-PapersOnLine (IFAC World Congress 2023), 56(2):3954–3959, 2023. https://doi.org/10.1016/j.ifacol.2023.10.1334. arXiv:2301.12461.

[34] E. Lavretsky, T. E. Gibson. *Projection operator in adaptive systems*. arXiv:1112.4232, 2011.

[35] E. Lavretsky, K. A. Wise. *Robust and Adaptive Control: With Aerospace Applications*. Advanced Textbooks in Control and Signal Processing. Springer, London, 2013. https://doi.org/10.1007/978-1-4471-4396-3.

[36] F. L. Lewis, A. Yeşildirek, K. Liu. *Multilayer neural-net robot controller with guaranteed tracking performance*. IEEE Transactions on Neural Networks, 7(2):388–399, 1996. https://doi.org/10.1109/72.485674.

[37] F. L. Lewis, S. Jagannathan, A. Yeşildirek. *Neural Network Control of Robot Manipulators and Nonlinear Systems*. Taylor & Francis, London, 1999.

[38] P. Marion, R. Berthier. *Leveraging the two-timescale regime to demonstrate convergence of neural networks*. Advances in Neural Information Processing Systems 36 (NeurIPS 2023), pp. 64996–65029, 2023. arXiv:2304.09576.

[39] A. P. Morgan, K. S. Narendra. *On the stability of nonautonomous differential equations* $\dot{x} = [A + B(t)]x$, *with skew symmetric matrix* $B(t)$. SIAM Journal on Control and Optimization, 15(1):163–176, 1977. https://doi.org/10.1137/0315013.

[40] K. S. Narendra, A. M. Annaswamy. *A new adaptive law for robust adaptation without persistent excitation*. IEEE Transactions on Automatic Control, 32(2):134–145, 1987. https://doi.org/10.1109/TAC.1987.1104543.

[41] A. Nitanda, D. Wu, T. Suzuki. *Convex analysis of the mean field Langevin dynamics*. Proceedings of the 25th International Conference on Artificial Intelligence and Statistics (AISTATS), PMLR 151:9741–9757, 2022. arXiv:2201.10469.

[42] F. Otto, C. Villani. *Generalization of an inequality by Talagrand and links with the logarithmic Sobolev inequality*. Journal of Functional Analysis, 173(2):361–400, 2000. https://doi.org/10.1006/jfan.1999.3557.

[43] S. T. Paruchuri, J. Guo, A. J. Kurdila. *Kernel center adaptation in the reproducing kernel Hilbert space embedding method*. International Journal of Adaptive Control and Signal Processing, 36(7):1562–1583, 2022. https://doi.org/10.1002/acs.3407. arXiv:2009.02867.

[44] O. S. Patil, E. J. Griffis, W. A. Makumi, W. E. Dixon. *Simultaneous online system identification and control using composite adaptive Lyapunov-based deep neural networks*. arXiv:2311.13056, 2023 (revised 2026).

[45] O. S. Patil. *Adaptive control as an information gradient flow: excitation, constraints, energy, and stochastic learning*. arXiv:2610.04822, 2026.

[46] J. C. Perdomo, T. Zrnic, C. Mendler-Dünner, M. Hardt. *Performative prediction*. Proceedings of the 37th International Conference on Machine Learning (ICML), PMLR 119:7599–7609, 2020. arXiv:2002.06673.

[47] S. Plunder, B. Simeon. *Coupled systems of linear differential-algebraic and kinetic equations with application to the mathematical modelling of muscle tissue*. In T. Reis, S. Grundel, S. Schöps (eds.), Progress in Differential-Algebraic Equations II, Differential-Algebraic Equations Forum, pp. 357–395. Springer, 2020. https://doi.org/10.1007/978-3-030-53905-4_12. arXiv:1911.05468 (2019).

[48] S. Plunder, B. Simeon. *The mean-field limit for particle systems with uniform full-rank constraints*. Kinetic and Related Models, 16(6):884–912, 2023. https://doi.org/10.3934/krm.2023012. arXiv:2203.07249 (2022).

[49] M. M. Polycarpou. *Stable adaptive neural control scheme for nonlinear systems*. IEEE Transactions on Automatic Control, 41(3):447–451, 1996. https://doi.org/10.1109/9.486648.

[50] J.-B. Pomet, L. Praly. *Adaptive nonlinear regulation: estimation from the Lyapunov equation*. IEEE Transactions on Automatic Control, 37(6):729–740, 1992. https://doi.org/10.1109/9.256328.

[51] J. A. Sanders, F. Verhulst, J. Murdock. *Averaging Methods in Nonlinear Dynamical Systems*. 2nd edition, Applied Mathematical Sciences 59. Springer, New York, 2007. https://doi.org/10.1007/978-0-387-48918-6.

[52] R. M. Sanner, J.-J. E. Slotine. *Gaussian networks for direct adaptive control*. IEEE Transactions on Neural Networks, 3(6):837–863, 1992. https://doi.org/10.1109/72.165588.

[53] F. Santambrogio. *{Euclidean, metric, and Wasserstein} gradient flows: an overview*. Bulletin of Mathematical Sciences, 7(1):87–154, 2017. https://doi.org/10.1007/s13373-017-0101-1. arXiv:1609.03890.

[54] S. Sastry, M. Bodson. *Adaptive Control: Stability, Convergence, and Robustness*. Prentice Hall, Englewood Cliffs, NJ, 1989.

[55] N. J. Tatro, P.-Y. Chen, P. Das, I. Melnyk, P. Sattigeri, R. Lai. *Optimizing mode connectivity via neuron alignment*. Advances in Neural Information Processing Systems 33 (NeurIPS 2020), pp. 15300–15311, 2020. arXiv:2009.02439.

[56] J. N. Tsitsiklis, B. Van Roy. *An analysis of temporal-difference learning with function approximation*. IEEE Transactions on Automatic Control, 42(5):674–690, 1997. https://doi.org/10.1109/9.580874.

[57] F.-Y. Wang. *On estimation of the logarithmic Sobolev constant and gradient estimates of heat semigroups*. Probability Theory and Related Fields, 108(1):87–101, 1997. https://doi.org/10.1007/s004400050102.

[58] Y. Zhang, Q. Cai, Z. Yang, Y. Chen, Z. Wang. *Can temporal-difference and Q-learning learn representation? A mean-field theory*. Advances in Neural Information Processing Systems 33 (NeurIPS 2020), pp. 19680–19692, 2020. arXiv:2006.04761.