

LECTURE NOTES

AS AN AUDIT

GROUP THEORY APPLIED TO CONDENSED MATTER PHYSICS

PHY889

by

Dimitri Bonanni-Surprenant

Lecture by

Maia Vergniory



Sherbrooke's University
Faculty of Science
Department of Physics
September 13, 2024

Table des matières

1	Abstract Groups	3
1.1	Translation group	3
1.2	Subgroups	3
1.3	The order of an element	3
1.4	Cosets	4
1.5	Conjugate elements and Class	5
1.6	Invariant subgroup	6
1.7	The factor group	6
1.8	Isomorphism and Homomorphism of groups	7
1.9	Exercices	7

1 Abstract Groups

By denoting the elements of a group by certain symbols which obey a given rule of multiplication, we obtain the so-called abstract group. In this chapter, we will review some of its properties.

1.1 Translation group

Suppose that the group G consists of elements g_1, g_2, \dots, g_m . Let us multiply its elements on the right by some element g_ℓ , i.e., let's carry a right translation along the group.

$$g_1 g_\ell, g_2 g_\ell, \dots, g_m g_\ell \quad (1)$$

We see that each group element is encountered once and only once in this sequence. Suppose that an element could be encountered more than once such that $g_i g_\ell = g_j g_\ell$, then this would imply that $g_i = g_j$ by the existence of the inverse. Since there are as many elements as elements in the group, and they are all unique, then because a group is closed this implies that all elements are represented. Let g_ℓ be an arbitrary element of the group, it is clear that $(g_\ell g_\ell^{-1}) g_\ell = g_\ell$, and, consequently, the element g_ℓ also appears in the sequence. Since the number of elements in our sequence is the order of the group, each element can be found in the sequence only once. The sequence of elements $g_\ell g_1, g_\ell g_2, \dots, g_\ell g_m$ has the same properties.

1.2 Subgroups

A set of elements belonging to a group G , to which itself forms a group with the same multiplication rule, is a subgroup of G . The remainder of the group G cannot form a group since it will not contain the unit element.

1.3 The order of an element

Let us take an arbitrary element g_i of a group G and consider the powers of g_i, g_i^2, g_i^3, \dots . Since we are considering a finite group, the members of this sequence must appear repeatedly. Suppose for example that

$$g_i^{k_1} = g_i^{k_2} = g_i \quad k_2 > k_1 \quad (2)$$

$$\rightarrow g_i^{k_2} = g_i^{k_1} g_i^{k_2 - k_1} = g_i g_i^{k_2 - k_1} = g_i \implies g_i^{k_2 - k_1} = E. \quad (3)$$

The smallest exponent h for which $g_i^h = E$ is the order of the element g_i . The set of elements $g_i, g_i^2, \dots, g_i^h = E$ is the period of the element g_i . It is clear that the period of an element of the group is a subgroup of G . All the elements of this subgroup commute and consequently, the subgroup is Abelian.

If h is the order of the element g_j , then $g_j^{h-1} = g_j^{-1}$, therefore for finite groups, the existence of the inverse is a consequence of the group order properties.

Proof. Recall the definition of a group.

- Composition law. $\forall a, b \in G, a \cdot b \in G$.
- Associativity. $\forall a, b, c \in G, (a \cdot b) \cdot c = a \cdot (b \cdot c)$.
- Neutral element. $\exists E \in G \ni \forall a \in G, E \cdot a = a \cdot E = a$.
- Symmetry. $\forall a \in G \exists a^{-1} \in G \ni a \cdot a^{-1} = a^{-1} \cdot a = E$.

The goal of this proof is to show that for finite groups, we can relax the symmetry axiom as it is a consequence of the group order properties. This section already proved most of this proof. ■

1.4 Cosets

Let H be a subgroup of a group G , h_1, h_2, \dots, h_m , where m is the order of H . Let us now construct the following sequences of sets of elements of G .

- Let us take g_1 contained in G , which is not contained in H , and construct the set $g_1h_1, g_1h_2, \dots, g_1h_m$, which we denote by g_1H .
- Now we take another element of G , $g_2 \notin H$, and set up g_2H .
- \vdots
-

As a result, we obtain the sequence $H, g_1H, g_2H, \dots, g_{k-1}H$. The sets g_iH are the *left cosets* of the subgroup H . We shall show that the cosets defined above have no common elements. Let us suppose that the sets g_1H and g_2H have one common element.

$$g_1h_1 = g_2h_2 \rightarrow g_2 = g_1h_1h_2^{-1} = g_1h_3 \quad (4)$$

so that g_2 belongs to the g_1H set. But this conflicts with the original assumption that each element of a group G enters only one of the cosets. Since G contains n elements, and each of the cosets contains m elements, it follows that $m = \frac{n}{k}$. The number k is the index of the subgroup H . We see that the order of the subgroup is a divisor of the order of the group.

Similarly, we can do the same with the right cosets, in constructing the cosets, we have a choice in selecting the element g_i . It can be shown that for an acceptable choice of the elements g_i , we can obtain the same set of cosets and the same decomposition. This follows directly from the following theorem.

Two cosets g_iH and g_kH (being g_i, g_k any two elements of G) either coincide or have no common elements. If these sets have at least one common element $g_ih_\alpha = g_kh_\beta \rightarrow g_k = g_ih_\alpha h_\beta^{-1} \implies g_k \in g_iH$. However, any element of the set g_kH can be represented in the form $g_kh_j = g_ih_\alpha h_\beta h_\beta^{-1}h_j = g_ih_\gamma$ and will also belong to g_iH . The group G can therefore uniquely decompose into (left or right) cosets of subgroup H .

1.5 Conjugate elements and Class

Let g be an element of the group G and let us construct the element $g' = g_i g g_i^{-1}$, $g_i \in G$. The elements g and g' are said to be conjugate. Let us suppose that g_i runs over all elements of the group G . We then obtain n elements, some of which might be equal. Let the number of distinct elements be k , let us denote them by g_1, g_2, \dots, g_k . It is clear that this set includes all the elements of the group G , which are conjugate to the element g . All the elements of this set are mutually conjugate.

Let

$$g_1 = g_\alpha g g_\alpha^{-1} \quad (5)$$

$$g_2 = g_\beta g g_\beta^{-1}. \quad (6)$$

Then we have

$$g = g_\alpha^{-1} g_1 g_\alpha \quad (7)$$

$$g_2 = g_\beta g_\alpha^{-1} g_1 g_\alpha g_\beta^{-1} = g_\beta g_\alpha^{-1} g_1 (g_\beta g_\alpha^{-1})^{-1}. \quad (8)$$

The set of all the mutually conjugate elements form a *class*. Thus the elements g_1, g_2, \dots, g_k form a class of conjugate elements. The number of elements in a class is its order. Any finite group can be denoted into a number of classes of conjugate elements. The unit element of a group is a class by itself. All the elements of a given class have the same order. The product of the elements of two classes consists of whole classes.

$$C_i C_j = \sum_k h_{ijk} C_k \quad (9)$$

where k is the multiplicity of the class. If $g_p \in C_i C_j$, then the entire class C_p to which g_p belongs itself belongs to the set $C_i C_j$. Let

$$g_p = g_i g_j \cdot \begin{cases} g_i \in C_i \\ g_j \in C_j \end{cases} \quad (10)$$

We then have that for any $g \in G$,

$$g^{-1} g_p g = g^{-1} g_i g g^{-1} g_j g \in C_i C_j \quad (11)$$

it remains to show that each element of the class C_p enters the set $C_i C_j$ the same number of times. Suppose that the element g_p enters twice, $g_i g_j = g_p$ and $g'_i g'_j = g_p$, where $g_i \neq g'_i$ and $g_j \neq g'_j$. Each element $g'^{-1} g_p g'$ ($g' \in G$) will C_p then be contained in $C_i C_j$ at least twice.

$$g'^{-1} g_p g' = g'^{-1} (g_i g_j) g' = (g'^{-1} g_i g') (g'^{-1} g_j g') \quad (12)$$

$$g'^{-1} g_p g' = g'^{-1} (g'_i g'_j) g' = (g'^{-1} g'_i g') (g'^{-1} g'_j g') \quad (13)$$

if we consider the result, it follows that $g'^{-1} g_i g' \neq g'^{-1} g'_i g'$. The element $g'^{-1} g_p g'$ will not be encountered more than twice, since otherwise g_p would be encountered more than twice, which contradicts the original argument. ■

1.6 Invariant subgroup

Let H be a subgroup of the group G , and suppose that $g_i \in G$. Consider the set of elements $g_i H g_i^{-1}$, where g_i is fixed. That set is also a group, since all the group axioms are satisfied for it. If $g_i \in H$, then the similar subgroup will coincide with H . If $g_i \notin H$, then in general we obtain a subgroup of which is different from H . When the subgroup H coincide with all its similar subgroups, it is called an invariant subgroup or a normal divisor. An invariant subgroup is represented by the letter N . It follows from this definition that if an invariant subgroup contains an element $g \in G$, then it will also contain the entire class to which g belongs. The invariant subgroup consists of the whole class of the group.

For an invariant subgroup of G , the left and right cosets coincide.

$$g_i N = g_i N g_i^{-1} g_i = N g_i \quad \text{since } g_i N g_i^{-1} = N. \quad (14)$$

Any group has two trivial invariant subgroups: the group itself and the unit element. Groups which do not have invariant subgroups other than the trivial ones are called simple. It would be useful to prove that $NN = N$.

1.7 The factor group

Let N be an invariant subgroup of the group G . Let us decompose G into the cosets N ,

$$N, g_1 N, g_2 N, \dots, g_{k-1} N \quad (15)$$

and let's form a set $g_1 N g_2 N$, which consists of different elements $g_1 h_\alpha g_2 h_\beta$, where h_α and h_β run independently over the entire subgroup N . We can see that

$$g_1 N g_2 N = g_1 g_2 g_2^{-1} N g_2 N = g_1 g_2 N N = g_1 g_2 N = g_3 N \quad (16)$$

if the set $g_1 N g_2 N$ is called the product of sets $g_1 N$ and $g_2 N$, then the product of two cosets of N will again give a coset N , since the multiplication of a coset of N by N on the right and left does not change these cosets.

$$N g_1 N = g_1 g_1^{-1} N g_1 N = g_1 N N = g_1 N \quad (17)$$

For each coset $g_i N$, there is a coset $g_i^{-1} N$ such that the product is equal to N .

$$g_i^{-1} N g_i N = N N = N \quad (18)$$

The cosets of an invariant subgroup can be regarded as the elements of a new group in which N plays the role of the unit element. This group is what we call *the factor group* or *the quotient group* of the invariant subgroup. Its order is equal to the order of N .

1.8 Isomorphism and Homomorphism of groups

If between the elements of two groups there is a one-to-one correspondence which preserves group multiplication, then the groups are *isomorphic*. Let G and \tilde{G} be two isomorphic groups, then if elements g_i and g_k of G correspond to the elements \tilde{g}_i and \tilde{g}_k of \tilde{G} , i.e., $g_i \leftarrow \tilde{g}_i$, $g_k \leftarrow \tilde{g}_k$, then $g_1 = g_i g_k \rightarrow \tilde{g}_1 = \tilde{g}_i \tilde{g}_k$. We can reduce the investigation of a given group to that of another group isomorphic to it.

If to each element of G there corresponds only one definite element of \tilde{G} , and to each element of \tilde{G} a number of elements of G , and this correspondence is preserved under group multiplication, \tilde{G} is *homomorphic* to G . Homomorphism has the following properties:

- If \tilde{G} is homomorphic to G , then the element of G corresponds to the unit element of \tilde{G} .
- The mutually reciprocal elements of G correspond to the mutually reciprocal elements of \tilde{G} .

$$g_1 g_k = E \quad \tilde{g}_i \tilde{g}_k = \tilde{E}. \quad (19)$$

- All elements of G , to which corresponds to the unit element of \tilde{G} form an invariant subgroup N of the group G . Elements g'_1, g'_2, \dots, g'_j of G correspond to \tilde{E} of \tilde{G} . The product $g'_i g'_k$ correspond to $\tilde{E} \tilde{E} = \tilde{E}$. $g'_i g'_k = g'_\ell$ and the set g'_1, g'_2, \dots, g'_j is closed with respect to group multiplication.

from (a), it contains a unit element, but since \tilde{E} is the inverse of itself and because of (b) for each element g'_ℓ we can find the inverse. From $\tilde{g} \tilde{E} \tilde{g}^{-1} = \tilde{E}$, where \tilde{g} is an arbitrary element of \tilde{G} , it follows that $g g_i g^{-1} = g'_f$ for an arbitrary element of G .

$$g'_1, g'_2, \dots, g'_s \quad (20)$$

forms an *invariant* subgroup of G .

1.9 Exercises

Consider the group S_6 , symmetric group of order 6, with the elements $\{E, A, B, C, D, F\}$.

	E	A	B	C	D	F
E	E	A	B	C	D	F
A	A	E	D	F	B	C
B	B	F	E	D	C	A
C	C	D	F	E	A	B
D	D	C	A	B	F	E
F	F	B	C	E	E	D

1. Find the order of all the elements.
2. Find the possible subgroups.

1 Abstract Groups

3. Divide the group into cosets, and verify that this can be done in a unique way.
4. Divide the group into classes of conjugate elements.
5. Find the invariant subgroups and verify that the left and right cosets are the same.
6. Write down the multiplication table for the corresponding factor groups.
7. Show that the abstract group has the following realizations
 - (a) Permutation group of 3 elements.
 - (b) Matrix group of order 2: S_6 can also be 2×2 matrices corresponding to rotations and reflections.

Recall the definition of the order of an element ([subsection 1.3](#)). Then we can compute for all elements the smallest exponent h such that $g^h = E$. From the table of multiplication, we can see that the order of all the elements except for the unit element and F and D is 2. The order of the unit element is 1. The order of D and F is 3.

Recall the definition of the subgroup ([subsection 1.2](#)). Then the subgroups are

$$\begin{aligned} &\{E\} \\ &\{A, E\}, \{B, E\}, \{C, E\} \\ &\{D, F, E\} \\ &\{A, B, C, D, E, F\} \end{aligned}$$

Recall the definition of the cosets ([subsection 1.4](#)). Let's start with $H = \{E\}$. The left and right cosets are then equal and form G . Let's continue with $H = \{A, E\}$, we then have the left cosets $\{D, B\}, \{F, C\}$.

References