

Optimal acquisition scheme for subwavelength localization microscopy of bleachable fluorophores: erratum

Edward Shore¹ and Alex Small^{2,*}

¹Mathematics and Statistics Department, California State Polytechnic University,
3801 West Temple Avenue, Pomona, California 91768, USA

²Physics Department, California State Polytechnic University, 3801 West Temple Avenue,
Pomona, California 91768, USA

*Corresponding author: arsmall@csupomona.edu

Received July 11, 2011; accepted July 11, 2011;
posted July 11, 2011 (Doc. ID 150681); published August 8, 2011

One of the plots in Fig. 2 of our previous Letter [Opt. Lett. **36**, 289 (2011)] was mislabeled with an incorrect parameter value and is corrected here. © 2011 Optical Society of America

OCIS codes: 180.2520, 100.6640.

In our recently published Letter [1], the third plot in Fig. 2 was incorrectly labeled. The parameter value used to generate that plot was $f_1 = 0.5$, not $f_1 = 0.05$. A corrected version of the figure is included (Fig. 1).

References

1. E. Shore and A. Small, Opt. Lett. **36**, 289 (2011).




Fig. 1. (This is a corrected version of Fig. 2 of [1].) p_γ versus E_2 for different b and f_1 values. E_2 ranges with different qualitative features are noted.