

ERASMUS UNIVERSITY ROTTERDAM

ERASMUS SCHOOL OF ECONOMICS

Master's Thesis Econometrics and Management Science - Business Analytics and Quantitative
Marketing

Median-of-Means Two-Stage Least Squares Regression

David von Schledorn Márquez (597334)



Supervisor:	Mikhail Zhelonkin
Second assessor:	-
Date final version:	-

The views stated in this thesis are those of the author and not necessarily those of the supervisor, second assessor, Erasmus School of Economics or Erasmus University Rotterdam.

Abstract

Abstract

1 Introduction

Foundational papers on Median-of-Means: Nemirovskij and Yudin (1983) and Jerrum, Valiant and Vazirani (1986). For giving the framework and template on the proofs: Devroye, Lerasle, Lugosi and Oliveira (2016) and Lugosi and Mendelson (2019).

2 Median-of-Means (MoM) Estimator

The following section is adapted from Devroye et al. (2016) and Lugosi and Mendelson (2019)

2.1 Sub-Gaussian Estimators

Definition 1. *Sub-Gaussian Estimators* Let $(X_i)_{i=1}^n$ be a random sample with $\mathbb{E}[X_i] = \mu$ and $\text{Var}[X_i] = \sigma^2$, and let $\bar{\mu}(X)$ denote an estimator for μ . For some constant $L > 0$, $\bar{\mu}(X)$ is said to be a sub-Gaussian estimator for μ if

$$\mathbb{P}\left(|\bar{\mu}(X) - \mu| > L\sigma\sqrt{\frac{\ln(1/\delta)}{n}}\right) \leq \delta \quad \text{for all } \delta \in (0, 1). \quad (1)$$

2.2 The Empirical Mean Estimator

Let $(X_i)_{i=1}^n$ be a random sample with $\mathbb{E}[X_i] = \mu$ and $\text{Var}[X_i] = \sigma^2 < \infty$. The empirical mean is defined as

$$\bar{\mu} = \frac{1}{n} \sum_{i=1}^n X_i.$$

By the Central Limit Theorem,

$$\sqrt{n}(\bar{\mu} - \mu) \xrightarrow{d} \mathcal{N}(0, \sigma^2),$$

which implies that for any fixed $\delta \in (0, 1)$,

$$\mathbb{P}\left(|\bar{\mu} - \mu| > \frac{\sigma\Phi^{-1}(1 - \delta/2)}{\sqrt{n}}\right) \rightarrow \delta \quad \text{as } n \rightarrow \infty.$$

Using the bound $\Phi^{-1}(1 - \delta/2) \leq \sqrt{2\ln(2/\delta)}$, it follows that

$$\lim_{n \rightarrow \infty} \mathbb{P}\left(|\bar{\mu} - \mu| > \sigma\sqrt{\frac{2\ln(2/\delta)}{n}}\right) \leq \delta,$$

so $\bar{\mu}$ is an asymptotically sub-Gaussian estimator in the sense of (1) with constant $L = \sqrt{2}$.

While $\bar{\mu}$ is asymptotically sub-Gaussian, it is not a sub-Gaussian estimator for finite samples;

Chebyshev's inequality gives

$$\mathbb{P}\left(|\bar{\mu} - \mu| > \sigma \sqrt{\frac{1}{n\delta}}\right) \leq \delta$$

shows that the dependence on δ is polynomial rather than logarithmic, which is substantially worse than (1) for small δ . As such, the empirical mean is insufficient as a sub-Gaussian estimator for random variables with only finite first and second moments.

2.3 The Median-of-Means Estimator

The median-of-means (MoM) estimator is not new, being proposed in various forms, first by Nemirovskij and Yudin (1983) and Jerrum et al. (1986). The algorithm works as follows

Algorithm 1: Median-of-Means Estimator

Input: Sample $(X_i)_{i=1}^n$, number of blocks k

Output: Estimate $\hat{\mu}$

Partition $(X_i)_{i=1}^n$ into k blocks $B_1, \dots, B_k$ of size $m = \lfloor n/k \rfloor$;

for $j = 1, \dots, k$ **do**

 Compute the block mean $\bar{\mu}_j = \frac{1}{m} \sum_{i \in B_j} X_i$;

return $\hat{\mu} = \text{median}(\bar{\mu}_1, \dots, \bar{\mu}_k)$;

The MoM estimator satisfies

$$\mathbb{P}\left(|\hat{\mu} - \mu| > \sigma \sqrt{\frac{4}{m}}\right) \leq e^{-k/8},$$

and setting $k = \lceil 8 \ln(1/\delta) \rceil$, this becomes

$$\mathbb{P}\left(|\hat{\mu} - \mu| > \sigma \sqrt{\frac{32 \ln(1/\delta)}{n}}\right) \leq \delta \quad \text{for all } \delta \in (0, 1) \quad (2)$$

which is sub-Gaussian with $L = \sqrt{32}$

3 Median-of-Means Two-Stage Least Squares (MoM-2SLS)

We first explore the MoM estimator in a univariate setting, with one instrument and one variable.

3.1 Univariate MoM-2SLS

In the univariate setting, the 2SLS estimator collapses to the Wald/IV estimator

$$\hat{\beta}_{IV} = \frac{\sum_{i=1}^n Z_i Y_i}{\sum_{i=1}^n Z_i X_i} = \frac{\bar{S}_{ZY}}{\bar{S}_{ZX}}.$$

This is a ratio of two sample means. To solve this, we can utilise two MoM-type estimators. The first replaces each sample mean with its MoM counterpart; we call this the ratio-of-medians estimator. The second returns the median of the Wald/IV estimate of each block, we call this the median-of-ratios estimator.

3.2 Setup

Let $(Y_i, X_i, Z_i)_{i=1}^n$ be an i.i.d. random sample satisfying the structural equation

$$Y_i = \beta X_i + \varepsilon_i,$$

where $\mathbb{E}[Z_i \varepsilon_i] = 0$ (exogeneity) and $\mu_{ZX} := \mathbb{E}[Z_i X_i] \neq 0$ (relevance). Define

$$\mu_{ZY} := \mathbb{E}[Z_i Y_i], \quad \sigma_{ZY}^2 := \text{Var}(Z_i Y_i), \quad \sigma_{ZX}^2 := \text{Var}(Z_i X_i), \quad \sigma_{Z\varepsilon}^2 := \text{Var}(Z_i \varepsilon_i),$$

all assumed finite. Both estimators partition the sample into k blocks $B_1, \dots, B_k$ of size $m = \lfloor n/k \rfloor$ and compute block means

$$\hat{S}_{ZY}^{(j)} = \frac{1}{m} \sum_{i \in B_j} Z_i Y_i, \quad \hat{S}_{ZX}^{(j)} = \frac{1}{m} \sum_{i \in B_j} Z_i X_i.$$

They differ only in how the block-level quantities are aggregated.

3.3 The Estimators

Definition 2 (Ratio-of-Medians (RoM)).

$$\hat{\beta}_{\text{RoM}} = \frac{\tilde{S}_{ZY}}{\tilde{S}_{ZX}}, \quad \text{where } \tilde{S}_{ZY} = \text{median}_j(\hat{S}_{ZY}^{(j)}), \quad \tilde{S}_{ZX} = \text{median}_j(\hat{S}_{ZX}^{(j)}).$$

The medians are taken separately and the ratio is formed afterwards.

Definition 3 (Median-of-Ratios (MoR)).

$$\hat{\beta}_{\text{MoR}} = \text{median}_j \left(\hat{\beta}^{(j)} \right), \quad \text{where } \hat{\beta}^{(j)} = \frac{\hat{S}_{ZY}^{(j)}}{\hat{S}_{ZX}^{(j)}}.$$

The ratio is formed within each block and the median is taken over the block-level estimates.

3.4 Concentration Results

Theorem 4 (Standard IV). *Suppose $n \geq 8\sigma_{ZX}^2/(\delta \mu_{ZX}^2)$, then*

$$\mathbb{P} \left[\left| \hat{\beta}_{\text{IV}} - \beta \right| > \frac{2\sqrt{2} \sigma_{Z\varepsilon}}{|\mu_{ZX}| \sqrt{\delta n}} \right] \leq \delta. \quad (3)$$

Theorem 5 (Ratio-of-Medians). *Let $k = \lceil 8 \ln(2/\delta) \rceil$ and $m = \lfloor n/k \rfloor$. Suppose $m \geq 4\sigma_{ZX}^2/\mu_{ZX}^2$. Then*

$$\mathbb{P} \left[\left| \hat{\beta}_{\text{RM}} - \beta \right| > \frac{2(\sigma_{ZY} + |\beta| \sigma_{ZX})}{|\mu_{ZX}| \sqrt{m} - 2\sigma_{ZX}} \right] \leq \delta. \quad (4)$$

Theorem 6 (Median-of-Ratios). *Let $k = \lceil 8 \ln(1/\delta) \rceil$ and $m = \lfloor n/k \rfloor$. Suppose $m \geq 32\sigma_{ZX}^2/\mu_{ZX}^2$. Then*

$$\mathbb{P} \left[\left| \hat{\beta}_{\text{MR}} - \beta \right| > \frac{4\sqrt{2} \sigma_{Z\varepsilon}}{|\mu_{ZX}| \sqrt{m}} \right] \leq \delta. \quad (5)$$

The proofs for these results can be found in the tablet notes.

The error bounds can then be written as

$$\left| \hat{\beta} - \beta \right| \leq C \cdot \frac{\sigma_{\text{eff}}}{|\mu_{ZX}|} \cdot g(\delta, n),$$

where C is an absolute constant, σ_{eff} is an effective standard deviation, and $g(\delta, n)$ captures the dependence on confidence and sample size. Note that this definition is directly comparable to 1. We utilise these definitions for the following comparisons.

3.5 Comparison of the Estimators

3.5.1 Numerical Impact

δ	$1/\sqrt{\delta}$ (IV)	$\sqrt{\ln(2/\delta)}$ (RoM)	$\sqrt{\ln(1/\delta)}$ (MoR)	IV / MoR ratio
0.10	3.16	1.73	1.52	2.1
0.05	4.47	1.92	1.73	2.6
0.01	10.0	2.30	2.15	4.7
10^{-3}	31.6	2.74	2.63	12.0
10^{-6}	10^3	3.80	3.72	269
10^{-9}	3.16×10^4	4.60	4.55	6950

3.5.2 Effective Variance

The RoM and IV bound have $\sigma_{ZY} + |\beta|\sigma_{ZX}$ in the numerator while the MoR has $\sigma_{Z\varepsilon}$. Using $Z_i Y_i = \beta Z_i X_i + Z_i \varepsilon_i$, it can be shown that

$$\sigma_{Z\varepsilon} \leq \sigma_{ZY} + |\beta|\sigma_{ZX}.$$

Thus, the MoR and IV bounds are tighter. The proof is found in the tablet notes.

3.5.3 Number of Blocks

3.5.4 Instrument Strength Condition

	Standard IV	RoM	MoR
Min. block size	-	$\frac{4\sigma_{ZX}^2}{\mu_{ZX}^2}$	$\frac{32\sigma_{ZX}^2}{\mu_{ZX}^2}$
Min. sample size	$\frac{8\sigma_{ZX}^2}{\delta \mu_{ZX}^2}$	$\frac{32\sigma_{ZX}^2}{\mu_{ZX}^2} \ln(2/\delta)$	$\frac{256\sigma_{ZX}^2}{\mu_{ZX}^2} \ln(1/\delta)$
δ dependence	Polynomial	Logarithmic	Logarithmic

The standard IV instrument strength condition depends inversely on $1/\delta$, meaning that achieving higher confidence requires proportionally more samples. The MoR and RoM estimators depend logarithmically on δ , which is a significant improvement over the standard IV.

Comparing the two MoM-based estimators, the RoM condition is eight times less stringent than the MoR condition.

3.5.5 Conclusion for the Univariate Case

no single estimator dominates. The standard IV has the tightest constants but the weakest instrument strength for moderate δ . Its polynomial tail makes it unsuitable for high-confidence inference. The MoR achieves the tightest sub-Gaussian bound, but requires the strongest instrument condition. The RoM has a mild instrument condition, but a looser effective variance.

References

- Devroye, L., Lerasle, M., Lugosi, G. & Oliveira, R. I. (2016). Sub-gaussian mean estimators.
- Jerrum, M. R., Valiant, L. G. & Vazirani, V. V. (1986). Random generation of combinatorial structures from a uniform distribution. *Theoretical computer science*, 43, 169–188.
- Lugosi, G. & Mendelson, S. (2019). Mean estimation and regression under heavy-tailed distributions: A survey. *Foundations of Computational Mathematics*, 19(5), 1145–1190.
- Nemirovskij, A. S. & Yudin, D. B. (1983). Problem complexity and method efficiency in optimization.