


IMAGES IN CLINICAL MEDICINE

Stephanie V. Sherman, M.D., *Editor*

Pupillary Light–Near Dissociation



Emanuele La Corte, M.D., Ph.D.

Alma Mater Studiorum University
of Bologna
Bologna, Italy
emanuele.lacorte2@unibo.it

Matteo Martinoni, M.D.

IRCCS Istituto Delle Scienze
Neurologiche di Bologna
Bologna, Italy


A video is
available at
[NEJM.org](https://www.nejm.org)

A 37-YEAR-OLD MAN PRESENTED TO THE NEUROSURGERY CLINIC WITH A 1-year history of blurry vision and 1 week of headache and difficulty looking upward. The neurologic examination was notable for vertical and horizontal diplopia, upward gaze palsy, and vertical misalignment of the eyes that varied with gaze position. In addition, his pupils only minimally constricted to light but constricted to near objects, a finding known as light–near dissociation (Panel A and video). Magnetic resonance imaging of the brain showed an enhancing dorsal midbrain and medial thalamic lesion with obstructive hydrocephalus (Panel B, arrow). Pupillary light–near dissociation is characterized by the absence of constriction to light with an intact accommodation reflex. Causes of light–near dissociation include dorsal rostral midbrain lesions, as in this case, and neurosyphilis, though the Argyll Robertson pupils associated with syphilis are also typically miotic and irregular at baseline. After noninvasive testing did not confirm the diagnosis, ventriculostomy and biopsy of the lesion were performed, and histologic analysis showed a nonsecreting germinoma. The patient was treated with platinum-based chemotherapy followed by radiation, which resulted in complete resolution of the lesion. At 1-year follow-up, his pupillary responses were normal to both light and accommodation; however, he had persistent diplopia and disconjugate gaze.

DOI: 10.1056/NEJMicm2031597

Copyright © 2021 Massachusetts Medical Society.