Name:

Mark: ____

Mini-math Div 3/4: Monday, December 7, 2020 (12 minutes)

1. Determine all horizontal and vertical asymptotes of the following functions, or indicate there are no such asymptotes. (Show your work for vertical asymptotes via a sign analysis)

(a) (2 points)
$$f(x) = \frac{3x^2 - 9x}{x^2 - 2x - 3}$$

Solution: Notice

$$\lim_{x \to \pm \infty} f(x) = \lim_{x \to \pm \infty} \frac{3 - \frac{9}{x}}{1 - \frac{2}{x} - \frac{3}{x^2}} = 3$$

so y = 3 is the only horizontal asymptote.

Notice $f(x) = \frac{3x(x-3)}{(x-3)(x+1)}$, so x=3 and x=-1 are potential vertical asymptotes.

Near -1^- , the fraction is of the form $\frac{(-)(-)}{(-)(0^-)}$, so $\lim_{x\to -1^-} f(x) = \infty$, while

$$\lim_{x \to 3} f(x) = \lim_{x \to 3} \frac{3x}{x+1} = \frac{9}{4}$$

so only x = -1 is a vertical asymptote.

(b) (2 points) $f(x) = \frac{2x^3 + 10x}{x^2 + 3x + 2}$

Solution: Notice

$$\lim_{x \to \pm \infty} f(x) = \lim_{x \to \pm \infty} \frac{2x + \frac{10}{x}}{1 + \frac{3}{x} + \frac{2}{x^2}} = \pm \infty$$

so there are no horizontal asymptotes.

Notice $f(x) = \frac{2x(x^2+5)}{(x+2)(x+1)}$, so x = -2 and x = -1 are potential vertical asymptotes.

Near -2^- , the fraction is of the form $\frac{(-)(+)}{(0^-)(-)}$, so $\lim_{x\to -2^-} f(x) = -\infty$, while near -1^- , the fraction is of the form $\frac{(-)(+)}{(+)(0^-)}$, so $\lim_{x\to -1^-} f(x) = \infty$. Then both x=-2,-1 are vertical asymptotes.

2. (2 points) Consider the function $f(x) = x^4 + 6x^3 + 12x^2 + x + 1$. Find the interval(s) on which f is concave up.

Solution: Differentiating twice,

$$f'(x) = 4x^3 + 18x^2 + 24x + 1,$$

$$f''(x) = 12x^2 + 36x + 24$$

We find the critical points: f''(x) always exists, and

$$0 = 12x^{2} + 36x + 24 = 12(x^{2} + 3x + 2) = 12(x+1)(x+2)$$

so the critical points are x = -2, -1.

On each subinterval, the second derivative has the following sign:

		-2		-1	
x+2	_		+		+
x + 1	_		_		+
f''(x)	+		_		+

Then f(x) is concave up on $(-\infty, -2)$ and $(-1, \infty)$.