

Numerical Methods

Numerical Linear Algebra

Martijn Boussé, Pieter Collins & Başak Sakçak

Department of Advanced Computing Sciences

Maastricht University

`m.bousse,pieter.collins,basak.sakcak@maastrichtuniversity.nl`

KEN1540 & KEN2540, Block 5, April-May 2025

Algebraic Methods	2
Gaussian Elimination	4
LU Factorisation	5
Sparsity	14
Norms and Conditioning	19
Iterative refinement	23
Iterative Methods	24
Iterative method	25
Jacobi	26
Gauss-Seidel	28
Over-Relaxation	34
Conjugate-Gradient	36
Conjugate-Gradient	37
Eigenvalues	43
Eigenvalues	44
Approximation	45
Power method	46
Inverse power method	50
Deflation	52
Orthogonalisation	53
Orthogonality	54
Orthogonalisation	55
The QR Method	59
QR method	60
Householder matrices	65
Givens matrices	71
Eigenvalue condition	76

Matrix form

System of linear algebraic equations:

$$\begin{aligned} 3x_1 - 7x_2 - 2x_3 + 2x_4 &= -9 \\ -3x_1 + 5x_2 + x_3 &= 5; \\ 6x_1 - 4x_2 + 2x_3 - 5x_4 &= 7; \\ -9x_1 + 5x_2 - 5x_3 + 6x_4 &= -19. \end{aligned}$$

Write using matrix form $Ax = b$ where

$$A = \begin{pmatrix} 3 & -7 & -2 & 2 \\ -3 & 5 & 1 & 0 \\ 6 & -4 & 2 & -5 \\ -9 & 5 & -5 & 6 \end{pmatrix}, \quad x = \begin{pmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{pmatrix}, \quad b = \begin{pmatrix} -9 \\ 5 \\ 7 \\ -19 \end{pmatrix}.$$

3 / 77

Gaussian Elimination

Solve by Gaussian elimination:

$$\begin{aligned} \left(\begin{array}{cccc|c} 3 & -7 & -2 & 2 & -9 \\ -3 & 5 & 1 & 0 & 5 \\ 6 & -4 & 2 & -5 & 7 \\ -9 & 5 & -5 & 6 & -19 \end{array} \right) &\sim \begin{matrix} r_1 \\ r_2 - (-1)r_1 \\ r_3 - (+2)r_1 \\ r_4 - (-3)r_1 \end{matrix} \left(\begin{array}{cccc|c} 3 & -7 & -2 & 2 & -9 \\ 0 & -2 & -1 & 2 & -4 \\ 0 & 10 & 6 & -9 & 25 \\ 0 & -16 & -11 & 12 & -46 \end{array} \right) \\ &\sim \begin{matrix} r_1 \\ r_2 \\ r_3 - (-5)r_2 \\ r_4 - (+8)r_2 \end{matrix} \left(\begin{array}{cccc|c} 3 & -7 & -2 & 2 & -9 \\ 0 & -2 & -1 & 2 & -4 \\ 0 & 0 & 1 & 1 & 5 \\ 0 & 0 & -3 & -4 & -14 \end{array} \right) \\ &\sim \begin{matrix} r_1 \\ r_2 \\ r_3 \\ r_4 - (-3)r_3 \end{matrix} \left(\begin{array}{cccc|c} 3 & -7 & -2 & 2 & -9 \\ 0 & -2 & -1 & 2 & -4 \\ 0 & 0 & 1 & 1 & 5 \\ 0 & 0 & 0 & -1 & 1 \end{array} \right) \end{aligned}$$

Solve by *backsubstitution* $x_4 = -1$, $x_3 = 6$, $x_2 = -2$, $x_1 = -3$.

4 / 77

LU Factorisation

Define L by letting L_{ij} be the multiple of row j subtracted from row i , and U the tableau matrix before backsubstitution.

$$L = \begin{pmatrix} 1 & 0 & 0 & 0 \\ -1 & 1 & 0 & 0 \\ 2 & -5 & 1 & 0 \\ -3 & 8 & -3 & 1 \end{pmatrix}, \quad U = \begin{pmatrix} 3 & -7 & -2 & 2 \\ 0 & -2 & -1 & 2 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & -1 \end{pmatrix}$$

Notice that $LU = A$!

$$LU = \begin{pmatrix} 1 & 0 & 0 & 0 \\ -1 & 1 & 0 & 0 \\ 2 & -5 & 1 & 0 \\ -3 & 8 & -3 & 1 \end{pmatrix} \begin{pmatrix} 3 & -7 & -2 & 2 \\ 0 & -2 & -1 & 2 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & -1 \end{pmatrix} = \begin{pmatrix} 3 & -7 & -2 & 2 \\ -3 & 5 & 1 & 0 \\ 6 & -4 & 2 & -5 \\ -9 & 5 & -5 & 6 \end{pmatrix} = A.$$

Solving $Ly = b$ gives $y = (-9, -4, 5, 1)^T$. Notice that this y is the right-hand column of the tableau before backsubstitution.

5 / 77

LU Factorisation

Factorization $A = LU$ where L is unit lower triangular $l_{ii} = 1$, $l_{ij} = 0$ for $j > i$ and U is upper triangular $u_{ij} = 0$ for $i > j$.

Backsubstitution The linear system $LUx = b$ can be solved by computing $Ly = b$ and $Ux = y$. Since L and U are triangular, the linear systems can be easily solved by *backsubstitution*

$$y_i = b_i - \sum_{j < i} l_{ij} y_j; \quad x_i = (y_i - \sum_{j > i} u_{ij} x_j) / u_{ii}.$$

6 / 77

LU Factorisation—Complexity

Complexity Computing the LU -factorisation:

$$n(n-1) + (n-1)(n-2) + \cdots + 3 \cdot 2 + 2 \cdot 1 = (n^3 - n)/3 \sim (n^3/3) = O(n^3).$$

Solving the system $Ly = b$ requires

$$(n-1) + (n-2) + \cdots + 2 + 1 = (n^2 - n)/2$$

operations, and solving $Ux = y$ requires

$$n + (n-1) + \cdots + 2 + 1 = (n^2 + n)/2$$

operations, so computing x from L, U, b requires n^2 operations.

Hence for large systems, most of the work in solving $Ax = b$ lies in computing the LU -factorisation.

Once the LU -factorisation has been computed for a given A , the systems $Ax_i = b_i$ can easily be solved for different vectors b_i .

7 / 77

LU Factorisation—Pivoting

Let $A^{(k-1)}$ be the matrix obtained after working on the first $k - 1$ columns.

Pivoting If $A_{kk}^{(k-1)}$ is equal to zero, then Gaussian elimination needs to swap rows.

Obtain a factorisation $PA = LU$, where P is a *permutation matrix*.

Partial pivoting Make the *pivot* element $A_{kk}^{(k-1)}$ the largest of $A_{kj}^{(k-1)}$ for $j \geq k$.

Scaled partial pivoting Define scale factor $s_k = \max_{1 \leq j \leq k} |a_{kj}|$.

Choose first row to maximise $|a_{k1}|/s_k = |a_{k1}|/\max_{1 \leq j \leq k} |a_{kj}|$.

8 / 77

LU Factorisation—Pivoting Example (Omit)

Example Compute the LU-factorisation of A using partial pivoting and solve $Ax = b$ where

$$A = \begin{pmatrix} 2.11 & -4.21 & 0.921 \\ 1.09 & 0.987 & 0.832 \\ 4.01 & 10.2 & -1.12 \end{pmatrix}, \quad b = \begin{pmatrix} 2.01 \\ 4.21 \\ -3.09 \end{pmatrix}.$$

Swap row 1 and row 3 and eliminate x_1 :

$$\begin{aligned} T_1 A &= \begin{pmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{pmatrix} \begin{pmatrix} 2.11 & -4.21 & 0.921 \\ 1.09 & 0.987 & 0.832 \\ 4.01 & 10.2 & -1.12 \end{pmatrix} = \begin{pmatrix} 4.01 & 10.2 & -1.12 \\ 1.09 & 0.987 & 0.832 \\ 2.11 & -4.21 & 0.921 \end{pmatrix} \\ &= \begin{pmatrix} 1 & 0 & 0 \\ 0.272 & 1 & 0 \\ 0.526 & 0 & 1 \end{pmatrix} \begin{pmatrix} 4.01 & 10.2 & -1.12 \\ 0 & -1.79 & 1.14 \\ 0 & -9.58 & 1.51 \end{pmatrix} = L_1 U_1 \end{aligned}$$

9 / 77

LU Factorisation—Pivoting Example (Omit)

Example $T_1 A = L_1 U_1$ given by

$$\begin{pmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{pmatrix} \begin{pmatrix} 1.09 & 0.987 & 0.832 \\ 2.11 & -4.21 & 0.921 \\ 4.01 & 10.2 & -1.12 \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 \\ 0.272 & 1 & 0 \\ 0.526 & 0 & 1 \end{pmatrix} \begin{pmatrix} 4.01 & 10.2 & -1.12 \\ 0 & -1.79 & 1.14 \\ 0 & -9.58 & 1.51 \end{pmatrix}$$

Swap row 2 and row 3, so $T_2 T_1 A = (T_2 L_1 T_2^{-1})(T_2 U_1)$, and eliminate x_2 :

$$\begin{aligned} T_2 T_1 A &= \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix} \begin{pmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{pmatrix} \begin{pmatrix} 2.11 & -4.21 & 0.921 \\ 1.09 & 0.987 & 0.832 \\ 4.01 & 10.2 & -1.12 \end{pmatrix} \\ &= \begin{pmatrix} 1 & 0 & 0 \\ 0.526 & 1 & 0 \\ 0.272 & 0 & 1 \end{pmatrix} \begin{pmatrix} 4.01 & 10.2 & -1.12 \\ 0 & -9.58 & 1.51 \\ 0 & -1.79 & 1.14 \end{pmatrix} \\ &= \begin{pmatrix} 1 & 0 & 0 \\ 0.526 & 1 & 0 \\ 0.272 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0.186 & 1 \end{pmatrix} \begin{pmatrix} 4.01 & 10.2 & -1.12 \\ 0 & -9.58 & 1.51 \\ 0 & 0 & 0.855 \end{pmatrix} \\ &= (T_2 L_1 T_2^{-1}) L_2 U_2 \end{aligned}$$

10 / 77

LU Factorisation—Pivoting Example (Omit)

Example Obtain $PA = LU$ with

$$\begin{aligned} PA &= \begin{pmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix} \begin{pmatrix} 2.11 & -4.21 & 0.921 \\ 1.09 & 0.987 & 0.832 \\ 4.01 & 10.2 & -1.12 \end{pmatrix} = \begin{pmatrix} 4.01 & 10.2 & -1.12 \\ 2.11 & -4.21 & 0.921 \\ 1.09 & 0.987 & 0.832 \end{pmatrix} \\ &= \begin{pmatrix} 1 & 0 & 0 \\ 0.526 & 1 & 0 \\ 0.272 & 0.186 & 1 \end{pmatrix} \begin{pmatrix} 4.01 & 10.2 & -1.12 \\ 0 & -9.58 & 1.51 \\ 0 & 0 & 0.855 \end{pmatrix} = LU \end{aligned}$$

Solve $Ax = b$ using $PAx = LUx = Pb$, so $Ux = y$ where $Ly = Pb$.

$$Pb = \begin{pmatrix} -3.09 \\ 2.01 \\ 4.21 \end{pmatrix}; \quad y = L \backslash Pb = \begin{pmatrix} -3.09 \\ 3.64 \\ 4.37 \end{pmatrix}; \quad x = U \backslash y = \begin{pmatrix} -0.428 \\ 0.427 \\ 5.11 \end{pmatrix}.$$

11 / 77

LU Factorisation—Pivoting Example (Omit)

Example Compute y by backsubstitution:

$$Ly = \begin{pmatrix} 1 & 0 & 0 \\ 0.526 & 1 & 0 \\ 0.272 & 0.186 & 1 \end{pmatrix} \begin{pmatrix} y_1 \\ y_2 \\ y_3 \end{pmatrix} = \begin{pmatrix} -3.09 \\ 2.01 \\ 4.21 \end{pmatrix} = Pb$$

$$y_1 = -3.09;$$

$$0.526y_1 + y_2 = 2.01 \implies$$

$$\begin{aligned} y_2 &= 2.01 - 0.526y_1 = 2.01 - 0.526 \times (-0.309) \\ &= 3.64 \end{aligned}$$

$$0.272y_1 + 0.186y_2 + y_3 = 4.21 \implies$$

$$\begin{aligned} y_3 &= 4.21 - 0.272y_1 - 0.186y_2 = 4.21 - 0.272 \times (-0.309) - 0.186 \times 3.64 \\ &= 4.37 \end{aligned}$$

12 / 77

LU Factorisation—Pivoting Example (Omit)

Example Compute x by backsubstitution:

$$Ux = \begin{pmatrix} 4.01 & 10.2 & -1.12 \\ 0 & -9.58 & 1.51 \\ 0 & 0 & 0.855 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} -3.09 \\ 3.64 \\ 4.37 \end{pmatrix} = y$$

$$0.855x_3 = 4.37 \implies$$

$$x_3 = 4.37 \div 0.855 \\ = 5.11$$

$$-9.58x_2 + 1.51x_3 = 3.64 \implies$$

$$x_2 = (3.64 - 1.51x_3) \div (-9.58) = (3.64 - 1.51 \times 5.11) \div (-9.58) \\ = 0.427$$

$$4.01x_1 + 10.2x_2 - 1.12x_3 = -3.09 \implies$$

$$x_1 = (-3.09 - 10.2x_2 + (-1.12)x_3) \div 4.01 \\ = (-3.09 - 10.2 \times 0.427 + 1.12 \times 5.11) \div 4.01 \\ = -0.428$$

13 / 77

Sparsity

Sparsity A matrix is *sparse* if it has many zero elements.

Fill-in The inverse of a sparse matrix is usually dense.

Tridiagonal matrices A is *tridiagonal* if $a_{ij} = 0$ for $|i - j| > 1$.

LU-factorisation *preserves* the zeros for a *banded* matrix!

Complexity Computing $A^{-1}b$ if A is tridiagonal requires n^2 operations; solving $LUx = b$ requires $\sim 3n$ operations!

14 / 77

Sparsity

Example For the given tridiagonal matrix, the inverse is

$$\begin{pmatrix} 4 & 1 & 0 & 0 & 0 \\ 1 & 4 & 1 & 0 & 0 \\ 0 & 1 & 4 & 1 & 0 \\ 0 & 0 & 1 & 4 & 1 \\ 0 & 0 & 0 & 1 & 4 \end{pmatrix}^{-1} = \begin{pmatrix} 0.268 & -0.072 & 0.019 & -0.005 & 0.001 \\ -0.072 & 0.287 & -0.077 & 0.021 & -0.005 \\ 0.019 & -0.077 & 0.288 & -0.077 & 0.019 \\ -0.005 & 0.021 & -0.077 & 0.287 & -0.072 \\ 0.001 & -0.005 & 0.019 & -0.072 & 0.268 \end{pmatrix}$$

and the LU-factorisation is

$$\begin{pmatrix} 1 & 0 & 0 & 0 & 0 \\ 0.250 & 1 & 0 & 0 & 0 \\ 0 & 0.267 & 1 & 0 & 0 \\ 0 & 0 & 0.268 & 1 & 0 \\ 0 & 0 & 0 & 0.268 & 1 \end{pmatrix} \begin{pmatrix} 4.000 & 1 & 0 & 0 & 0 \\ 0 & 3.750 & 1 & 0 & 0 \\ 0 & 0 & 3.733 & 1 & 0 \\ 0 & 0 & 0 & 3.732 & 1 \\ 0 & 0 & 0 & 0 & 3.732 \end{pmatrix}$$

Clearly, solving $LUx = b$ by backsubstitution is faster than computing $A^{-1}x$.

15 / 77

Sparsity

Example For the LU-factorisation

$$LU = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 \\ 0.250 & 1 & 0 & 0 & 0 \\ 0 & 0.267 & 1 & 0 & 0 \\ 0 & 0 & 0.268 & 1 & 0 \\ 0 & 0 & 0 & 0.268 & 1 \end{pmatrix} \begin{pmatrix} 4.000 & 1 & 0 & 0 & 0 \\ 0 & 3.750 & 1 & 0 & 0 \\ 0 & 0 & 3.733 & 1 & 0 \\ 0 & 0 & 0 & 3.732 & 1 \\ 0 & 0 & 0 & 0 & 3.732 \end{pmatrix}$$

we have $(LU)^{-1} = U^{-1}L^{-1}$ given by

$$\begin{pmatrix} 0.250 & -0.067 & 0.018 & -0.005 & 0.001 \\ 0 & 0.267 & -0.071 & 0.019 & -0.005 \\ 0 & 0 & 0.268 & -0.072 & 0.019 \\ 0 & 0 & 0 & 0.268 & -0.072 \\ 0 & 0 & 0 & 0 & 0.268 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 & 0 & 0 \\ -0.250 & 1 & 0 & 0 & 0 \\ 0.067 & -0.267 & 1 & 0 & 0 \\ -0.018 & 0.071 & -0.268 & 1 & 0 \\ 0.005 & -0.019 & 0.072 & -0.268 & 1 \end{pmatrix}$$

Clearly, solving $LUx = b$ by backsubstitution is faster than computing L^{-1} , U^{-1} and $x = U^{-1}(L^{-1}b)$

16 / 77

Symmetric matrices

Symmetric matrices A symmetric matrix ($A = A^T$) can be factorised $A = LDL^T$ where L is unit-lower-triangular and D is diagonal.

Positive-definite matrices A matrix is *positive definite* if $x^T Ax > 0$ whenever $x \neq 0$; equivalently, if all eigenvalues are positive, or if $A = LDL^T$ with D having strictly-positive diagonal elements.

Cholesky factorisation If A is positive definite, then there is an upper-triangular matrix U such that $A = U^T U$ (or a lower-triangular matrix L such that $A = LL^T$.)

17 / 77

Symmetric matrices—Example (Omit)

Example Compute the LDL^T factorisation:

$$\begin{aligned} A = \begin{pmatrix} 2 & 1 & 0 \\ 1 & 2 & 1 \\ 0 & 1 & 2 \end{pmatrix} &= \begin{pmatrix} 1 & 0 & 0 \\ 1/2 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 2 & 1 & 0 \\ 0 & 3/2 & 1 \\ 0 & 1 & 2 \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 \\ 1/2 & 1 & 0 \\ 0 & 2/3 & 1 \end{pmatrix} \begin{pmatrix} 2 & 1 & 0 \\ 0 & 3/2 & 1 \\ 0 & 0 & 4/3 \end{pmatrix} \\ &= \begin{pmatrix} 1 & 0 & 0 \\ 1/2 & 1 & 0 \\ 0 & 2/3 & 1 \end{pmatrix} \begin{pmatrix} 2 & 0 & 0 \\ 0 & 3/2 & 0 \\ 0 & 0 & 4/3 \end{pmatrix} \begin{pmatrix} 1 & 1/2 & 0 \\ 0 & 1 & 2/3 \\ 0 & 0 & 1 \end{pmatrix} = LDL^T \end{aligned}$$

The Cholesky factorisation is given by $A = U^T U$ where

$$U = D^{1/2} L^T = \begin{pmatrix} \sqrt{2} & 0 & 0 \\ 0 & \sqrt{3/2} & 0 \\ 0 & 0 & \sqrt{4/3} \end{pmatrix} \begin{pmatrix} 1 & 1/2 & 0 \\ 0 & 1 & 2/3 \\ 0 & 0 & 1 \end{pmatrix} = \begin{pmatrix} \sqrt{2/1} & \sqrt{1/2} & 0 \\ 0 & \sqrt{3/2} & \sqrt{2/3} \\ 0 & 0 & \sqrt{4/3} \end{pmatrix}$$

18 / 77

Vector and matrix norms

Properties A *norm* is a measure of the *magnitude* of a vector (or matrix).

A vector norm $\|\cdot\|$ is a function $\mathbb{R}^n \rightarrow \mathbb{R}$ which satisfies

$$\begin{aligned}\|v\| &\geq 0, \quad \|v\| = 0 \iff v = 0; \\ \|\alpha v\| &= |\alpha| \cdot \|v\|; \quad \|u + v\| \leq \|u\| + \|v\|.\end{aligned}$$

A matrix norm additionally satisfies

$$\|AB\| \leq \|A\| \cdot \|B\|.$$

Given a vector norm $\|\cdot\|_*$, the corresponding matrix norm is

$$\|A\|_* = \max\{\|Ax\|_* \mid \|x\|_* = 1\}$$

and satisfies

$$\|Ax\|_* \leq \|A\|_* \times \|x\|_*$$

20 / 77

Vector and matrix norms

p-norms Important vector norms are

$$\begin{aligned}\|v\|_p &:= \left(\sum_{i=1}^n |v_i|^p\right)^{1/p}. \\ \|v\|_\infty &:= \lim_{p \rightarrow \infty} \|v\|_p.\end{aligned}$$

Note

$$\begin{aligned}\|v\|_1 &= \sum_{i=1}^n |v_i|; \\ \|v\|_2 &= \sqrt{\sum_{i=1}^n v_i^2} = \sqrt{v \cdot v}; \\ \|v\|_\infty &= \max_{i=1, \dots, n} |v_i|.\end{aligned}$$

The two-norm $\|v\|_2$ gives the Euclidean length of v . The *uniform* norm $\|v\|_\infty$ gives the maximum absolute value of the components, and is usually easiest to compute.

The corresponding matrix norms are

$$\begin{aligned}\|A\|_1 &= \max_{j=1, \dots, n} \sum_{i=1}^m |a_{ij}|; \\ \|A\|_2 &= \max(\text{eig}(A^T A))^{1/2}; \\ \|A\|_\infty &= \max_{i=1, \dots, m} \sum_{j=1}^n |a_{ij}|.\end{aligned}$$

21 / 77

Condition number

Conditioning For a given vector norm $\|\cdot\|$ and corresponding matrix norm, define the matrix *condition number* $K(A) := \|A\| \times \|A^{-1}\|$.

Suppose \tilde{x} is an approximate solution to $Ax = b$. Then

$$\|\tilde{x} - x\| = \|A^{-1}(A\tilde{x} - Ax)\| \leq \|A^{-1}\| \|A\tilde{x} - b\|$$

so the error satisfies:

$$\|\tilde{x} - x\| \leq K(A) \frac{\|A\tilde{x} - b\|}{\|A\|}.$$

Further, since $\|A\| \cdot \|x\| \geq \|Ax\| = \|b\|$, we have $1/\|x\| \leq \|A\|/\|b\|$, so the relative error satisfies:

$$\frac{\|\tilde{x} - x\|}{\|x\|} \leq K(A) \frac{\|A\tilde{x} - b\|}{\|b\|}$$

Setting $\tilde{b} = A\tilde{x}$, then

$$\frac{\|\tilde{x} - x\|}{\|x\|} \leq K(A) \frac{\|\tilde{b} - b\|}{\|b\|}$$

22 / 77

Iterative refinement

Approximate solution Suppose \tilde{x} is an approximate solution to $Ax = b$.

Refinement Then

$$A(x - \tilde{x}) = Ax - A\tilde{x} = b - A\tilde{x} \approx 0,$$

so

$$x = \tilde{x} + A^{-1}(b - A\tilde{x}).$$

Accuracy If $A^{-1}b$ can be computed less accurately than Ax , refinement typically improves the accuracy of a solution.

23 / 77

Iterative Methods

24 / 77

General iterative method

Fixed-point For the linear system $Ax = b$, write $A = D + E$, where D is “easy” to invert.

Then $Ax = Dx + Ex = b$, so $Dx = b - Ex$ and

$$x = D^{-1}(b - Ex).$$

Alternatively, we can write

$$x = x - D^{-1}(Ax - b).$$

Update Attempt to improve x using the update

$$x' = D^{-1}(b - Ex) = x - D^{-1}(Ax - b).$$

Iteration Use this as a basis for an iterative method

$$x^{(n+1)} = D^{-1}(b - Ex^{(n)}) = x^{(n)} - D^{-1}(Ax^{(n)} - b).$$

25 / 77

Jacobi method

Fixed-point formula We have $\sum_{j=1}^n a_{ij}x_j = b_i$ for $i = 1, \dots, n$.

Write $a_{ii}x_i + \sum_{j=1, j \neq i}^n a_{ij}x_j = b_i$. Rearranging gives

$$x_i = \frac{b_i - \sum_{j \neq i} a_{ij}x_j}{a_{ii}}.$$

We can use this as the basis for an *iterative* method.

Jacobi Method Iterate using

$$x'_i = \frac{b_i - \sum_{j \neq i} a_{ij}x_j}{a_{ii}}.$$

An alternative formula is

$$x'_i = x_i - \frac{\sum_j a_{ij}x_j - b_i}{a_{ii}}.$$

In matrix form

$$x' = D^{-1}(b - Ex) = x - D^{-1}(Ax - b)$$

where D is the diagonal of A and $E = A - D$.

26 / 77

Jacobi method

Example Solve $Ax = b$ using the Jacobi method starting at $x^{(0)} = 0$ for

$$A = \begin{pmatrix} 6 & 2 & 0 \\ 3 & 5 & -1 \\ -2 & 1 & 4 \end{pmatrix}, \quad b = \begin{pmatrix} 3 \\ 4 \\ 1 \end{pmatrix}.$$

$$x_1^{(1)} = (b_1 - a_{12}x_2^{(0)} - a_{13}x_3^{(0)})/a_{11} = (3 - 2 \times 0.0 - 0 \times 0.0)/6 = 0.500;$$

$$x_2^{(1)} = (b_2 - a_{21}x_1^{(0)} - a_{23}x_3^{(0)})/a_{22} = (4 - 3 \times 0.0 - (-1) \times 0.0)/5 = 0.800;$$

$$x_3^{(1)} = (b_3 - a_{31}x_1^{(0)} - a_{32}x_2^{(0)})/a_{33} = (1 - (-2) \times 0.0 - 1 \times 0.0)/4 = 0.250.$$

$$x_1^{(2)} = (b_1 - a_{12}x_2^{(1)} - a_{13}x_3^{(1)})/a_{11} = (3 - 2 \times 0.800 - 0 \times 0.250)/6 = 0.233;$$

$$x_2^{(2)} = (b_2 - a_{21}x_1^{(1)} - a_{23}x_3^{(1)})/a_{22} = (4 - 3 \times 0.500 + 1 \times 0.250)/5 = 0.550;$$

$$x_3^{(2)} = (b_3 - a_{31}x_1^{(1)} - a_{32}x_2^{(1)})/a_{33} = (1 + 2 \times 0.500 - 1 \times 0.800)/4 = 0.300.$$

Continuing yields

$$x^{(3)} = \begin{pmatrix} 0.3167 \\ 0.7200 \\ 0.2292 \end{pmatrix}, \quad x^{(4)} = \begin{pmatrix} 0.2600 \\ 0.6558 \\ 0.2283 \end{pmatrix}, \quad x^{(5)} = \begin{pmatrix} 0.2814 \\ 0.6897 \\ 0.2160 \end{pmatrix}.$$

Estimate error $\|x^{(5)} - x\| \lesssim \|x^{(5)} - x^{(4)}\| = 0.024$. Residual $\|Ax^{(5)} - b\| = 0.076$.

27 / 77

Gauss-Seidel method

Gauss-Seidel Method Rather than update all the x_i simultaneously, we can update in turn. Then

$$x'_i = \frac{b_i - \sum_{j=1}^{i-1} a_{ij}x'_j - \sum_{j=i+1}^n a_{ij}x_j}{a_{ii}}.$$

In other words,

$$\text{for } i = 1, \dots, n, \text{ set } x_i = (b_i - \sum_{j \neq i} a_{ij}x_j)/a_{ii} = x_i - (\sum_{j=1}^n a_{ij}x_j - b_i)/a_{ii}.$$

This is more easily implemented in code:

```
for i=1:n, x(i)=x(i)-(A(i,:)*x-b(i))/A(i,i); end;
```

or using an explicit loop:

```
for i=1:n,  
    ri=-b(i);  
    for j=1:n, ri=ri+A(i,j)*x(j); end;  
    x(i)=x(i)-ri/A(i,i);  
end;
```

28 / 77

Gauss-Seidel method

Example Solve $Ax = b$ using the Gauss-Seidel method for:

$$A = \begin{pmatrix} 6 & 2 & 0 \\ 3 & 5 & -1 \\ -2 & 1 & 4 \end{pmatrix}, \quad b = \begin{pmatrix} 3 \\ 4 \\ 1 \end{pmatrix}, \quad x^{(0)} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}$$

$$x_1^{(1)} = (b_1 - a_{12}x_2^{(0)} - a_{13}x_3^{(0)})/a_{11} = (3 - 2 \times 0.0000 - 0 \times 0.0000)/6 = 0.5000;$$

$$x_2^{(1)} = (b_2 - a_{21}x_1^{(1)} - a_{23}x_3^{(0)})/a_{22} = (4 - 3 \times 0.5000 - (-1) \times 0.0000)/5 = 0.5000;$$

$$x_3^{(1)} = (b_3 - a_{31}x_1^{(1)} - a_{32}x_2^{(1)})/a_{33} = (1 - (-2) \times 0.5000 - 1 \times 0.5000)/4 = 0.3750.$$

$$x_1^{(2)} = (b_1 - a_{12}x_2^{(1)} - a_{13}x_3^{(1)})/a_{11} = (3 - 2 \times 0.5000 - 0 \times 0.3750)/6 = 0.3333;$$

$$x_2^{(2)} = (b_2 - a_{21}x_1^{(2)} - a_{23}x_3^{(1)})/a_{22} = (4 - 3 \times 0.3333 + 1 \times 0.3750)/5 = 0.6750;$$

$$x_3^{(2)} = (b_3 - a_{31}x_1^{(2)} - a_{32}x_2^{(2)})/a_{33} = (1 + 2 \times 0.3333 - 1 \times 0.6750)/4 = 0.2479.$$

$$x^{(1)} = \begin{pmatrix} 0.50000 \\ 0.50000 \\ 0.37500 \end{pmatrix}, \quad x^{(2)} = \begin{pmatrix} 0.33333 \\ 0.67500 \\ 0.2479 \end{pmatrix}, \quad x^{(3)} = \begin{pmatrix} 0.27500 \\ 0.68458 \\ 0.21635 \end{pmatrix}, \quad x^{(4)} = \begin{pmatrix} 0.27181 \\ 0.68019 \\ 0.21568 \end{pmatrix},$$

Convergence is rapid.

29 / 77

Gauss-Seidel method

Example For

$$A = \begin{pmatrix} 3 & -7 & -2 & 2 \\ -3 & 5 & 1 & 0 \\ 6 & -4 & 2 & -5 \\ -9 & 5 & -5 & 6 \end{pmatrix}, \quad b = \begin{pmatrix} -9 \\ 5 \\ 7 \\ -19 \end{pmatrix}, \quad x^{(0)} = \begin{pmatrix} 0 \\ 0 \\ 0 \\ 0 \end{pmatrix}$$

the Gauss-Seidel method gives

$$x^{(1)} = \begin{pmatrix} -3.000 \\ 1.000 \\ 3.500 \\ -3.167 \end{pmatrix}, \quad x^{(2)} = \begin{pmatrix} 6.778 \\ -2.500 \\ 3.083 \\ -2.417 \end{pmatrix}, \quad x^{(3)} = \begin{pmatrix} -11.944 \\ 6.950 \\ -30.958 \\ 14.069 \end{pmatrix}, \quad x^{(4)} = \begin{pmatrix} -4.857 \\ -6.925 \\ 119.365 \\ -66.743 \end{pmatrix}$$

so does not converge!

30 / 77

Gauss-Seidel method

Matrix form Write $A = L + D + U$, where L is strictly lower-triangular, D is diagonal and U is strictly upper triangular.

i.e. $L_{i,j} = 0$ if $i \leq j$, $D_{i,j} = 0$ if $i \neq j$, $U_{i,j} = 0$ if $i \geq j$.

Note that L, U here are not the L and U of the LU factorisation!!

Then the Gauss-Seidel method is given by $x' = D^{-1}(b - Lx' + Ux)$, and rearranging gives

$$\begin{aligned} x' &= (L + D)^{-1}(b - Ux) \\ &= x - (L + D)^{-1}(Ax - b). \end{aligned}$$

31 / 77

Convergence

Theorem An iterative method $x' = Tx + r$ converges if $\|T\| < 1$ for some matrix norm $\|\cdot\|$.

Definition A matrix A is *diagonally-dominant* if $|a_{ii}| > \sum_{j \neq i} |a_{ij}|$ for all i .

Theorem (Convergence of Jacobi / Gauss-Seidel) If A is diagonally-dominant, then the Jacobi and Gauss-Seidel iterations converge to the solution of $Ax = b$.

Preconditioning Can precondition A by multiplying by a matrix P to obtain $(PA)x = Pb$.

Approximate inverse Since if $J \approx I$, then J is diagonally-dominant, precondition by $P \approx A^{-1}$ to obtain $J = PA \approx I$.

32 / 77

Preconditioning

Example Let

$$A = \begin{pmatrix} 3 & -7 & -2 & 2 \\ -3 & 5 & 1 & 0 \\ 6 & -4 & 2 & -5 \\ -9 & 5 & -5 & 6 \end{pmatrix}, \quad b = \begin{pmatrix} -9 \\ 5 \\ 7 \\ -19 \end{pmatrix}, \quad P = \begin{pmatrix} 0 & 0 & -1 & -1 \\ 1 & 2 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix}.$$

Then

$$PA = \begin{pmatrix} 3 & -1 & 3 & -1 \\ -3 & 3 & 0 & 2 \\ 6 & -4 & 2 & -5 \\ 3 & -7 & -2 & 2 \end{pmatrix}, \quad Pb = \begin{pmatrix} 12 \\ 1 \\ 7 \\ -9 \end{pmatrix}.$$

Applying the Gauss-Seidel method to $(PA)x = (Pb)$ gives iterates:

$$x^{(1)} = \begin{pmatrix} 4.0 \\ 4.3 \\ 0.2 \\ 4.8 \end{pmatrix}, \quad x^{(2)} = \begin{pmatrix} 6.9 \\ 4.0 \\ 2.9 \\ 2.1 \end{pmatrix}, \quad x^{(4)} = \begin{pmatrix} 1.7 \\ 1.0 \\ 4.2 \\ 0.8 \end{pmatrix}, \quad x^{(8)} = \begin{pmatrix} -1.69 \\ -1.15 \\ 5.50 \\ -0.40 \end{pmatrix}, \quad x^{(12)} = \begin{pmatrix} -2.63 \\ -1.76 \\ 5.86 \\ -0.85 \end{pmatrix}.$$

Method converges slowly to $x^{(\infty)} = (-3, -2, 6, -1)^T$.

33 / 77

Successive Over-Relaxation Method

Successive Over-Relaxation Use slightly larger update in the Gauss-Seidel method:

$$\begin{aligned} x'_i &= x_i - \omega \frac{\sum_{j<i} a_{ij}x'_j + \sum_{j>i} a_{ij}x_j - b_i}{a_{ii}} \\ &= (1 - \omega)x_i + \omega \frac{b_i - \sum_{j<i} a_{ij}x'_j - \sum_{j>i} a_{ij}x_j}{a_{ii}}. \end{aligned}$$

with $\omega \gtrsim 1$.

Typically ω is taken to be in the range $1.1 \leq \omega \leq 1.3$.

Implement in Matlab as:

```
for i=1:n, x(i)=x(i)-omega*(A(i,:)*x-b(i))/A(i,i); end;
```

Explicitly in matrix form,

$$x' = x - \omega (\omega L + D)^{-1} (Ax - b).$$

34 / 77

Successive Over-Relaxation Method

Example Solve $Ax = b$ using the successive over-relaxation method with

$$A = \begin{pmatrix} 6 & 2 & 0 \\ 3 & 5 & -1 \\ -2 & 1 & 4 \end{pmatrix}, \quad b = \begin{pmatrix} 3 \\ 4 \\ 1 \end{pmatrix}, \quad x^{(0)} = 0, \quad \omega = 1.1.$$

First step gives $x^{(1)} = (0.5500, 0.5170, 0.4353)^T$. Second step:

$$\begin{aligned} x_1^{(2)} &= (1 - \omega)x_1^{(1)} + \omega(b_1 - a_{12}x_2^{(1)} - a_{13}x_3^{(1)})/a_{11} \\ &= -0.1 \times 0.5500 + 1.1 \times (3 - 2 \times 0.5170 - 0 \times 0.4353)/6 = 0.3054; \\ x_2^{(2)} &= (1 - \omega)x_2^{(1)} + \omega(b_2 - a_{21}x_1^{(2)} - a_{23}x_3^{(1)})/a_{22} \\ &= -0.1 \times 0.5170 + 1.1 \times (4 - 3 \times 0.3054 - (-1) \times 0.4353 - 4)/5 = 0.7225; \\ x_3^{(2)} &= (1 - \omega)x_3^{(1)} + \omega(b_3 - a_{31}x_1^{(2)} - a_{32}x_2^{(2)})/a_{33} \\ &= -0.1 \times 0.4353 + 1.1 \times (1 - (-2) \times 0.3054 - 1 \times 0.7225)/4 = 0.2008. \end{aligned}$$

Further iterates give:

$$x^{(3)} = \begin{pmatrix} 0.25455 \\ 0.68392 \\ 0.20684 \end{pmatrix}, \quad x^{(4)} = \begin{pmatrix} 0.27377 \\ 0.67642 \\ 0.21888 \end{pmatrix}, \quad x^{(5)} = \begin{pmatrix} 0.27460 \\ 0.67927 \\ 0.21734 \end{pmatrix}, \quad x^{(\infty)} = \begin{pmatrix} 0.27358 \\ 0.67925 \\ 0.21698 \end{pmatrix}.$$

35 / 77

Conjugate-Gradient

36 / 77

Conjugate-Gradient Method

Method for solving $Ax = b$ where A is positive-definite.

Idea Construct sequence x_k minimising residual $\|Ax_k - b\|_2$ in $\text{span}\{b, Ab, \dots, A^{k-1}b\}$.

Use *inner products*

$$\langle u, S, v \rangle = \sum_{i,j=1}^n u_i S_{ij} v_j \text{ and } \langle u, v \rangle = \langle u, I, v \rangle = \sum_i^n u_i v_i.$$

Algorithm

Initialise

$$x_0 = 0, \quad r_0 = b - Ax_0, \quad v_1 = r_0;$$

Iterate for $k = 1, 2, \dots$

$$s_k = \langle r_{k-1}, r_{k-1} \rangle / \langle r_{k-2}, r_{k-2} \rangle, \quad s_1 \text{ unused.}$$

$$v_k = r_{k-1} + s_k v_{k-1}, \quad v_1 = r_0;$$

$$t_k = \langle r_{k-1}, r_{k-1} \rangle / \langle v_k, A, v_k \rangle,$$

$$x_k = x_{k-1} + t_k v_k,$$

$$r_k = r_{k-1} - A t_k v_k,$$

Note $r_k = b - Ax_k$ and $\langle v_i, A, v_j \rangle = 0$ for $i \neq j$.

37 / 77

Conjugate-Gradient Method

Example Solve $Ax = b$ using the conjugate-gradient method, where

$$A = \begin{pmatrix} 6 & 3 & -1 \\ 3 & 5 & 2 \\ -1 & 2 & 4 \end{pmatrix}, \quad b = \begin{pmatrix} 3 \\ 4 \\ 1 \end{pmatrix}.$$

$$x_0 = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}, \quad r_0 = b = \begin{pmatrix} 3 \\ 4 \\ 1 \end{pmatrix};$$

$$v_1 = r_0 = \begin{pmatrix} 3 \\ 4 \\ 1 \end{pmatrix}$$

$$t_1 = \frac{\langle r_0, r_0 \rangle}{\langle v_1, Av_1 \rangle} = \frac{26}{220} = 0.11818$$

$$x_1 = x_0 + t_1 v_1 = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix} + 0.11818 \times \begin{pmatrix} 3 \\ 4 \\ 1 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix} + \begin{pmatrix} 0.35455 \\ 0.47273 \\ 0.11818 \end{pmatrix} = \begin{pmatrix} 0.35455 \\ 0.47273 \\ 0.11818 \end{pmatrix}$$

$$r_1 = r_0 - At_1 v_1 = \begin{pmatrix} 3 \\ 4 \\ 1 \end{pmatrix} - \begin{pmatrix} 6 & 3 & -1 \\ 3 & 5 & 2 \\ -1 & 2 & 4 \end{pmatrix} \times \begin{pmatrix} 0.35455 \\ 0.47273 \\ 0.11818 \end{pmatrix} = \begin{pmatrix} -0.427273 \\ 0.336364 \\ -0.063636 \end{pmatrix}$$

38 / 77

Conjugate-Gradient Method

Example

$$v_1 = \begin{pmatrix} 3.00000 \\ 4.00000 \\ 1.00000 \end{pmatrix}, \quad x_1 = \begin{pmatrix} 0.35455 \\ 0.47273 \\ 0.11818 \end{pmatrix}, \quad r_1 = \begin{pmatrix} -0.427273 \\ 0.336364 \\ -0.063636 \end{pmatrix}.$$

$$s_2 = \frac{\langle r_1, r_1 \rangle}{\langle r_0, r_0 \rangle} = \frac{0.29975}{26.00000} = 0.011529$$

$$v_2 = r_1 + s_2 v_1 = \begin{pmatrix} -0.427273 \\ 0.336364 \\ -0.063636 \end{pmatrix} + 0.011529 \times \begin{pmatrix} 3.00000 \\ 4.00000 \\ 1.00000 \end{pmatrix} = \begin{pmatrix} -0.39269 \\ 0.38248 \\ -0.05211 \end{pmatrix}$$

$$t_2 = \frac{\langle r_1, r_1 \rangle}{\langle v_2, Av_2 \rangle} = \frac{0.29975}{0.64572} = 0.46422$$

$$x_2 = x_1 + t_2 v_2 = \begin{pmatrix} 0.355 \\ 0.473 \\ 0.118 \end{pmatrix} + 0.464 \times \begin{pmatrix} -0.393 \\ 0.382 \\ -0.052 \end{pmatrix} = \begin{pmatrix} 0.355 \\ 0.473 \\ 0.118 \end{pmatrix} + \begin{pmatrix} -0.1823 \\ 0.1776 \\ -0.0242 \end{pmatrix} = \begin{pmatrix} 0.17225 \\ 0.65028 \\ 0.09399 \end{pmatrix}$$

$$r_2 = r_1 - At_2 v_2 = \begin{pmatrix} -0.4273 \\ 0.3364 \\ -0.0636 \end{pmatrix} - \begin{pmatrix} 6 & 3 & -1 \\ 3 & 5 & 2 \\ -1 & 2 & 4 \end{pmatrix} \times \begin{pmatrix} -0.1823 \\ 0.1776 \\ -0.0242 \end{pmatrix} = \begin{pmatrix} 0.109625 \\ 0.043850 \\ -0.504277 \end{pmatrix}$$

39 / 77

Conjugate-Gradient Method

Example

$$v_2 = \begin{pmatrix} -0.39269 \\ 0.38248 \\ -0.05211 \end{pmatrix}, \quad x_2 = \begin{pmatrix} 0.17225 \\ 0.65028 \\ 0.09399 \end{pmatrix}, \quad r_2 = \begin{pmatrix} 0.109625 \\ 0.043850 \\ -0.504277 \end{pmatrix}.$$

$$s_3 = \frac{\langle r_2, r_2 \rangle}{\langle r_1, r_1 \rangle} = \frac{0.26824}{0.29975} = 0.89486$$

$$v_3 = r_2 + s_3 v_2 = \begin{pmatrix} -0.24177 \\ 0.38612 \\ -0.55091 \end{pmatrix}$$

$$t_3 = \frac{\langle r_2, r_2 \rangle}{\langle v_3, A v_3 \rangle} = \frac{0.26824}{0.63278} = 0.42390$$

$$\text{Solution } x = x_3 = x_2 + t_3 v_3 = \begin{pmatrix} 0.06977 \\ 0.81395 \\ -0.13953 \end{pmatrix}$$

$$r_3 = r_2 - A t_3 v_3 = \begin{pmatrix} 0.0 \\ 0.0 \\ 0.0 \end{pmatrix} = b - Ax \text{ (residual)}$$

40 / 77

Conjugate-Gradient Method

Convergence The conjugate-gradient method yields the exact result (up to roundoff errors) for an $n \times n$ system after n steps.

Termination Often, good accuracy is achieved after $m \ll n$ steps! This can be seen if the residual r is small.

Sparse systems If A is sparse, with $p \ll n^2$ nonzero elements, then each step takes $O(p)$ operations.

Efficiency For many practical problems, the conjugate-gradient method only needs $O(mp)$ steps for good accuracy, much less than $O(n^3)$ for Gaussian elimination!

41 / 77

Preconditioned Conjugate-Gradient (Non-examinable)

Preconditioning Choose a *preconditioning* matrix P .

Apply the conjugate-gradient method to the equations

$$(PAP^T)y = Pb$$

and set

$$x = P^T y.$$

A typical choice is $P = (\text{diag}(A))^{1/2}$.

42 / 77

Eigenvalues and eigenvectors

Eigenvalues and Eigenvectors If $Av = \lambda v$ and $v \neq 0$, then λ is an *eigenvalue* of A with corresponding eigenvector v .

Similarity Matrices A and B are *similar* if $B = PAP^{-1}$ for some invertible matrix P .

If $Av = \lambda v$, then $B(Pv) = (PAP^{-1})Pv = PA v = P(\lambda v) = \lambda(Pv)$, so Pv is an eigenvector of B with eigenvalue λ .

Triangular If A is a lower- or upper-triangular matrix, then the eigenvalues of A are the diagonal entries;
 $\lambda_i = a_{ii}$

Diagonal If D is a diagonal matrix, then the eigenvectors are the standard unit basis vectors e_i , with
 $De_i = d_{ii}e_i$.

Notation Often write eigenvalues in order, $|\lambda_1| \geq |\lambda_2| \geq \dots \geq |\lambda_n|$, with corresponding eigenvectors v_i .

44 / 77

Approximation theorems

Gersgorin Circle Theorem For $i = 1, \dots, n$, there exists an eigenvalue λ_i of A within the circle

$$\{z \in \mathbb{C} \mid |z - a_{ii}| \leq \sum_{j=1, j \neq i}^n |a_{ij}|\}.$$

Similarly, for $j = 1, \dots, n$, there exists an eigenvalue λ_j of A in the circle

$$\{z \in \mathbb{C} \mid |z - a_{jj}| \leq \sum_{i=1, i \neq j}^n |a_{ij}|\}.$$

Example

$$A = \begin{pmatrix} 10 & -2 & 1 \\ 1 & 3 & -1 \\ 0 & 1 & -2 \end{pmatrix}.$$

The eigenvalues $\lambda_{1,2,3}$ satisfy:

$$|\lambda_1 - 10| \leq |-2| + |1| = 3, \quad |\lambda_2 - 3| \leq |1| + |-1| = 2, \quad |\lambda_3 - (-2)| \leq 1.$$

If $\lambda_{1,2,3}$ are real, then $\lambda_1 \in [7, 13]$, $\lambda_2 \in [1, 5]$, $\lambda_3 \in [-3, -1]$.

By considering the first column, see $|\lambda_1 - 10| \leq 1$.

45 / 77

The power method

Power method Iterate

$$y^{(n)} = Ax^{(n)}, \quad x^{(n+1)} = y^{(n)} / \pm \|y^{(n)}\|.$$

Note $x^{(n)} = \pm A^n x^{(0)} / \|A^n x^{(0)}\|$.

Although any norm can be used, typically use supremum norm $\|y\|_\infty$, choosing sign so that maximum absolute value of x is one:

$$x^{(n+1)} = y^{(n)} / y_{i_{\text{abs max}}}^{(n)} \quad \text{where } |y_{i_{\text{abs max}}}^{(n)}| \geq |y_i^{(n)}| \text{ for all } i.$$

Take eigenvalue approximation

$$\mu^{(n)} = (Ax^{(n)})_{i_{\text{abs max}}} / x_{i_{\text{abs max}}}^{(n)} = y_{i_{\text{abs max}}}^{(n)} / x_{i_{\text{abs max}}}^{(n)}.$$

It is usually more accurate, notably for symmetric matrices, to take the alternative eigenvalue approximation

$$\mu^{(n)} = \frac{x^{(n)T} Ax^{(n)}}{x^{(n)T} x^{(n)}} = \frac{x^{(n)} \cdot y^{(n)}}{x^{(n)} \cdot x^{(n)}}.$$

If the Euclidean norm is used, then $\|x^{(n)}\|_2 = 1$, and the formula reduces to

$$\mu^{(n)} = x^{(n)T} Ax^{(n)} = x^{(n)} \cdot y^{(n)}.$$

46 / 77

The power method

Theorem If A has an eigenvalue λ_1 such that $|\lambda_1| > |\lambda_i|$ for all other eigenvalues, then the power method converges, with $\lim_{n \rightarrow \infty} x^{(n)} = v_1$ and $\lim_{n \rightarrow \infty} \mu^{(n)} = \lambda_1$.

Proof.

For simplicity, suppose \mathbb{R}^n has basis $\{v_1, v_2, \dots, v_n\}$ of eigenvectors of A .

Write $x = \alpha_1 v_1 + \alpha_2 v_2 + \dots + \alpha_n v_n$. Then

$$\begin{aligned} A^k x &= A^k (\alpha_1 v_1 + \alpha_2 v_2 + \dots + \alpha_n v_n) \\ &= \alpha_1 A^k v_1 + \alpha_2 A^k v_2 + \dots + \alpha_n A^k v_n \\ &= \alpha_1 \lambda_1^k v_1 + \alpha_2 \lambda_2^k v_2 + \dots + \alpha_n \lambda_n^k v_n \\ &= \lambda_1^k (\alpha_1 v_1 + \alpha_2 (\lambda_2 / \lambda_1)^k v_2 + \dots + \alpha_n (\lambda_n / \lambda_1)^k v_n). \end{aligned}$$

Hence $\lim_{k \rightarrow \infty} A^k x / \lambda_1^k = \alpha_1 v_1$, so

$$\lim_{k \rightarrow \infty} x^{(k)} = \lim_{k \rightarrow \infty} \frac{A^k x}{\pm \|A^k x\|} = \lim_{k \rightarrow \infty} \pm \frac{A^k x / \lambda_1^k}{\|A^k x / \lambda_1^k\|} = \pm \frac{\alpha_1 v_1}{\|\alpha_1 v_1\|} = \pm \hat{v}_1.$$

47 / 77

The power method

Example

$$A = \begin{pmatrix} 1 & 2 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix}, \quad x = \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix}.$$

$$y^{(0)} = Ax^{(0)} = \begin{pmatrix} 1 & 2 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix} \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} = \begin{pmatrix} 3 \\ 1 \\ 1 \end{pmatrix}; \quad x^{(1)} = \frac{y^{(0)}}{\|y^{(0)}\|_{\infty}} = \begin{pmatrix} 1 \\ 1/3 \\ 1/3 \end{pmatrix} = \begin{pmatrix} 1.0000 \\ 0.3333 \\ 0.3333 \end{pmatrix}$$

$$y^{(1)} = Ax^{(1)} = \begin{pmatrix} 1 & 2 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix} \begin{pmatrix} 1.0000 \\ 0.3333 \\ 0.3333 \end{pmatrix} = \begin{pmatrix} 1.6667 \\ 0.3333 \\ 1.0000 \end{pmatrix}; \quad x^{(2)} = \frac{y^{(1)}}{\|y^{(1)}\|_{\infty}} = \begin{pmatrix} 1.0000 \\ 0.2000 \\ 0.6000 \end{pmatrix}$$

$$y^{(2)} = Ax^{(2)} = \begin{pmatrix} 1 & 2 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix} \begin{pmatrix} 1.0000 \\ 0.2000 \\ 0.6000 \end{pmatrix} = \begin{pmatrix} 1.4000 \\ 0.6000 \\ 1.0000 \end{pmatrix}; \quad x^{(3)} = \frac{y^{(2)}}{\|y^{(2)}\|_{\infty}} = \begin{pmatrix} 1.0000 \\ 0.4286 \\ 0.7143 \end{pmatrix}$$

$$y^{(3)} = Ax^{(3)} = \begin{pmatrix} 1 & 2 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix} \begin{pmatrix} 1.0000 \\ 0.4286 \\ 0.7143 \end{pmatrix} = \begin{pmatrix} 1.8571 \\ 0.7143 \\ 1.0000 \end{pmatrix}; \quad x^{(4)} = \frac{y^{(3)}}{\|y^{(3)}\|_{\infty}} = \begin{pmatrix} 1.0000 \\ 0.3846 \\ 0.5385 \end{pmatrix}$$

48 / 77

The power method

Example

$$y^{(4)} = Ax^{(4)} = \begin{pmatrix} 1 & 2 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix} \begin{pmatrix} 1.0000 \\ 0.3846 \\ 0.5385 \end{pmatrix} = \begin{pmatrix} 1.7692 \\ 0.5385 \\ 1.0000 \end{pmatrix}; \quad x^{(5)} = \frac{y^{(4)}}{\|y^{(4)}\|_{\infty}} = \begin{pmatrix} 1.0000 \\ 0.3043 \\ 0.5652 \end{pmatrix}$$

$$y^{(5)} = Ax^{(5)} = \begin{pmatrix} 1 & 2 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix} \begin{pmatrix} 1.0000 \\ 0.3846 \\ 0.5385 \end{pmatrix} = \begin{pmatrix} 1.6087 \\ 0.5652 \\ 1.0000 \end{pmatrix}.$$

Estimate

$$v \approx x^{(5)} = \begin{pmatrix} 1.0000 \\ 0.3043 \\ 0.5652 \end{pmatrix}; \quad \lambda \approx \mu^{(5)} = \frac{(Ax^{(5)})_{i_{\text{abs max}}}}{(x^{(5)})_{i_{\text{abs max}}}} = (Ax^{(5)})_1 = 1.6087.$$

Alternative estimate

$$v \approx \hat{x}^{(5)} = \frac{x^{(5)}}{\|x^{(5)}\|_2} = \begin{pmatrix} 0.8415 \\ 0.2561 \\ 0.4756 \end{pmatrix}; \quad \lambda \approx \hat{\mu}^{(5)} = \frac{x^{(5)T}Ax^{(5)}}{x^{(5)T}x^{(5)}} = \hat{x}^{(5)T}A\hat{x}^{(5)} = 1.6613.$$

Actual eigenvalue/vector $\lambda = 1.6841$, $v = (1.0000, 0.3514, 0.6216)^T$ or normalised $\hat{v} = (0.8138, 0.2859, 0.5059)^T$.

49 / 77

Inverse power method

Theorem If λ is an eigenvalue of A , then $1/(\lambda - \mu)$ is an eigenvalue of $(A - \mu I)^{-1}$ with the same eigenvector v . Hence if $(A - \mu I)^{-1}v = \kappa v$, then $1/(\lambda - \mu) = \kappa$, so $\lambda = \mu + 1/\kappa$.

Proof. $(A - \mu I)v = Av - \mu Iv = \lambda v - \mu v = (\lambda - \mu)v$,
so $(A - \mu I)^{-1}v = (\lambda - \mu)^{-1}v$. □

Inverse power method To estimate an eigenvalue $\lambda \approx \mu$, apply the power method to $(A - \mu I)^{-1}$.

Iterate

$$y^{(n)} = (A - \mu I)^{-1}x^{(n)}, \quad x^{(n+1)} = y^{(n)} / \pm \|y^{(n)}\|.$$

Estimate

$$\lambda \approx \lambda^{(n)} = \mu + 1/\kappa^{(n)} \text{ where } \kappa^{(n)} = y_i^{(n)} / x_i^{(n)}.$$

In practise, first compute LU-factorisation $A - \mu I = L_\mu U_\mu$, and solve

$$(A - \mu I)y^{(n)} = L_\mu U_\mu y^{(n)} = x^{(n)}.$$

Can update μ at each step, setting $\mu^{(n+1)} = \mu^{(n)} + 1/\kappa^{(n)}$. This speeds up convergence, but requires an expensive matrix factorisation at each step.

50 / 77

Inverse power method

Example

$$A = \begin{pmatrix} 2 & -1 & 1 \\ -1 & 3 & -2 \\ 1 & 2 & 3 \end{pmatrix}; \quad \mu = 1.5. \quad \text{Use } x^{(0)} = \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}.$$

$$A - \mu I = \begin{pmatrix} 0.5 & -1 & 1 \\ -1 & 1.5 & -2 \\ 1 & 2 & 1.5 \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 \\ -2.0 & 1 & 0 \\ 2.0 & -8.0 & 1 \end{pmatrix} \begin{pmatrix} 0.5 & -1.0 & 1.0 \\ 0 & -0.5 & 0.0 \\ 0 & 0 & -0.5 \end{pmatrix} = L_\mu U_\mu$$

$$(A - \mu I)^{-1} = \begin{pmatrix} 0.5 & -1 & 1 \\ -1 & 1.5 & -2 \\ 1 & 2 & 1.5 \end{pmatrix}^{-1} = \begin{pmatrix} 50 & 28 & 4 \\ -4 & -2 & 0 \\ -28 & -16 & -2 \end{pmatrix}$$

$$y^{(0)} = (A - \mu I)^{-1}x^{(0)} = \begin{pmatrix} 50.0000 \\ -4.0000 \\ -28.0000 \end{pmatrix}; \quad x^{(1)} = \frac{y^{(0)}}{y_{i_{\text{abs max}}}^{(0)}} = \begin{pmatrix} 1.0000 \\ -0.0800 \\ -0.5600 \end{pmatrix}$$

$$y^{(1)} = (A - \mu I)^{-1}x^{(1)} = \begin{pmatrix} 45.5200 \\ -3.8400 \\ -25.6000 \end{pmatrix}; \quad x^{(2)} = \frac{y^{(1)}}{y_{i_{\text{abs max}}}^{(1)}} = \begin{pmatrix} 1.0000 \\ -0.0844 \\ -0.5624 \end{pmatrix}$$

$$y^{(2)} = (A - \mu I)^{-1}x^{(2)} = \begin{pmatrix} 45.3884 \\ -3.8313 \\ -25.5255 \end{pmatrix}; \quad x^{(3)} = \frac{y^{(2)}}{y_{i_{\text{abs max}}}^{(2)}} = \begin{pmatrix} 1.0000 \\ -0.0844 \\ -0.5624 \end{pmatrix}$$

Eigenvalue/vector $\lambda \approx (Ax^{(3)})_1 = 1.5220$, $v \approx (1.0000, -0.0844, -0.5624)$.

51 / 77

Deflation (Off-syllabus)

Deflation Suppose A has eigenvalue/vector pairs (λ_i, v_i) .

If x is a vector such that $x^T v_1 = 1$, and $B = A - \lambda_1 v_1 x^T$,

then B has eigenvalues 0 and $\lambda_i - \lambda_1$ for $i = 2, \dots, n$

with eigenvectors w_i satisfying $v_i = (\lambda_i - \lambda_1)w_i + \lambda_1(x^T w_i)v_1$.

Wielandt deflation Define x by $x_j = a_{k,j}/(\lambda_1(v_1)_k)$ for some k . Then the k^{th} row of B is identically 0, so $(w_i)_k = 0$ for all $i = 2, \dots, n$.

Orthogonalisation

53 / 77

Orthogonality

Normal vectors A vector v is *normal* if $\|v\|_2 := \sqrt{\sum_{i=1}^n v_i^2} = 1$, equivalently if $v \cdot v = 1$.

Orthogonal vectors Vectors $\{v_1, \dots, v_n\}$ are *orthogonal* if $v_i \cdot v_j = 0$ for all $i \neq j$.

Orthonormal vectors Vectors $\{v_1, \dots, v_n\}$ are *orthonormal* if they are orthogonal and each is normal.

Orthogonal matrices A matrix Q is *orthogonal* if $Q^{-1} = Q^T$; equivalently, if the columns of Q are orthonormal vectors.

54 / 77

Gram-Schmidt orthogonalisation

Gram-Schmidt orthogonalisation Let x_1, \dots, x_m be vectors in \mathbb{R}^m .

Recursively define

$$v_i = x_i - \sum_{j=1}^{i-1} \frac{x_i \cdot v_j}{v_j \cdot v_j} v_j = x_i - \sum_{j=1}^{i-1} (x_i \cdot u_j) u_j; \quad u_i = \frac{v_i}{\|v_i\|_2}.$$

Theorem The v_i are an orthogonal set, and the u_i are an orthonormal set, such that for all $k = 1, \dots, m$,
 $\text{span}\{x_1, \dots, x_k\} = \text{span}\{u_1, \dots, u_k\} = \text{span}\{v_1, \dots, v_k\}.$

Proof. Fix i and assume $\{u_1, \dots, u_{i-1}\}$ are orthonormal. Then for $i > j$,

$$v_i \cdot u_j = x_i \cdot u_j - \sum_{k=1}^{i-1} (x_i \cdot u_k) u_k \cdot u_j = x_i \cdot u_j - (x_i \cdot u_j)(u_j \cdot u_j) = 0,$$

so $u_i \cdot u_j = (v_i / \|v_i\|_2) \cdot u_j = (v_i \cdot u_j) / \|v_i\|_2 = 0.$ □

55 / 77

Gram-Schmidt orthogonalisation

Example Apply the Gram-Schmidt orthogonalisation procedure to:

$$x_1 = \begin{pmatrix} 2 \\ -1 \\ 1 \end{pmatrix}, \quad x_2 = \begin{pmatrix} -1 \\ 3 \\ 2 \end{pmatrix}, \quad x_3 = \begin{pmatrix} 1 \\ -2 \\ 3 \end{pmatrix}.$$

$$v_1 = x_1 = \begin{pmatrix} 2 \\ -1 \\ 1 \end{pmatrix}; \quad v_2 = x_2 - \frac{x_2 \cdot v_1}{v_1 \cdot v_1} v_1 = \begin{pmatrix} -1 \\ 3 \\ 2 \end{pmatrix} - \frac{-3}{6} \begin{pmatrix} 2 \\ -1 \\ 1 \end{pmatrix} = \begin{pmatrix} 0 \\ 5/2 \\ 5/2 \end{pmatrix};$$

$$v_3 = x_3 - \frac{x_3 \cdot v_1}{v_1 \cdot v_1} v_1 - \frac{x_3 \cdot v_2}{v_2 \cdot v_2} v_2 = \begin{pmatrix} 1 \\ -2 \\ 3 \end{pmatrix} - \frac{7}{6} \begin{pmatrix} 2 \\ -1 \\ 1 \end{pmatrix} - \frac{5/2}{25/2} \begin{pmatrix} 0 \\ 5/2 \\ 5/2 \end{pmatrix} = \begin{pmatrix} -4/3 \\ -4/3 \\ 4/3 \end{pmatrix}.$$

$$\|v_1\|_2 = \sqrt{6}; \quad \|v_2\|_2 = 5/\sqrt{2}; \quad \|v_3\|_2 = 4/\sqrt{3}.$$

$$u_1 = \frac{v_1}{\|v_1\|_2} = \frac{1}{\sqrt{6}} \begin{pmatrix} 2 \\ -1 \\ 1 \end{pmatrix}; \quad u_2 = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 \\ 1 \\ 1 \end{pmatrix}; \quad u_3 = \frac{1}{\sqrt{3}} \begin{pmatrix} -1 \\ -1 \\ 1 \end{pmatrix}.$$

Check orthogonality:

$$u_1 \cdot u_2 = \frac{2 \times 0 + (-1) \times 1 + 1 \times 1}{\sqrt{6} \times \sqrt{2}} = 0; \quad u_1 \cdot u_3 = \frac{-2 + 1 + 1}{\sqrt{18}} = 0; \quad u_2 \cdot u_3 = 0.$$

QR factorisation

QR factorisation In the Gram-Schmidt orthogonalisation, define $r_{i,i} = \|v_i\|_2$ and $r_{i,j} = x_j \cdot u_i = x_j \cdot v_i / \|v_i\|_2$. Then

$$v_j = x_j - \sum_{i=1}^{j-1} r_{i,j} u_i, \text{ and } u_i = v_i / r_{i,i}$$

so

$$x_j = r_{j,j} u_j + \sum_{i=1}^{j-1} r_{i,j} u_i.$$

Let

$$A = (x_1, \dots, x_n), \quad Q = (u_1, \dots, u_n) \text{ and } (R)_{i,j} = r_{i,j} \text{ for } i \leq j.$$

Then Q is orthogonal, R is upper-triangular, and

$$A = QR.$$

QR factorisation

Example Compute the QR-factorisation of:

$$A = \begin{pmatrix} 2 & -1 & 1 \\ -1 & 3 & -2 \\ 1 & 2 & 3 \end{pmatrix}$$

From the Gram-Schmidt orthogonalisation,

$$Q = \begin{pmatrix} 2/\sqrt{6} & 0 & -1/\sqrt{3} \\ -1/\sqrt{6} & 1/\sqrt{2} & -1/\sqrt{3} \\ 1/\sqrt{6} & 1/\sqrt{2} & 1/\sqrt{3} \end{pmatrix} = \begin{pmatrix} 0.816 & 0.000 & -0.577 \\ -0.408 & 0.707 & -0.577 \\ 0.408 & 0.707 & 0.577 \end{pmatrix}$$

$$R = \begin{pmatrix} 6/\sqrt{6} & -3/\sqrt{6} & 7/\sqrt{6} \\ 0 & 5/\sqrt{2} & 1/\sqrt{2} \\ 0 & 0 & 4/\sqrt{3} \end{pmatrix} = \begin{pmatrix} 2.449 & -1.225 & 2.858 \\ 0 & 3.536 & 0.707 \\ 0 & 0 & 2.309 \end{pmatrix}.$$

The QR Method**The QR method**

The QR Method The QR method is an iterative algorithm for finding all the eigenvalues of A .

Set $A^{(0)} = A$. Iteratively find $Q^{(n)}, R^{(n)}$ such that $A^{(n)} = Q^{(n)} R^{(n)}$ and set $A^{(n+1)} = R^{(n)} Q^{(n)}$.

Theorem Assuming eigenvalues of A have distinct absolute value, $A^{(n)}$ converges to an upper-triangular matrix with the eigenvalues of A on the diagonal.

The QR method

Example Use the QR method to approximate eigenvalues of:

$$A = \begin{pmatrix} 4 & -1 & 1 \\ -1 & 3 & -2 \\ 1 & -2 & 3 \end{pmatrix}.$$

$$Q^{(0)} = \begin{pmatrix} -0.9428 & -0.3244 & -0.0765 \\ 0.2357 & -0.8111 & 0.5353 \\ -0.2357 & 0.4867 & 0.8412 \end{pmatrix}, \quad R^{(0)} = \begin{pmatrix} -4.2426 & 2.1213 & -2.1213 \\ 0 & -3.0822 & 2.7578 \\ 0 & 0 & 1.3765 \end{pmatrix}$$

$$A^{(1)} = R^{(0)}Q^{(0)} = \begin{pmatrix} 5.0000 & -1.3765 & -0.3244 \\ -1.3765 & 3.8421 & 0.6699 \\ -0.3244 & 0.6699 & 1.1579 \end{pmatrix}$$

$$A^{(2)} = R^{(1)}Q^{(1)} = \begin{pmatrix} 5.6667 & -0.9406 & 0.0640 \\ -0.9406 & 3.3226 & -0.1580 \\ 0.0640 & -0.1580 & 1.0108 \end{pmatrix}$$

$$A^{(3)} = \begin{pmatrix} 5.909 & -0.514 & -0.011 \\ -0.514 & 3.090 & 0.045 \\ -0.011 & 0.045 & 1.001 \end{pmatrix} \quad A^{(4)} = \begin{pmatrix} 5.977 & -0.263 & 0.002 \\ -0.263 & 3.023 & -0.014 \\ 0.002 & -0.014 & 1.000 \end{pmatrix}$$

Hence $\lambda_1 = 5.977 \pm 0.265$, $\lambda_2 = 3.023 \pm 0.277$, $\lambda_3 = 1.000 \pm 0.016$.

61 / 77

The QR method—Properties (Non-examinable)

Conjugacies Note

$$A^{(k)} = Q^{(k)}R^{(k)} = R^{(k-1)}Q^{(k-1)};$$

$$Q^{(k)}A^{(k+1)} = Q^{(k)}R^{(k)}Q^{(k)} = A^{(k)}Q^{(k)}.$$

Similarity Set $P^{(k)} = Q^{(0)}Q^{(1)} \dots Q^{(k-1)}$. Then

$$P^{(k)}A^{(k)} = AP^{(k)}.$$

Equivalently,

$$A = P^{(k)}A^{(k)}(P^{(k)})^{-1}$$

Power Set $S^{(k)} = R^{(k)} \dots R^{(1)}R^{(0)}$. Then

$$A^k = P^{(k)}S^{(k)}.$$

62 / 77

The QR method—Convergence (Non-examinable)

Convergence Since $P^{(k)}$ is orthogonal and $S^{(k)}$ upper-triangular, we have

$$A^k e_1 = P^{(k)} S^{(k)} e_1 = s_{1,1}^{(k)} P^{(k)} e_1,$$

$$A^k e_2 = P^{(k)} S^{(k)} e_2 = P^{(k)} (s_{1,2}^{(k)} e_1 + s_{2,2}^{(k)} e_2) = s_{1,2}^{(k)} P^{(k)} e_1 + s_{2,2}^{(k)} P^{(k)} e_2.$$

We deduce

$$P^{(k)} e_i \in \text{span}\{A^k e_1, \dots, A^k e_i\}$$

and is orthogonal to $\{A^k e_1, \dots, A^k e_{i-1}\}$.

Writing $e_i = \sum \alpha_{i,j} v_j$ gives

$$A^k e_i = \sum \alpha_{i,j} \lambda_j^k v_j.$$

Then

$$P^{(k)} e_1 \propto A^k e_1 = \alpha_{1,1} \lambda_1^k (v_1 + \sum (\alpha_{1,j}/\alpha_{1,1}) (\lambda_j/\lambda_1)^k v_j) \sim v_1 \text{ as } k \rightarrow \infty.$$

Similarly, we see that

$$\lim_{k \rightarrow \infty} P^{(k)} e_i \in \text{span}\{v_1, v_2, \dots, v_i\}$$

and is orthogonal to $\{v_1, \dots, v_{i-1}\}$.

63 / 77

The QR method (Non-examinable)

Shifted QR The QR method can be *shifted* i.e. applied to $A^{(n)} - \mu^{(n)} I$ for scalars $\mu^{(n)}$, as

$$A^{(n)} - \mu^{(n)} I = Q^{(n)} R^{(n)}; \quad A^{(n+1)} = R^{(n)} Q^{(n)} + \mu^{(n)} I.$$

This can accelerate convergence, especially when eigenvalues have nearly the same absolute value.

Partial QR The QR method can also be modified to find only *some* of the eigenvalues of A by iterating $Q^{(k)} R^{(k)} = A Q^{(k-1)}$, where the $Q^{(k)}$ are n -by- m matrices with orthonormal columns, and the $R^{(k)}$ are m -by- m upper-triangular matrices.

64 / 77

Householder matrices (Advanced)

Householder matrices A Householder matrix is a matrix of the form

$$H = I - 2 \frac{v v^T}{v^T v}$$

or

$$H = I - 2 w w^T \text{ where } \|w\|_2 = 1.$$

Symmetry A Householder matrix is symmetric, since

$$H^T = (I - 2 w w^T)^T = I^T - 2 (w w^T)^T = I^T - 2 (w^T)^T w^T = I - 2 w w^T.$$

Orthogonality A Householder matrix is orthogonal, since

$$\begin{aligned} H^T H &= H H = (I - 2 w w^T)(I - 2 w w^T) \\ &= I - 2 w w^T - 2 w w^T + 4 w w^T w w^T = I - 4 w w^T + 4 w (w^T w) w^T = I \end{aligned}$$

Theorem If $H = I - 2 w w^T$ is a Householder matrix, then $H = H^T = H^{-1}$.

65 / 77

Householder matrices (Advanced)

Example Take $v = (0, 2, -1, 1)^T$, $w = v/\sqrt{6}$,

$$H = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & -\frac{1}{3} & \frac{2}{3} & -\frac{2}{3} \\ 0 & \frac{2}{3} & \frac{2}{3} & \frac{1}{3} \\ 0 & -\frac{2}{3} & \frac{1}{3} & \frac{2}{3} \end{pmatrix} = \frac{1}{3} \begin{pmatrix} 3 & 0 & 0 & 0 \\ 0 & -1 & 2 & -2 \\ 0 & 2 & 2 & 1 \\ 0 & -2 & 1 & 2 \end{pmatrix}$$

66 / 77

Householder matrices (Advanced)

Upper-Hessenberg form For a matrix A , aim to find an orthogonal matrix Q such that $Q^T A Q$ has *upper-Hessenberg* form $A_{i,j} = 0$ for $i > j + 1$.

$$\begin{pmatrix} * & * & * & * & * \\ * & * & * & * & * \\ 0 & * & * & * & * \\ 0 & 0 & * & * & * \\ 0 & 0 & 0 & * & * \end{pmatrix}$$

Note that if A is symmetric and upper-Hessenberg, then it is *tridiagonal* $A_{i,j} = 0$ for $|i - j| > 1$.

$$\begin{pmatrix} * & * & 0 & 0 & 0 \\ * & * & * & 0 & 0 \\ 0 & * & * & * & 0 \\ 0 & 0 & * & * & * \\ 0 & 0 & 0 & * & * \end{pmatrix}$$

Upper-Hessenberg form greatly increases the efficiency of the QR method; time $O(n^2)$ per step instead of $O(n^3)$, and $O(n)$ for symmetric matrices.

67 / 77

Householder matrices (Advanced)

Conversion to upper-Hessenberg form First find a Householder matrix $H = H_1$ such that $H_1^T A H_1$ has first column $(H_1^T A H_1)_{i,1} = 0$ for $i \geq 3$:

1. Set $\alpha = (\sum_{i=2}^n a_{i,1}^2)^{1/2}$, $v_1 = 0, v_2 = a_{2,1} \pm \alpha, v_i = a_{i,1}$ for $i \geq 3$.
Typically choose sign so that $v_2 = a_{2,1} + \text{sgn}(a_{2,1})\alpha$.
2. Take $w = v/r$ where $r = \|v\|_2 = (2\alpha(\alpha \pm a_{2,1}))^{1/2}$.
3. Set $H = I - 2ww^T$, so $H_{i,1} = 0$ for $i > 2$.

Continue by applying the method to the sub-matrix $(H_1^T A H_1)_{2:n,2:n}$ to find a Householder matrix H_2 such that $H_2^T H_1^T A H_1 H_2$ has $H_{i,j} = 0$ for $j = 1, 2$ and $i > j + 1$.

Note that in practise, we compute

$$H A = (I - 2ww^T)A = A - (2w)(w^T A)$$

which takes $O(n^2)$ operations, whereas computing H first and then $H A$ would take $O(n^3)$ operations.

68 / 77

Householder matrices (Advanced)

Example

$$A = \begin{pmatrix} 4 & 1 & -2 & 2 \\ 1 & 2 & 0 & 1 \\ -2 & 0 & 3 & -2 \\ 2 & 1 & -2 & -1 \end{pmatrix}.$$

$$\alpha_1 = -\operatorname{sgn}(a_{21})(\sum_{i=2}^4 a_{2i}^2)^{1/2} = -\sqrt{9} = -3.$$

$$r_1 = (\frac{1}{2}\alpha_1(\alpha_1 - a_{2,1}))^{1/2} = (-\frac{1}{2}\alpha_1(a_{2,1} - \alpha_1))^{1/2} = \sqrt{\frac{1}{2}3(3+1)} = \sqrt{6}.$$

$$v_1 = \begin{pmatrix} 0 \\ 1 - (-3) \\ -2 \\ 2 \end{pmatrix} = \begin{pmatrix} 0 \\ 4 \\ -2 \\ 2 \end{pmatrix}; \quad w_1 = v_1/2r_1 = \frac{1}{\sqrt{6}} \begin{pmatrix} 0 \\ 2 \\ -1 \\ 1 \end{pmatrix}.$$

$$H_1 = I - 2w_1w_1^T = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & -\frac{1}{3} & \frac{2}{3} & -\frac{2}{3} \\ 0 & \frac{2}{3} & \frac{1}{3} & \frac{1}{3} \\ 0 & -\frac{2}{3} & \frac{1}{3} & \frac{1}{3} \end{pmatrix} = \frac{1}{3} \begin{pmatrix} 3 & 0 & 0 & 0 \\ 0 & -1 & 2 & -2 \\ 0 & 2 & 1 & 1 \\ 0 & -2 & 1 & 1 \end{pmatrix}$$

69 / 77

Householder matrices (Advanced)

Example (continued)

$$H_1^T A H_1 = \frac{1}{9} \begin{pmatrix} 36 & -27 & 0 & 0 \\ -27 & 30 & 9 & 12 \\ 0 & 9 & 15 & -12 \\ 0 & 12 & -12 & -9 \end{pmatrix} = \begin{pmatrix} 4 & -3 & 0 & 0 \\ -3 & \frac{10}{3} & 1 & \frac{4}{3} \\ 0 & 1 & \frac{5}{3} & -\frac{4}{3} \\ 0 & \frac{4}{3} & -\frac{4}{3} & -1 \end{pmatrix}$$

$$\alpha_2 = \pm(\sum_{i=3}^4 a_{i,2}^2)^{1/2} = -\sqrt{1^2 + (\frac{4}{3})^2} = -\frac{5}{3}; \quad (v_2)_3 = a_{2,3} - \alpha_2 = \frac{8}{3}$$

$$v_2 = \frac{4}{3} \begin{pmatrix} 0 \\ 0 \\ 2 \\ 1 \end{pmatrix}; \quad w_2 = \frac{1}{\sqrt{5}} \begin{pmatrix} 0 \\ 0 \\ 2 \\ 1 \end{pmatrix}; \quad H_2 = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & -\frac{3}{5} & -\frac{4}{5} \\ 0 & 0 & -\frac{4}{5} & \frac{3}{5} \end{pmatrix} = \frac{1}{5} \begin{pmatrix} 5 & 0 & 0 & 0 \\ 0 & 5 & 0 & 0 \\ 0 & 0 & -3 & -4 \\ 0 & 0 & -4 & 3 \end{pmatrix}$$

$$H_2^T (H_1^T A H_1) H_2 = \begin{pmatrix} 4.0000 & -3.0000 & 0 & 0 \\ -3.0000 & 3.3333 & -1.6667 & 0 \\ 0 & -1.6667 & -1.3200 & 0.9067 \\ 0 & 0 & 0.9067 & 1.9867 \end{pmatrix}$$

70 / 77

Givens matrices (Advanced)

Givens rotation A Givens rotation is an matrix of the form

$$[G_{k,l}(\alpha, \beta)]_{i,j} = \begin{cases} \alpha & \text{if } i = j = k \text{ or } i = k = l \\ \beta & \text{if } i = l \text{ and } j = k \\ -\beta & \text{if } i = k \text{ and } j = l \\ 0 & \text{otherwise.} \end{cases}$$

where $\alpha^2 + \beta^2 = 1$. We can write $\alpha = \cos \theta$ and $\beta = \sin \theta$ for some θ .

Inverse Givens rotation matrices are orthogonal, with inverse

Examples

$$G_{k,l}(\theta)^{-1} = G_{k,l}(\theta)^T = G_{k,l}(-\theta).$$

$$G_{1,2}(\theta) = \begin{pmatrix} \cos \theta & -\sin \theta & 0 & 0 \\ \sin \theta & \cos \theta & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \quad G_{2,4}(\theta) = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & \cos \theta & 0 & -\sin \theta \\ 0 & 0 & 1 & 0 \\ 0 & \sin \theta & 0 & \cos \theta \end{pmatrix}$$

71 / 77

Givens matrices (Advanced)

We can implement the QR algorithm for upper-Hessenberg A by a sequence of Givens rotations

$$A = QR = G_{1,2}(\theta_1) G_{2,3}(\theta_2) G_{3,4}(\theta_3) R$$

so

$$R = G_{3,4}^{-1}(\theta_3) G_{2,3}^{-1}(\theta_2) G_{1,2}^{-1}(\theta_1) A$$

and

$$\begin{aligned} RQ &= R G_{1,2}(\theta_1) G_{2,3}(\theta_2) G_{3,4}(\theta_3) \\ &= G_{3,4}^{-1}(\theta_3) G_{2,3}^{-1}(\theta_2) G_{1,2}^{-1}(\theta_1) A G_{1,2}(\theta_1) G_{2,3}(\theta_2) G_{3,4}(\theta_3) \end{aligned}$$

Note that pre-multiplying A by G_{kl} (or G_{kl}^{-1}) changes rows k and l to linear combinations of each other, and leaves all other rows unchanged.

Likewise, post-multiplying by G_{kl} changes columns k and l , and leaves all other columns unchanged.

72 / 77

Givens matrices (Advanced)

Example

$$A = \begin{pmatrix} 4 & 1 & 7 \\ 3 & 7 & 9 \\ 0 & 12 & 2 \end{pmatrix}$$

Take Givens rotation (with $r_1 = (a_{11}^2 + a_{21}^2)^{1/2}$)

$$G_1 = \begin{pmatrix} a_{11}/r_1 & -a_{21}/r_1 & 0 \\ a_{21}/r_1 & a_{11}/r_1 & 0 \\ 0 & 0 & 1 \end{pmatrix} = \begin{pmatrix} \frac{4}{5} & -\frac{3}{5} & 0 \\ \frac{3}{5} & \frac{4}{5} & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

$$G_1^{-1}A = \begin{pmatrix} \frac{4}{5} & \frac{3}{5} & 0 \\ -\frac{3}{5} & \frac{4}{5} & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 4 & 1 & 7 \\ 3 & 7 & 9 \\ 0 & 12 & 2 \end{pmatrix} = \begin{pmatrix} 5 & 5 & 11 \\ 0 & 5 & 3 \\ 0 & 12 & 2 \end{pmatrix}.$$

$$G_2 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & \frac{5}{13} & -\frac{12}{13} \\ 0 & \frac{12}{13} & \frac{5}{13} \end{pmatrix}; \quad R = G_2^{-1}(G_1^{-1}A) = \begin{pmatrix} 5 & 5 & 11 \\ 0 & 13 & 3 \\ 0 & 0 & -2 \end{pmatrix}.$$

Givens matrices (Advanced)**Example (continued)**

$$\begin{aligned}
RQ &= RG_1G_2 = (RG_1)G_2 \\
&= \begin{pmatrix} 5 & 5 & 11 \\ 0 & 13 & 3 \\ 0 & 0 & -2 \end{pmatrix} \begin{pmatrix} \frac{4}{5} & -\frac{3}{5} & 0 \\ \frac{3}{5} & \frac{4}{5} & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & \frac{5}{13} & -\frac{12}{13} \\ 0 & \frac{12}{13} & \frac{5}{13} \end{pmatrix} \\
&= \begin{pmatrix} 7 & 1 & 11 \\ 7\frac{4}{5} & 10\frac{2}{5} & 3 \\ 0 & 0 & -2 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & \frac{5}{13} & -\frac{12}{13} \\ 0 & \frac{12}{13} & \frac{5}{13} \end{pmatrix} \\
&= \begin{pmatrix} 7 & 10\frac{7}{13} & 3\frac{4}{13} \\ 7\frac{4}{5} & 6\frac{10}{13} & -9\frac{29}{65} \\ 0 & -1\frac{11}{13} & -\frac{10}{13} \end{pmatrix} = \begin{pmatrix} 7.000 & 10.538 & 3.308 \\ 7.800 & 6.769 & -8.446 \\ 0 & -1.846 & -0.769 \end{pmatrix}
\end{aligned}$$

74 / 77

Givens matrices (Advanced)

If A is tridiagonal, then $A = QR$ with $r_{i,j} = 0$ for $i > j$ or $j > i + 2$, so R is banded with three nontrivial bands on the main diagonal and above ($3n$ nonzero entries).

However, Q is upper-Hessenberg, and has $n^2/2$ nonzero entries, even though it is the product of $n - 1$ Givens rotations, which are sparse.

We therefore perform the QR-method using Givens rotations, and do not directly construct Q .

75 / 77

Eigenvalue condition number (Non-examinable)

Eigenvalue condition numbers Suppose λ is a simple eigenvalue of A , $Ax = \lambda x$ and $A^T y = \lambda y$ with $\|x\|_2 = \|y\|_2 = 1$. Suppose $\|F\|_2 = 1$. Then if $\lambda(\epsilon)$ continuous to $A + \epsilon F$, we have $|\lambda'(\epsilon)| = |Y^T F x| / |Y^T x| \leq 1 / |Y \cdot x|$, with the bound attained for $F = yx^T$. Define $s(\lambda) = |y^T x|$ the condition of the eigenvalue.

[From Golub & Van Loon]

76 / 77

Eigenvalue conditioning (Non-examinable)

Eigenvalue conditioning If μ is eigenvalue of perturbation $A + E$ of a nondefective matrix A , then

$$|\mu - \lambda_k| \leq \text{cond}_2(V) \|E\|_2$$

where λ_k is closest eigenvalue of A to μ , and V is the matrix of eigenvectors of A .

[From Michael T. Heath: Scientific Computing: An Introductory Survey]

77 / 77