

Numerical Methods

Differentiation and Integration

Martijn Boussé, Pieter Collins & Başak Sakçak

Department of Advanced Computing Sciences

Maastricht University

`m.bousse,pieter.collins,basak.sakcak@maastrichtuniversity.nl`

KEN1540 & KEN2540, Block 5, April-May 2025

Differentiation	2
Numerical differentiation	3
Differentiation techniques	4
Finite-difference	5
Centred difference	8
Forward difference	15
Five-point schemes	18
Second derivative	21
Appendix: Derivations	27
Three-point schemes	28
Five-point schemes	30
Error estimates	33
Numerical Integration	36
Integration in Matlab	37
Riemann sums	38
Midpoint and Trapezoid Rules	39
Midpoint rule	40
Trapezoid rule	47
Simpson's Rule	53
Simpson's rule	54
Newton-Cotes rules	60
Gaussian quadrature	61
Romberg Integration	62
Extrapolation	63
Romberg integration	65
Adaptive Integration	71
Adaptive integration	72
Adaptive trapezoid	73
Adaptive Simpson	76

Exact and numerical differentiation

Exact differentiation For most functions, the derivatives can be computed exactly using the formulae from Calculus.

e.g. $f(x) = x \sin(x) \implies f'(x) = x \cos(x) + \sin(x).$

Numerical differentiation The aim of numerical differentiation is to estimate the derivatives $f'(x)$ of a function f at a point x for cases where exact computation is impossible or impractical, such as:

- The function is determined by experimentally-measured data i.e. $f(x_i) = y_i$.
- The function is determined by numerically-computed data e.g. the numerical solution of a differential equation.
- The function is specified by black-box code, and only supports the operation of evaluation at points.
- Explicitly computing the derivative would result in a formula which is too complicated to be efficiently stored and/or evaluated.

3 / 78

Numerical differentiation techniques

Finite-difference Compute approximations to derivatives using weighted sums/differences of function values. Formulae are based on polynomial interpolation at nearby data points.

Automatic differentiation (Off-syllabus) A modern technique for computing derivatives exactly without constructing a symbolic formula explicitly. Typically requires access to source code.

4 / 78

Finite-difference approximation

Derivative definition Recall from Calculus that:

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h},$$

Difference approximation Estimate the derivative from the defining formula with $|h|$ small:

$$f'(x) \approx \frac{f(x+h) - f(x)}{h}$$

Note that h may be positive or negative!

5 / 78

Forward/backward difference

Example For $f(x) = 1/(1+x^2)$, estimate $f'(x)$ at $x = 2$ using the forward difference approximation with $h = 0.5, 0.1, 0.01, -0.1$.

$$f'(2) \approx \frac{f(2+0.5) - f(2)}{0.5} = \frac{\frac{1}{1+2.5^2} - \frac{1}{1+2^2}}{0.5} = \frac{\frac{1}{7.25} - \frac{1}{5}}{0.5} = \frac{0.138 - 0.200}{0.5} = -0.124$$

$$f'(2) \approx \frac{f(2+0.1) - f(2)}{0.1} = \frac{\frac{1}{1+2.1^2} - \frac{1}{1+2^2}}{0.1} = \frac{0.1848 - 0.2000}{0.1} = -0.1516$$

$$f'(2) \approx \frac{f(2+0.01) - f(2)}{0.01} = \frac{\frac{1}{1+2.01^2} - \frac{1}{1+2^2}}{0.01} = \frac{0.19841 - 0.20000}{0.01} = -0.15912$$

$$f'(2) \approx \frac{f(2-0.1) - f(2)}{-0.1} = \frac{\frac{1}{1+1.9^2} - \frac{1}{1+2^2}}{-0.1} = \frac{0.2169 - 0.2000}{-0.1} = -0.1692$$

Exact value

$$f'(x) = \frac{-2x}{(1+x^2)^2}; \quad f'(2) = \frac{-2 \times 2}{(1+2^2)^2} = \frac{-4}{25} = -0.1600.$$

Error is fairly large... What are the asymptotics for small h ? Can we do better??

6 / 78

2-point forward difference

Forward difference error By Taylor's theorem, for some ξ between x and $x+h$,

$$f(x+h) = f(x) + hf'(x) + h^2 f''(\xi)/2.$$

Rearranging gives

$$f'(x) = \frac{f(x+h) - f(x)}{h} - \frac{h}{2} f''(\xi).$$

Error term

$$\left| f'(x) - \frac{f(x+h) - f(x)}{h} \right| = \frac{h}{2} |f''(\xi)|.$$

Note that Taylor's theorem gives no way of finding the point ξ . (If it did, we could compute the derivative exactly!)

However, the error term does show the error is $O(h)$.

Backward difference error For some ξ between $x-h$ and x ,

$$f'(x) = \frac{f(x) - f(x-h)}{h} + \frac{h}{2} f''(\xi).$$

7 / 78

Centred difference

Centered difference by quadratic interpolation

Recall interpolation of f at $x - h, x, x + h$ by a quadratic polynomial:

$$p(y) = f(x) + \frac{f(x+h) - f(x-h)}{2h}(y-x) + \frac{f(x-h) - 2f(x) + f(x+h)}{2h^2}(y-x)^2.$$

Compute the derivative:

$$p'(y) = \frac{f(x+h) - f(x-h)}{2h} + \frac{f(x-h) - 2f(x) + f(x+h)}{2h^2} \times 2(y-x);$$
$$p'(x) = \frac{f(x+h) - f(x-h)}{2h}.$$

Centred difference formula Estimate $f'(x)$ by $p'(x)$:

$$f'(x) \approx \frac{f(x+h) - f(x-h)}{2h}.$$

This is called the *three-point* centred difference formula, even though only two function values are needed! The formula is 'missing' the term $0 \times f(x)$.

8 / 78

Centred difference

Centred difference by Taylor series Combine forwards and backwards difference approximations:

$$f(x+h) = f(x) + hf'(x) + h^2 f''(x)/2 + O(h^3)$$

$$f(x-h) = f(x) - hf'(x) + h^2 f''(x)/2 + O(h^3)$$

$$f(x+h) - f(x-h) = 2hf'(x) + O(h^3)$$

Hence

$$f'(x) = \frac{f(x+h) - f(x-h)}{2h} + O(h^2).$$

Centred difference formula

$$f'(x) \approx \frac{f(x+h) - f(x-h)}{2h}.$$

9 / 78

Centred difference error

Centered difference error estimate For some ξ between $x - h$ and $x + h$,

$$f'(x) = \frac{f(x+h) - f(x-h)}{2h} - \frac{h^2}{6} f'''(\xi).$$

Note method is exact for a quadratic polynomial!

Derivation by Taylor's theorem (Advanced)

$$f(x+h) = f(x) + hf'(x) + h^2 f''(x)/2 + h^3 f'''(\xi_+)/6$$

$$f(x-h) = f(x) - hf'(x) + h^2 f''(x)/2 - h^3 f'''(\xi_-)/6$$

$$f(x+h) - f(x-h) = 2hf'(x) + h^3(f'''(\xi_+) + f'''(\xi_-))/6$$

Since $(f'''(\xi_+) + f'''(\xi_-))/2$ lies between $f'''(\xi_+)$ and $f'''(\xi_-)$, there exists ξ such that $f'''(\xi) = (f'''(\xi_+) + f'''(\xi_-))/2$, so

$$f(x+h) - f(x-h) = 2hf'(x) + h^3 f'''(\xi)/3.$$

10 / 78

Centred difference

Example Use the centred difference method with $h = 0.1, 0.01$ to estimate $f'(x)$ at $x = 2$.

$$\begin{aligned} f'(2) &\approx \frac{f(2+0.1) - f(2-0.1)}{2 \times 0.1} = \frac{\frac{1}{1+(2+0.1)^2} - \frac{1}{1+(2-0.1)^2}}{2 \times 0.1} = \frac{\frac{1}{5.41} - \frac{1}{3.61}}{0.2} \\ &= (0.18484 - 0.21792)/0.2 = -0.16039. \\ f'(2) &\approx \frac{f(2+0.01) - f(2-0.01)}{2 \times 0.01} = \frac{\frac{1}{1+(2+0.01)^2} - \frac{1}{1+(2-0.01)^2}}{2 \times 0.01} = \frac{\frac{1}{5.0401} - \frac{1}{3.9601}}{0.02} \\ &= (0.19840876 - 0.2016088)/0.02 = -0.1600038. \end{aligned}$$

11 / 78

Centred difference

Error dependence Estimate $f'(2)$ by using the three-point centred-difference method with $h = 10^{-n}$ for $n = 0, \dots, 14$.

h	f'	$ e $
10^{-0}	-0.200000000000000	$0.040000000000000 \approx 4 \times 10^{-2}$
10^{-1}	-0.160384280736645	$0.000384280736645 \approx 4 \times 10^{-4}$
10^{-2}	-0.160003840028156	$0.000003840028156 \approx 4 \times 10^{-6}$
10^{-3}	-0.160000038399985	$0.000000038399985 \approx 4 \times 10^{-8}$
10^{-4}	-0.160000000384158	$0.000000000384158 \approx 4 \times 10^{-10}$
10^{-5}	-0.16000000004601	$0.00000000004601 \approx 5 \times 10^{-12}$
10^{-6}	-0.160000000012928	$0.000000000012928 \approx 1 \times 10^{-11}$
10^{-7}	-0.15999999915783	$0.000000000084217 \approx 8 \times 10^{-11}$
10^{-8}	-0.159999999360672	$0.000000000639328 \approx 6 \times 10^{-10}$
10^{-9}	-0.160000013238459	$0.000000013238459 \approx 1 \times 10^{-8}$
10^{-10}	-0.15999929971733	$0.00000070028267 \approx 7 \times 10^{-8}$
10^{-11}	-0.15999791193854	$0.00000208806146 \approx 2 \times 10^{-7}$
10^{-12}	-0.160024771211909	$0.000024771211908 \approx 2 \times 10^{-5}$
10^{-13}	-0.159872115546023	$0.000127884453977 \approx 1 \times 10^{-4}$
10^{-14}	-0.160982338570648	$0.000982338570648 \approx 1 \times 10^{-3}$

Small changes in computed $f(x \pm h)$ cause large change in derivative for small h !!

12 / 78

Truncation and roundoff error

Example For $f(x) = 1/(3+x^2)$, have $f'(x) = -2x/(3+x^2)^2$ and $f'''(x) = 24x(3-x^2)/(3+x^2)^4$, with $f'(2.0) = -0.081632$ (6 dp). Rounding to 3 dp gives

x	1.9	2.0	2.1
$y \approx f(x)$	0.151	0.143	0.135

Using the three-point centred-difference approximation to find $f'(2.0)$ yields $f'(2.0) \approx -0.08$, which has an absolute error of 1.63×10^{-3} (2 sf).

Since for $\xi \in [1.9, 2.1]$, $-0.024 \leq f'''(\xi) \leq -0.014$ for all ξ , we can estimate the truncation error as

$$\sup_{\xi \in [1.9, 2.1]} |f'''(\xi)| \frac{h^2}{6} \approx 0.024 \times \frac{0.1^2}{6} = 5 \times 10^{-5}.$$

The roundoff error is

$$\left| \frac{f(x+h) - f(x-h)}{2h} - \frac{0.135 - 0.151}{2 \times 0.1} \right| \leq 1.67 \times 10^{-3}.$$

Hence the main source of error is rounding in the y -values, and *not* the truncation error of the approximation.

13 / 78

Centred difference

Example Estimate $f'(x)$ for $x = 2.0, 2.2, 2.4, 2.6, 2.8, 3.0$ from the following data:

x	2.0	2.2	2.4	2.6	2.8	3.0
$f(x)$	1.386	1.735	2.101	2.484	2.883	3.296

$$f'(2.2) \approx (f(2.4) - f(2.0))/(2 \times 0.2) = (2.101 - 1.386)/0.4 = 1.79$$

$$f'(2.4) \approx (f(2.6) - f(2.2))/(2 \times 0.2) = (2.484 - 1.735)/0.4 = 1.87$$

$$f'(2.6) \approx (f(2.8) - f(2.4))/(2 \times 0.2) = (2.883 - 2.101)/0.4 = 1.96$$

$$f'(2.8) \approx (f(3.0) - f(2.6))/(2 \times 0.2) = (3.296 - 2.484)/0.4 = 2.03$$

Can't estimate $f'(2.0)$ and $f'(3.0)$ using centred difference!

Need a different formula for use at endpoints.

14 / 78

Forward/backward difference

Endpoints of domain For f defined on interval $[a, b]$, cannot use centred difference at/near endpoints a, b , as $f(a - h)$ and $f(b + h)$ are not defined!

Three-point forward-difference formula

$$f'(x) \approx \frac{-f(x + 2h) + 4f(x + h) - 3f(x)}{2h}.$$

Three-point forward-difference error estimate

$$f'(x) = \frac{-f(x + 2h) + 4f(x + h) - 3f(x)}{2h} + \frac{h^2}{3} f'''(\xi).$$

Three-point backward-difference Take h negative in forward-difference formula:

$$f'(x) \approx \frac{f(x - 2h) - 4f(x - h) + 3f(x)}{2h}.$$

15 / 78

Forward/backward difference

Example For $f(x) = 1/(1 + x^2)$, estimate $f'(2)$ using the forward-difference formula with $h = 0.1$.

$$\begin{aligned} f'(2) &\approx \frac{-f(2 + 2 \times 0.1) + 4f(2 + 0.1) - 3f(2)}{2h} = \frac{-f(2.2) + 4f(2.1) - 3f(2.0)}{2 \times 0.1} \\ &= \frac{-0.171 + 4 \times 1.85 - 3 \times 2.00}{0.2} = -0.159307 \end{aligned}$$

Exact value -1.6 ; error 7.9×10^{-4} , roughly twice that for centred difference.

16 / 78

Forward/backward difference

Example Estimate $f'(x)$ for $x = 2.0, 2.2, 2.4, 2.6, 2.8, 3.0$ from the following data:

x	2.0	2.2	2.4	2.6	2.8	3.0
$f(x)$	1.386	1.735	2.101	2.484	2.883	3.296

Use centred-difference where possible:

$$\begin{aligned}f'(2.0) &\approx (-f(2.4) + 4f(2.2) - 3f(2.0))/(2 \times 0.2) \\&= (-2.101 + 4 \times 1.735 - 3 \times 1.386)/0.4 = 1.70\end{aligned}$$

$$f'(2.2) \approx (f(2.4) - f(2.0))/(2 \times 0.2) = (2.101 - 1.386)/0.4 = 1.79$$

$$f'(2.4) \approx (f(2.6) - f(2.2))/(2 \times 0.2) = (2.484 - 1.735)/0.4 = 1.87$$

$$f'(2.6) \approx (f(2.8) - f(2.4))/(2 \times 0.2) = (2.883 - 2.101)/0.4 = 1.96$$

$$f'(2.8) \approx (f(3.0) - f(2.6))/(2 \times 0.2) = (3.296 - 2.484)/0.4 = 2.03$$

$$\begin{aligned}f'(3.0) &\approx (f(2.6) - 4f(2.8) + 3f(3.0))/(2 \times 0.2) \\&= (2.484 - 4 \times 2.883 + 3 \times 3.296)/0.4 = 2.10\end{aligned}$$

17 / 78

Five-point centred difference

Five-point centred difference formula

$$f'(x) \approx \frac{f(x-2h) - 8f(x-h) + 8f(x+h) - f(x+2h)}{12h}$$

Five-point centred difference error estimate

$$f'(x) = \frac{f(x-2h) - 8f(x-h) + 8f(x+h) - f(x+2h)}{12h} + \frac{h^4}{30} f^{(5)}(\xi)$$

Example Estimate $f'(x)$ for $f(x) = 1/(1+x^2)$ with $x = 2$ and $h = 0.1$.

$$\begin{aligned}f'(2) &\approx \frac{f(1.8) - 8f(1.9) + 8f(2.1) - f(2.2)}{12 \times 0.1} \\&= \frac{0.2358491 - 8 \times 0.2169197 + 8 \times 0.1848429 - 0.1712329}{1.2} \\&= -0.1599989 \text{ (7dp)}\end{aligned}$$

Error 1.1×10^{-6} , much better than three-point error 3.9×10^{-4} .

18 / 78

Five-point difference formulae

Five-point forward difference

$$f'(x) = \frac{-25f(x) + 48f(x+h) - 36f(x+2h) + 16f(x+3h) - 3f(x+4h)}{12h} + \frac{h^4}{5}f^{(5)}(\xi)$$

Five-point asymmetric difference

$$f'(x) = \frac{-3f(x-h) - 10f(x) + 18f(x+h) - 6f(x+2h) + f(x+3h)}{12h} - \frac{h^4}{20}f^{(5)}(\xi)$$

Five-point centred difference

$$f'(x) = \frac{f(x-2h) - 8f(x-h) + 8f(x+h) - f(x+2h)}{12h} + \frac{h^4}{30}f^{(5)}(\xi)$$

Backward-difference formulae can be obtained by taking h negative.

19 / 78

Five-point difference

Example Estimate $f'(x)$ for $x = 2.0, 2.2, 2.4$ from the following data:

x	2.0	2.2	2.4	2.6	2.8	3.0
$f(x)$	1.386	1.735	2.101	2.484	2.883	3.296

Use five-point formulae:

$$\begin{aligned}f'(2.0) &\approx (-25f(2.0) + 48f(2.2) - 36f(2.4) + 16f(2.6) - 3f(2.8))/(12 \times 0.2) \\&= (-25 \times 1.386 + 48 \times 1.735 - 36 \times 2.101 + 16 \times 2.484 - 3 \times 2.883)/2.4 \\&= (-34.650 + 83.280 - 75.636 + 39.744 - 8.649)/2.4 \\&= 4.089/2.4 = 1.704 \text{ (3dp)}\end{aligned}$$

$$\begin{aligned}f'(2.2) &\approx (-3f(2.0) - 10f(2.2) + 18f(2.4) - 6f(2.6) + f(2.8))/(12 \times 0.2) \\&= (-3 \times 1.386 - 10 \times 1.735 + 18 \times 2.101 - 6 \times 2.484 + 2.883)/2.4 \\&= (-4.158 - 17.350 + 37.818 - 14.904 + 2.883)/2.4 \\&= 4.289/2.4 = 1.787 \text{ (3dp)}\end{aligned}$$

$$\begin{aligned}f'(2.4) &\approx (f(2.0) - 8f(2.2) + 8f(2.6) - f(2.8))/(12 \times 0.2) \\&= (1.386 - 8 \times 1.735 + 8 \times 2.484 - 2.883)/2.4 \\&= (1.386 - 13.880 + 19.872 - 2.883)/2.4 = 4.495/2.4 = 1.873 \text{ (3dp)}\end{aligned}$$

20 / 78

Second derivative

Second derivative by quadratic interpolation

$$p(y) = f(x) + \frac{f(x+h) - f(x-h)}{2h}(y-x) + \frac{f(x-h) - 2f(x) + f(x+h)}{2h^2}(y-x)^2.$$

Compute the derivatives:

$$p'(y) = \frac{f(x+h) - f(x-h)}{2h} + \frac{f(x-h) - 2f(x) + f(x+h)}{2h^2} \times 2(y-x);$$
$$p''(y) = \frac{f(x-h) - 2f(x) + f(x+h)}{2h^2} \times 2.$$

Three-point second-derivative formula Estimate $f''(x)$ by $p''(x)$:

$$f''(x) \approx \frac{f(x-h) - 2f(x) + f(x+h)}{h^2}.$$

21 / 78

Second derivative

Example For $f(x) = 1/(1+x^2)$, estimate $f''(2)$ using $h = 0.1, 0.01$.

$$f''(2) \approx \frac{f(2.1) - 2f(2) + f(1.9)}{0.1^2} = \frac{\frac{1}{5.41} - 2 \times \frac{1}{5} + \frac{1}{4.61}}{0.01}$$
$$\approx \frac{0.184843 - 2 \times 0.200000 + 0.216920}{0.01} = \frac{0.001763}{0.01} = 0.1763.$$
$$f''(2) \approx \frac{f(2.01) - 2f(2) + f(1.99)}{0.01^2} = \frac{\frac{1}{5.0401} - 2 \times \frac{1}{5} + \frac{1}{4.9601}}{0.0001}$$
$$\approx \frac{0.1984087617 - 2 \times 0.200000000 + 0.2016088385}{0.0001}$$
$$= \frac{0.0000176003}{0.0001} = 0.176003.$$

The exact value is $f''(2) = \frac{22}{125} = 0.176$.

22 / 78

Second derivative

Second derivative with error term

$$f''(x) = \frac{f(x+h) - 2f(x) + f(x-h)}{h^2} - f^{(4)}(\xi)h^2/12.$$

Second derivative by Taylor series (Non-examinable)

$$f(x+h) = f(x) + hf'(x) + h^2f''(x)/2 + h^3f'''(x)/6 + \dots$$

$$f(x-h) = f(x) - hf'(x) + h^2f''(x)/2 - h^3f'''(x)/6 + \dots$$

$$f(x+h) + f(x-h) = 2f(x) + 2h^2f''(x)/2 + 2h^4f^{(4)}(x)/24 + O(h^5)$$

Hence

$$f(x+h) - 2f(x) + f(x-h) = 2h^2f''(x)/2 + 2h^4f^{(4)}(x)/24 + O(h^5).$$

23 / 78

Second derivative

Three-point schemes for second derivative

$$f''(x) = \frac{f(x-h) - 2f(x) + f(x+h)}{h^2} - \frac{1}{12}h^2 f^{(4)}(\xi)$$

$$f''(x) = \frac{f(x) - 2f(x+h) + f(x+2h)}{h^2} - hf^{(3)}(\xi)$$

Five-point schemes for second derivative

$$f''(x) = \frac{-f(x-2h) + 16f(x-h) - 30f(x) + 16f(x+h) - f(x+2h)}{12h^2} + \frac{1}{90}h^4 f^{(6)}(\xi)$$

$$f''(x) = \frac{11f(x-h) - 20f(x) + 6f(x+h) + 4f(x+2h) - f(x+3h)}{12h^2} + \frac{1}{12}h^3 f^{(5)}(\xi)$$

$$f''(x) = \frac{35f(x) - 104f(x+h) + 114f(x+2h) - 56f(x+3h) + 11f(x+4h)}{12h^2} - \frac{5}{6}h^3 f^{(5)}(\xi)$$

Note that the centred-difference approximations have a higher order!

24 / 78

Higher derivatives (Non-examinable)

Five-point schemes for third derivative

$$f'''(x) = \frac{-f(x-2h) + 2f(x-h) - 2f(x+h) + f(x+2h)}{2h^3} - \frac{1}{4}h^2 f^{(5)}(\xi)$$

$$f'''(x) = \frac{-3f(x-h) + 10f(x) - 12f(x+h) + 6f(x+2h) - f(x+3h)}{2h^3} + \frac{1}{4}h^2 f^{(5)}(\xi)$$

$$f'''(x) = \frac{-5f(x) + 18f(x+h) - 24f(x+2h) + 14f(x+3h) - 3f(x+4h)}{2h^3} + \frac{7}{4}h^2 f^{(5)}(\xi)$$

Five-point scheme for fourth derivative

$$f''''(x) = \frac{f(x-2h) - 4f(x-h) + 6f(x) - 4f(x+h) + f(x+2h)}{h^4} - \frac{1}{6}h^2 f^{(6)}(\xi)$$

25 / 78

Exercise

Exercise Estimate $f'(2)$, $f'(3)$ and $f''(3)$ for $f(x) = 1/x$ using

x	2.0	2.5	3.0	3.5	4.0
$f(x)$	0.500	0.400	0.333	0.286	0.250

Estimate the errors by $|f^{(n)}(\xi)| \leq n!/2^{n+1}$.

26 / 78

**Appendix: Derivations and
Error Estimates
(Non-examinable)**

27 / 78

Derivation of three-point schemes

Let $q(y)$ interpolate f at $x - h$, x and $x + h$. Then

$$q(y) = f(x) + \frac{f(x+h) - f(x-h)}{2h}(y-x) + \frac{f(x+h) - 2f(x) + f(x-h)}{2h^2}(y-x)^2$$

Approximate $f'(x) \approx q'(x)$ and $f''(x) \approx q''(x)$

$$f'(x) \approx \frac{f(x+h) - f(x-h)}{2h}; \quad f''(x) \approx \frac{f(x+h) - 2f(x) + f(x-h)}{h^2}.$$

and

$$\begin{aligned} f'(x+h) &\approx q'(x+h) = \frac{f(x+h) - f(x-h)}{2h} + \frac{f(x+h) - 2f(x) + f(x-h)}{h} \\ &= \frac{3f(x+h) - 4f(x) + f(x-h)}{2h} \\ f'(x) &\approx \frac{3f(x) - 4f(x-h) + f(x-2h)}{2h} = \frac{-3f(x) + 4f(x+h) - f(x+2h)}{2h} \end{aligned}$$

28 / 78

Derivation of three-point schemes

Let $q(y)$ be a quadratic polynomial

$$q(y) = a_0 + a_1(y-x) + a_2(y-x)^2; \quad q'(y) = a_1 + 2a_2(y-x).$$

If q interpolates f at $x - h$, x and $x + h$, then

$$\begin{aligned} f(x) &= q(x) = a_0; \\ f(x+h) &= q(x+h) = a_0 + a_1h + a_2h^2; \\ f(x+2h) &= q(x+2h) = a_0 + 2a_1h + 4a_2h^2. \end{aligned}$$

Eliminate a_2 and a_0 by

$$f(x+2h) - 4f(x+h) = -2a_1h - 3a_0 = -2a_1h - 3f(x)$$

Approximate

$$f'(x) \approx q'(x) = a_1 = \frac{-f(x+2h) + 4f(x+h) - 3f(x)}{2h}$$

29 / 78

Derivation of five-point scheme by standard basis

Let $q(y)$ be a quartic polynomial

$$q(y) = a_0 + a_1(y - x) + a_2(y - x)^2 + a_3(y - x)^3 + a_4(y - x)^4.$$

If q interpolates f at $x - 2h, x - h, x, x + h$ and $x + 2h$, then

$$f(x - 2h) = q(x - 2h) = a_0 - 2a_1h + 4a_2h^2 - 8a_3h^3 + 16a_4h^4;$$

$$f(x - h) = q(x - h) = a_0 - a_1h + a_2h^2 - a_3h^3 + a_4h^4;$$

$$f(x) = q(x) = a_0;$$

$$f(x + h) = q(x + h) = a_0 + a_1h + a_2h^2 + a_3h^3 + a_4h^4;$$

$$f(x + 2h) = q(x + 2h) = a_0 + 2a_1h + 4a_2h^2 + 8a_3h^3 + 16a_4h^4.$$

Eliminate a_0, a_2 and a_4 by

$$f(x + 2h) - f(x - 2h) = 4a_1h + 16a_3h^3; \quad f(x + h) - f(x - h) = 2a_1h + 2a_3h^3.$$

Eliminate a_3 by

$$(f(x + 2h) - f(x - 2h)) - 8(f(x + h) - f(x - h)) = -12a_1h.$$

Approximate

$$f'(x) \approx q'(x) = a_1 = \frac{f(x - 2h) - 8f(x - h) + 8f(x + h) - f(x + 2h)}{12h}$$

30 / 78

Derivation of five-point scheme by Lagrange basis

Let $x_i = ih$ for $i = -2, -1, 0, 1, 2$.

Let l_{-2} be the Lagrange basis polynomial

$$l_{-2}(y) = \frac{(y + h)y(y - h)(y - 2h)}{24h^4} = \frac{y^4 - 2hy^3 - h^2y^2 + 2h^3y}{24h^4}$$

satisfying

$$l_{-2}(-2h) = 1, \quad l_{-2}(-h) = l_{-2}(0) = l_{-2}(h) = l_{-2}(2h) = 0$$

Since the term in y has coefficient $2h^3/24h^4$, have $l'_{-2}(0) = 1/12h$.

Let l_{-1} be the Lagrange basis polynomial

$$l_{-1}(y) = \frac{(y + 2h)y(y - h)(y - 2h)}{-6h^4} = \frac{-y^4 + hy^3 + 4h^2y^2 - 4h^3y}{6h^4}$$

satisfying

$$l_{-1}(-h) = 1, \quad l_{-1}(-2h) = l_{-1}(0) = l_{-1}(h) = l_{-1}(2h) = 0$$

Since the term in y has coefficient $-4h^3/6h^4$, have $l'_{-1}(0) = -8/12h$.

Similarly, $l'_0(0) = 0$, $l'_1(0) = 8/12h$ and $l'_2(0) = -1/12h$.

31 / 78

Derivation of five-point scheme by Lagrange basis

If q is a quartic interpolating y at x_{-2}, x_{-1}, x_0, x_1 and x_2 , then

$$q(y) = f(x_{-2})l_{-2}(y) + f(x_{-1})l_{-1}(y) + f(x_0)l_0(y) + f(x_1)l_1(y) + f(x_2)l_2(y)$$

so

$$\begin{aligned} q'(0) &= f(x_{-2})l'_{-2}(0) + f(x_{-1})l'_{-1}(0) + f(x_0)l'_0(0) + f(x_1)l'_1(0) + f(x_2)l'_2(0) \\ &= \frac{f(x_{-2}) - 8f(x_{-1}) + 8f(x_1) - f(x_2)}{12h} \end{aligned}$$

Hence approximate

$$f'(0) \approx q'(0) = \frac{f(-2h) - 8f(-h) + 8f(h) - f(2h)}{12h}$$

By shifting the x -axis, find

$$f'(x) \approx \frac{f(x-2h) - 8f(x-h) + 8f(x+h) - f(x+2h)}{12h}$$

32 / 78

Error estimate for centred difference

Error estimate for centred difference Fix x, h . A polynomial p interpolating f at $x-h, x, x+h$ and also satisfying $p'(x) = f'(x)$ is given by

$$\begin{aligned} p(y) &= f(x) + f'(x)(y-x) + \frac{f(x-h) - 2f(x) + f(x+h)}{2h^2}(y-x)^2 \\ &\quad + \left(\frac{f(x+h) - f(x-h)}{2h^3} - \frac{f'(x)}{h^2} \right)(y-x)^3. \end{aligned}$$

Let $g(y) = p(y) - f(y)$. By the interpolation conditions, $g(x) = g'(x) = g(x-h) = g(x+h) = 0$, so $g'''(\xi) = 0$ for some $\xi \in [x-h, x+h]$.

For this ξ , we have $p'''(\xi) = f'''(\xi)$, or

$$6 \left(\frac{f(x+h) - f(x-h)}{2h^3} - \frac{f'(x)}{h^2} \right) = f'''(\xi).$$

Rearranging gives

$$f'(x) = \frac{f(x+h) - f(x-h)}{2h} - \frac{f'''(\xi)}{6}h^2.$$

33 / 78

Error estimate for centred difference

Error estimate for centred difference Fix x, h . A polynomial p interpolating f at $x-h, x, x+h$ is given by

$$p(y) = f(x) + \frac{f(x+h) - f(x-h)}{2h}(y-x) + \frac{f(x-h) - 2f(x) + f(x+h)}{2h^2}(y-x)^2.$$

Note that

$$p'(x) = \frac{f(x+h) - f(x-h)}{2h}$$

A cubic q satisfying $q(x-h) = q(x) = q(x+h) = 0$ and $q'(x) = 1$ is

$$q(y) = (y-x)(1 - (y-x)^2/h^2).$$

Let $E = f'(x) - p'(x)$, and Let $g(y) = p(y) + Eq(y) - f(y)$.

Then $g(x) = g(x-h) = g(x+h) = g'(x) = 0$, so $g'''(\xi) = 0$ for some $\xi \in [x-h, x+h]$.

For this ξ , we have $Eq'''(\xi) = f'''(\xi)$, or

$$-6/h^2 E = -\frac{6}{h^2} \left(f'(x) - \frac{f(x+h) - f(x-h)}{2h} \right) = f'''(\xi)$$

Rearranging gives

$$f'(x) = \frac{f(x+h) - f(x-h)}{2h} - \frac{f'''(\xi)}{6}h^2.$$

Error estimate guess using polynomial

Let

$$\begin{aligned} p(y) &= \frac{(y+2h)(y+h)y(y-h)(y-2h)}{4h^4} \\ &= \frac{(y^2-4h^2)(y^2-h^2)y}{4h^4} = \frac{y^5-5h^2y^3+4h^4y}{4h^4} \end{aligned}$$

Then $p(y) = 0$ when $y = -2h, -h, 0, h, 2h$, but $p'(0) = 1$ and $p^{(5)}(y) = 30/h^4$.

Hence the error in approximating $p'(0)$ by the five-point centered difference formula is $p^{(5)}(\xi)h^4/30$.

This error also holds for any non-polynomial:

$$f'(x) = \frac{f(x-2h) - 8f(x-h) + 8f(x+h) - f(x+2h)}{12h} + \frac{1}{30}f^{(5)}(\xi)h^4$$

Integration in Matlab

The Matlab command `integral(f,a,b)` computes the definite integral

$$\int_a^b f(x) dx.$$

Example Use Matlab to compute

$$\int_1^4 \frac{1}{1+x^2} dx$$

Solution

```
f=@(x)1./(1+x.^2); a=1; b=4;
I=integral(f,a,b)
```

or even

```
I=integral(@(x)1./(1+x.^2),1,4)
```

Matlab yields the answer 0.540419500270584.

This is equal to the exact answer $\text{atan}(4) - \text{atan}(1)$ to 15 dp.

37 / 78

Riemann sums

Partition Partition interval $[a, b]$ by $P = \{a = x_0 < x_1 < x_2 < \dots < x_{n-1} < x_n = b\}$.

Define spacing $h_i = \Delta x_i = x_{i+1} - x_i$.

Maximum spacing $\|P\| = \max_{i=0,\dots,n-1} (x_{i+1} - x_i)$.

Riemann sum Choose $c_i \in [x_i, x_{i+1}]$ for $i = 0, \dots, n-1$. Approximate

$$\int_a^b f(x) dx \approx \sum_{i=0}^{n-1} (x_{i+1} - x_i) f(c_i)$$

Riemann integral Limit of Riemann sums as $\|P\| \rightarrow 0$.

38 / 78

Midpoint and Trapezoid Rules

Midpoint rule

Single interval Approximate f over the interval $[a, b]$ by the constant $f(c)$ where $c = (a + b)/2$.

Simple midpoint rule Then

$$\int_a^b f(x) dx \approx M(f, [a, b]) := \int_a^b f(c) dx = (b - a)f(c) = (b - a)f\left(\frac{a+b}{2}\right).$$

Example For $f(x) = x^4$ on $[0, 1]$,

$$\int_0^1 f(x) dx \approx M(f, [0, 1]) = (1 - 0)f\left(\frac{0+1}{2}\right) = (1 - 0)f\left(\frac{1}{2}\right) = 1 \times \frac{1}{16} = 1/16.$$

The exact value is $\int_0^1 f(x) dx = 1/5$; relative error 69%!

40 / 78

Midpoint rule

Partition Divide $[a, b]$ into n subintervals with nodes $a = x_0 < x_1 < \dots < x_n = b$.

Let $P = \{x_0, x_1, \dots, x_n\}$ and $h_i = x_{i+1} - x_i$.

Composite midpoint rule

$$\int_a^b f(x) dx \approx M(f, P) := \sum_{i=0}^{n-1} (x_{i+1} - x_i) f\left(\frac{x_i + x_{i+1}}{2}\right).$$

Writing $x_{i+1/2} = (x_i + x_{i+1})/2$,

$$\int_a^b f(x) dx \approx M(f, P) := \sum_{i=0}^{n-1} (x_{i+1} - x_i) f(x_{i+1/2}).$$

Equally-spaced nodes If $h_i \equiv h = (b - a)/n$, then $x_i = a + hi$ and

$$\int_a^b f(x) dx \approx M_n(f, [a, b]) := \frac{b-a}{n} \sum_{i=0}^{n-1} f\left(\frac{x_i + x_{i+1}}{2}\right) = h \sum_{i=0}^{n-1} f\left(a + \left(i + \frac{1}{2}\right)h\right).$$

Alternative indexing: $M_n(f, [a, b]) = h \sum_{i=1}^n f\left(a + \left(i - \frac{1}{2}\right)h\right)$.

41 / 78

Midpoint rule

Example Use the midpoint rule with $P = \{1.0, 1.5, 2.0, 3.0, 3.5, 4.0\}$ to estimate $\int_a^b f(x) dx$ with $f(x) = 1/(1 + x^2)$, $a = 1$, $b = 4$.

Table of function values $f(x_{i+1/2})$ for $i = 0, \dots, 4$:

$x_{i+1/2}$	1.25	1.75	2.5	3.25	3.75
$f(x_{i+1/2})$	0.39024	0.24615	0.13793	0.08649	0.06639

Approximation

$$\begin{aligned}
M(f, P) &= (x_1 - x_0)f(x_{\frac{1}{2}}) + (x_2 - x_1)f(x_{1\frac{1}{2}}) + (x_3 - x_2)f(x_{2\frac{1}{2}}) \\
&\quad + (x_4 - x_3)f(x_{3\frac{1}{2}}) + (x_5 - x_4)f(x_{4\frac{1}{2}}) \\
&= (1.5 - 1.0)f(1.25) + (2.0 - 1.5)f(1.75) + (3.0 - 2.0)f(2.5) \\
&\quad + (3.5 - 3.0)f(3.25) + (4.0 - 3.5)f(3.75) \\
&= 0.5 \times 0.39024 + 0.5 \times 0.24615 + 1.0 \times 0.13793 + 0.5 \times 0.08649 + 0.5 \times 0.06639 \\
&= 0.19512 + 0.12308 + 0.13793 + 0.04324 + 0.03320 = 0.53257 \text{ (5dp)}
\end{aligned}$$

Exact answer $0.54041950\dots$, error 7.9×10^{-3} , relative error 1.5%.

42 / 78

Midpoint rule error analysis (non-examinable)

Error analysis For interval $[-h/2, +h/2]$, let $E = \int_{-h/2}^{h/2} f(x)dx - hf(0)$. Let $p(x) = f(0) + xf'(0) + 12Ex^2/h^3$, $g(x) = p(x) - f(x)$.

Clearly

$$g(0) = 0 \quad \text{and} \quad g'(0) = 0.$$

Note that

$$\int_{-h/2}^{h/2} dx = h, \quad \int_{-h/2}^{h/2} x dx = 0 \quad \text{and} \quad \int_{-h/2}^{h/2} x^2 dx = h^3/12.$$

Hence

$$\int_{-h/2}^{h/2} p(x)dx = hf(0) + 12E/h^3 \times h^3/12 = hf(0) + E = \int_{-h/2}^{h/2} f(x)dx.$$

Therefore $\int_{-h/2}^{h/2} g(x) = 0$, so g has a zero s in $(-h/2, h/2)$.

If the only zero of g is at 0, then $g''(0) = 0$ since g must change sign somewhere, and $g'(0) = 0$. If $s \neq 0$, then by Rolle's theorem, g' has a zero between s and 0, so g'' has a zero at $\xi \in (-h/2, h/2)$.

$$g''(\xi) = 24E/h^3 - f''(\xi) = 0 \implies E = h^3 f''(\xi)/24.$$

43 / 78

Midpoint rule error

Error term For the simple midpoint rule, there exists $\xi \in [a, b]$ such that

$$\int_a^b f(x) dx = (b-a)f\left(\frac{a+b}{2}\right) + \frac{1}{24}f''(\xi)(b-a)^3.$$

For the composite midpoint rule with n equal subdivisions of width h , there exists $\xi \in [a, b]$ such that

$$\int_a^b f(x) dx = \frac{b-a}{n} \sum_{i=0}^{n-1} f(x_{i+1/2}) + \frac{b-a}{24} h^2 f''(\xi).$$

44 / 78

Midpoint rule error bound (Advanced)

Norm of a partition For a partition P recall $\|P\| := \sup_{i=1, \dots, n} (x_i - x_{i-1})$.

Error bounds An upper bound for the error of the midpoint rule is

$$\left| \int_a^b f(x)dx - M(f, P) \right| \leq \frac{b-a}{24} \|P\|^2 \sup_{\xi \in [a, b]} |f''(\xi)|.$$

For a partition into n equal subintervals of $[a, b]$ with width h , then

$$\left| \int_a^b f(x)dx - M_n(f, [a, b]) \right| \leq \frac{b-a}{24} h^2 \sup_{\xi \in [a, b]} |f''(\xi)|.$$

Suppose B_2 is a constant such that

$$\forall \xi \in [a, b], |f''(\xi)| \leq B_2.$$

Then for $\|P\| \leq h$, an upper bound for the error is

$$\left| \int_a^b f(x)dx - M(f, P) \right| \leq \frac{b-a}{24} h^2 B_2.$$

45 / 78

Midpoint rule error bound (Advanced)

Example How many subdivisions suffice to evaluate $\int_0^1 x^2 dx$ to an accuracy of $\epsilon = 10^{-3}$ using the midpoint rule?

Note that $|f''(\xi)| = 2$ for all $\xi \in [0, 1]$, so $B_2 := \sup_{\xi \in [0,1]} |f''(\xi)| = 2$.

For an error bound of at most ϵ , need

$$\frac{(b-a)^3 B_2}{24n^2} \leq \epsilon.$$

Hence

$$n^2 \geq \frac{(b-a)^3 B_2}{24\epsilon} = \frac{1^3 \times 2}{24 \times 10^{-3}} = 10^3/12 = 83.\dot{3}$$

so require $n = 10$ with step-size $h = 0.1$.

46 / 78

Trapezoid rule

Single interval Interpolate f by a linear function l between a and b , so

$$l(x) = f(a) + \frac{x-a}{b-a}(f(b) - f(a)).$$

Note

$$\int_a^b \frac{x-a}{b-a} dx = \left[\frac{(x-a)^2}{2(b-a)} \right]_a^b = \frac{(b-a)^2}{2(b-a)} = \frac{b-a}{2}.$$

Hence

$$\begin{aligned} \int_a^b f(x) dx &\approx \int_a^b f(a) + \frac{x-a}{b-a}(f(b) - f(a)) dx \\ &= (b-a)f(a) + \frac{b-a}{2}(f(b) - f(a)) = (b-a) \frac{f(a) + f(b)}{2}. \end{aligned}$$

This is the area of the trapezoid with vertices $\{(a, 0), (b, 0), (a, f(a)), (b, f(b))\}$!

Simple trapezoid rule

$$\int_a^b f(x) dx \approx T(f, [a, b]) := (b-a) \frac{f(a) + f(b)}{2}$$

Example For $f(x) = x^4$ on $[0, 1]$,

$$1/5 = \int_0^1 f(x) dx \approx T(f, [0, 1]) = (f(0) + f(1))/2 = (0 + 1)/2 = 1/2.$$

47 / 78

Trapezoid rule

Composite trapezoid rule For partition $P = \{x_0, x_1, \dots, x_n\}$, approximate the integral on each subinterval by the simple trapezoid rule:

$$\int_a^b f(x) dx \approx T(f, P) := \sum_{i=0}^{n-1} (x_{i+1} - x_i) \frac{f(x_i) + f(x_{i+1})}{2}.$$

Equally-spaced nodes With $h_i \equiv h = (b-a)/n$ and $x_i = a + hi$,

$$\begin{aligned} \int_a^b f(x) dx &\approx T_n(f, [a, b]) := \frac{b-a}{n} \left(\frac{1}{2}f(x_0) + \sum_{i=1}^{n-1} f(x_i) + \frac{1}{2}f(x_n) \right) \\ &= h \left(\frac{1}{2}f(a) + \sum_{i=1}^{n-1} f(a + hi) + \frac{1}{2}f(b) \right) \end{aligned}$$

48 / 78

Trapezoid rule

Example Use the trapezoid rule with $n = 6$ to estimate

$$\int_1^4 \frac{1}{1+x^2} dx$$

Have $a = 1$, $b = 4$, $h = (b - a)/n = (4 - 1)/6 = 0.5$, $x_i = 1.0 + 0.5i$.

Table of function values $f(x) = 1/(1 + x^2)$:

x_i	1.0	1.5	2.0	2.5	3.0	3.5	4.0
$f(x_i)$	0.50000	0.30769	0.20000	0.13793	0.10000	0.07547	0.05882

Approximation

$$\begin{aligned} T_6(f, [1, 4]) &= h\left(\frac{1}{2}f(x_0) + f(x_1) + f(x_2) + f(x_3) + f(x_4) + f(x_5) + \frac{1}{2}f(x_6)\right) \\ &= 0.5 \times \left(\frac{1}{2} \times 0.50000 + 0.30769 + 0.20000 + 0.13793 \right. \\ &\quad \left. + 0.10000 + 0.07547 + \frac{1}{2} \times 0.05882\right) \\ &= \frac{1}{2} (0.25000 + 1.23077 + 0.40000 + 0.55172 + 0.20000 + 0.30189 + 0.02941) \\ &= \frac{1}{2} \times 1.10051 = 0.55025 \text{ (5dp)} \end{aligned}$$

Exact answer $0.54041950 \dots$, error 9.8×10^{-3} , relative error 1.8%.

49 / 78

Trapezoid rule error analysis (Non-examinable)

Error analysis For a single interval $[0, h]$, set $E = \int_0^h f(x) dx - h \frac{f(0)+f(h)}{2}$. Note

$$\int_0^h x(h-x) dx = [hx^2/2 - x^3/3]_0^h = h^3/2 - h^3/3 = h^3/6.$$

Let

$$g(x) = f(0) + x \frac{f(h)-f(0)}{h} + 6Ex(h-x)/h^3 - f(x).$$

Then

$$\int_0^h g(x) dx = hf(0) + \frac{h^2}{2h}(f(h) - f(0)) + E - \int_0^h f(x) dx = 0.$$

By the mean value theorem, there exists $s \in (0, h)$ such that $g(s) = 0$. Then g has roots $0, s, h$ in $[0, h]$. By Rolle's theorem, g' has two distinct roots in $(0, h)$, and g'' has a root ξ in $(0, h)$. Hence

$$g''(\xi) = -12E/h^3 - f''(\xi) = 0$$

So

$$E = \int_0^h f(x) dx - h \frac{f(0)+f(h)}{2} = -f''(\xi)h^3/12$$

50 / 78

Trapezoid rule error

Error term For the simple trapezoid rule, there exists $\xi \in [a, b]$ such that

$$\int_a^b f(x) dx = (b-a) \frac{f(a)+f(b)}{2} - \frac{1}{12} f''(\xi)(b-a)^3.$$

For the composite trapezoid rule with n equal subdivisions of width h , there exists $\xi \in [a, b]$ such that

$$\int_a^b f(x) dx = T_n(f, [a, b]) - \frac{b-a}{12} h^2 f''(\xi).$$

51 / 78

Trapezoid rule error bound (Advanced)

Error bound An upper bound for the error of the trapezoid rule is

$$\left| \int_a^b f(x) dx - T(f, P) \right| \leq \frac{b-a}{12} \|P\|^2 \sup_{\xi \in [a,b]} |f''(\xi)|.$$

For a partition into n equal subintervals of $[a, b]$ with width h , then

$$\left| \int_a^b f(x) dx - T_n(f, [a, b]) \right| \leq \frac{b-a}{12} h^2 \sup_{\xi \in [a,b]} |f''(\xi)|.$$

Example For $f(x) = 1/(1+x^2)$, $a = 1$, $b = 4$, find $f''(x) = 2(4x^2 - 1)/(1+x^2)^3$ and $\sup_{\xi \in [1,4]} |f''(\xi)| = |f''(1)| = 0.75$.

So for $n = 6$, have $h = 0.5$ and error bound

$$E \leq \frac{(b-a)}{12} h^2 \sup_{\xi \in [a,b]} |f''(\xi)| = \frac{4-1}{12} \times 0.5^2 \times 0.75 = \frac{3}{64} = 4.7 \times 10^{-2} \text{ (2sf)}.$$

The actual error is 9.8×10^{-3} , well below the error bound.

52 / 78

Simpson's Rule

53 / 78

Simpson's rule

Single scaled interval Interpolate f by a quadratic polynomial at $-h, 0, h$.

$$p(x) = f(0) + \frac{f(h) - f(-h)}{2} \frac{x}{h} + \frac{f(h) - 2f(0) + f(-h)}{2} \frac{x^2}{h^2}$$

Then

$$\begin{aligned} \int_{-h}^h p(x) dx &= \int_{-h}^h f(0) + \frac{f(h) - f(-h)}{2} \frac{x}{h} + \frac{f(h) - 2f(0) + f(-h)}{2} \frac{x^2}{h^2} dx \\ &= 2 \int_0^h f(0) + \frac{f(h) - 2f(0) + f(-h)}{2} \frac{x^2}{h^2} dx \\ &= 2 \left[x f(0) + \frac{f(h) - 2f(0) + f(-h)}{2} \frac{x^3}{3h^2} \right]_0^h \\ &= 2 \left(h f(0) + \frac{f(h) - 2f(0) + f(-h)}{2} \frac{h^3}{3h^2} \right) \\ &= \frac{2h}{6} (f(-h) + 4f(0) + f(h)) \end{aligned}$$

54 / 78

Simpson's rule

Simple Simpson's rule

$$\int_a^b f(x) dx \approx S(f, [a, b]) = \frac{b-a}{6} (f(a) + 4f(\frac{a+b}{2}) + f(b)).$$

Error term

$$\int_a^b f(x) dx = \frac{b-a}{6} (f(a) + 4f(\frac{a+b}{2}) + f(b)) - \frac{f^{(4)}(\xi)}{2880} (b-a)^5.$$

55 / 78

Simpson's rule

Composite Simpson's rule; equally-spaced nodes For $n = 2m$ (even!) subintervals of length $h = (b - a)/n$ with $x_i = a + ih$:

$$\begin{aligned} S_n(f, [a, b]) &= \frac{h}{3} (f(x_0) + 4f(x_1) + 2f(x_2) + 4f(x_3) \\ &\quad + 2f(x_4) + \cdots + 2f(x_{n-2}) + 4f(x_{n-1}) + f(x_n)) \\ &= \frac{h}{3} \left(f(x_0) + 4 \sum_{i=1}^{n/2} f(x_{2i-1}) + 2 \sum_{i=1}^{n/2-1} f(x_{2i}) + f(x_n) \right) \end{aligned}$$

Error bound A bound for the error on subdividing $[a, b]$ into $n = 2m$ subintervals of length $h = (b - a)/n$ is

$$\left| \int_a^b f(x) dx - S_n(f, [a, b]) \right| \leq \frac{(b - a)}{180} h^4 \sup_{\xi \in [a, b]} |f^{(4)}(\xi)|$$

56 / 78

Simpson's rule

Simpson's rule, non-equal subdivisions For partition $P = a = x_0 < x_1 < \cdots < x_m = b$, set $x_{i+1/2} = (x_i + x_{i+1})/2$.

$$S(f, P) = \sum_{i=0}^{m-1} (x_{i+1} - x_i) \frac{f(x_i) + 4f(x_{i+1/2}) + f(x_{i+1})}{6}$$

Note that here we are evaluating f at points indexed

$$x_0, x_{\frac{1}{2}}, x_1, x_{1\frac{1}{2}}, x_2, \dots, x_{m-1}, x_{m-\frac{1}{2}}, x_m!$$

57 / 78

Simpson's rule

Polynomials For the third-order polynomial x^3 , Simpson's rule is exact.

$$\int_{-h}^h x^3 dx = 0; \quad S(x^3, [-h, +h]) = h((-h)^3 + h^3)/3 = 0.$$

For the fourth-order polynomial x^4 , the error E of using Simpson's rule is

$$\int_{-h}^h x^4 dx = 2h^5/5; \quad S(x^4, [-h, +h]) = 2h^5/3; \quad E = 4h^5/15$$

58 / 78

Simpson's rule

Example Use Simpson's rule with $n = 6$ to estimate

$$\int_1^4 \frac{1}{1+x^2} dx$$

Have $a = 1$, $b = 4$, $h = (b - a)/n = (4 - 1)/6 = 0.5$, $x_i = 1.0 + 0.5i$.

Table of function values $f(x) = 1/(1 + x^2)$:

x_i	1.0	1.5	2.0	2.5	3.0	3.5	4.0
$f(x_i)$	0.50000	0.30769	0.20000	0.13793	0.10000	0.07547	0.05882

Approximation

$$\begin{aligned}
 S_6(f, [1, 4]) &= \frac{h}{3} (f(x_0) + 4f(x_1) + 2f(x_2) + 4f(x_3) + 2f(x_4) + 4f(x_5) + f(x_6)) \\
 &= \frac{0.5}{3} (0.50000 + 4 \times 0.30769 + 2 \times 0.20000 + 4 \times 0.13793 \\
 &\quad + 2 \times 0.10000 + 4 \times 0.07547 + 0.05882) \\
 &= \frac{1}{6} (0.50000 + 1.23077 + 0.40000 + 0.55172 + 0.20000 + 0.30189 + 0.05882) \\
 &= \frac{1}{6} \times 3.24320 = 0.54053 \text{ (5dp)}
 \end{aligned}$$

Exact answer $0.54041950 \dots$, error 1.1×10^{-4} , relative error 0.02% .

59 / 78

Newton-Cotes formulae

Simpson's three-eighths rule

$$\begin{aligned}
 \int_a^b f(x) dx &= \frac{b-a}{8} (f(a) + 3f(\frac{2a+b}{3}) + 3f(\frac{a+2b}{3}) + f(b)) - \frac{(b-a)^5}{6480} f^{(4)}(\xi) \\
 &= \frac{3h}{8} (f(x_0) + 3f(x_1) + 3f(x_2) + f(x_3)) - \frac{(b-a)}{80} h^4 f^{(4)}(\xi).
 \end{aligned}$$

where $h = (b - a)/3$, $x_i = a + ih$.

Note that this formula uses three partition intervals, and has error $O(h^4)$, so can be used to apply Simpson's rule to the case of an odd number of partition intervals!

Sixth-order Newton-Cotes (Non-examinable)

$$\begin{aligned}
 \int_a^b f(x) dx &\approx \frac{b-a}{90} (7f(a) + 32f(\frac{3a+b}{4}) + 12f(\frac{a+b}{2}) + 32f(\frac{a+3b}{4}) + 7f(b)) \\
 &= \frac{2h}{45} (7f(x_0) + 32f(x_1) + 12f(x_2) + 32f(x_3) + 7f(x_4)) - \frac{2(b-a)}{945} h^6 f^{(6)}(\xi).
 \end{aligned}$$

where $h = (b - a)/4$, $x_i = a + ih$.

60 / 78

Gaussian quadrature (Non-examinable)

Two-point Gaussian quadrature

$$\begin{aligned}\int_{-1}^{+1} f(x) dx &= (f(-\sqrt{1/3}) + f(+\sqrt{1/3})) + \frac{b-a}{4320} h^4 f^{(4)}(\xi) \\ &\approx f(-0.57735027) + f(+0.57735027)\end{aligned}$$

where $b - a = h = 2$.

Three-point Gaussian quadrature

$$\begin{aligned}\int_{-1}^{+1} f(x) dx &= \left(\frac{5}{9}f(-\sqrt{3/5}) + \frac{8}{9}f(0) + \frac{5}{9}f(+\sqrt{3/5})\right) + \frac{b-a}{2016000} h^6 f^{(6)}(\xi) \\ &\approx 0.55555556f(-0.77459667) + 0.88888889f(0.00000000) \\ &\quad + 0.55555556f(+0.77459667)\end{aligned}$$

Four-point Gaussian quadrature

$$\begin{aligned}\int_{-1}^{+1} f(x) dx &\approx 0.34785485f(-0.86113631) + 0.65214515f(-0.33998104) \\ &\quad + 0.65214515f(+0.33998104) + 0.34785485f(+0.86113631)\end{aligned}$$

61 / 78

Romberg Integration

62 / 78

Richardson extrapolation

Richardson extrapolation Since the error of the trapezoid rule is $O(h^2)$, over a fixed interval $[a, b]$ we can estimate for $I(f) = \int_a^b f(x) dx$:

$$E_{T_{2n}} = I(f) - T_{2n}(f) \approx \frac{1}{4}(I(f) - T_n(f)) = \frac{1}{4}E_{T_n}.$$

Rearranging gives

$$T_{2n}(f) + E_{T_2} = I(f) \approx T_n(f) + 4E_{T_2} \implies E_{T_2} = \frac{1}{3}(T_{2n}(f) - T_n(f))$$

so

$$\begin{aligned}I(f) &\approx T_{2n}(f) + \frac{1}{3}(T_{2n}(f) - T_n(f)) \\ &= \frac{1}{3}(4T_{2n}(f) - T_n(f)) = \frac{4}{3}T_{2n}(f) - \frac{1}{3}T_n(f).\end{aligned}$$

This gives an improved estimate of the integral!

For the case $n = 1$,

$$\begin{aligned}\frac{4}{3}T_2(f) - \frac{1}{3}T_1(f) &= \frac{4}{3} \frac{b-a}{2} \left(\frac{1}{2}f(a) + f\left(\frac{a+b}{2}\right) + \frac{1}{2}f(b)\right) - \frac{1}{3} \frac{b-a}{1} \left(\frac{1}{2}f(a) + \frac{1}{2}f(b)\right) \\ &= \frac{b-a}{6} \left(f(a) + 4f\left(\frac{a+b}{2}\right) + f(b)\right).\end{aligned}$$

This is just Simpson's rule!

63 / 78

Richardson extrapolation

Richardson extrapolation of Simpson's rule Since the error of Simpson's rule is $O(h^4)$, we can estimate

$$S_{2n}(f) - I(f) \approx \frac{1}{16}(S_n(f) - I(f))$$

Rearranging gives

$$\begin{aligned} I(f) &\approx S_{2n}(f) + \frac{1}{15}(S_{2n}(f) - S_n(f)) \\ &= \frac{1}{15}(16S_{2n}(f) - S_n(f)) = \frac{16}{15}S_{2n}(f) - \frac{1}{15}S_n(f). \end{aligned}$$

This gives a further improved estimate of the integral!

This idea of using two approximations of the same order, and attempting to cancel the errors is known as *Richardson extrapolation*.

For integration, this can be continued indefinitely, yielding *Romberg integration*.

64 / 78

Romberg integration

Romberg integration For fixed n , let $h_n = \frac{b-a}{n}$ and T_n the trapezoid approximation

$$T_n(f, [a, b]) = h_n \left(\frac{1}{2}f(a) + \sum_{k=1}^{n-1} f(a + kh_n) + \frac{1}{2}f(b) \right).$$

Notice that we can reuse results of $f(\cdot)$ for $T_{2^{i-1}}$ in computing T_{2^i} since

$$T_{2m} = \frac{1}{2}T_m + h_{2m} \sum_{k=1}^m f(a + (k - \frac{1}{2})h_m).$$

Define initial Romberg estimates

$$R_{i,0} = T_{2^i}$$

Apply Richardson extrapolation to obtain higher-order estimates for $j > 0$:

$$\begin{aligned} R_{i,j} &= R_{i,j-1} + \frac{R_{i,j-1} - R_{i-1,j-1}}{4^j - 1} \\ &= \frac{4^j R_{i,j-1} - R_{i-1,j-1}}{4^j - 1} \\ &= \frac{4^j}{4^j - 1} R_{i,j-1} - \frac{1}{4^j - 1} R_{i-1,j-1} \end{aligned}$$

65 / 78

Romberg integration

Write as a table

$R_{0,0} = T_1$			
$R_{1,0} = T_2$	$R_{1,1} = \frac{4}{3}R_{1,0} - \frac{1}{3}R_{0,0}$		
$R_{2,0} = T_4$	$R_{2,1} = \frac{4}{3}R_{2,0} - \frac{1}{3}R_{1,0}$	$R_{2,2} = \frac{16}{15}R_{2,1} - \frac{1}{15}R_{1,1}$	
$R_{3,0} = T_8$	$R_{3,1} = \frac{4}{3}R_{3,0} - \frac{1}{3}R_{2,0}$	$R_{3,2} = \frac{16}{15}R_{3,1} - \frac{1}{15}R_{2,1}$	$R_{3,3} = \frac{64}{63}R_{3,2} - \frac{1}{63}R_{2,2}$

Alternatively, use the formulae

$R_{0,0} = T_1$			
$R_{1,0} = T_2$	$R_{1,1} = \frac{1}{3}(4R_{1,0} - R_{0,0})$		
$R_{2,0} = T_4$	$R_{2,1} = \frac{1}{3}(4R_{2,0} - R_{1,0})$	$R_{2,2} = \frac{1}{15}(16R_{2,1} - R_{1,1})$	
$R_{3,0} = T_8$	$R_{3,1} = \frac{1}{3}(4R_{3,0} - R_{2,0})$	$R_{3,2} = \frac{1}{15}(16R_{3,1} - R_{2,1})$	$R_{3,3} = \frac{1}{63}(64R_{3,2} - R_{2,2})$

66 / 78

Romberg integration

Example Estimate $\int_0^\pi \sin x \, dx$ using Romberg integration computing $R_{3,3}$.

$$f(0) = f(\pi) = 0.000000; \quad f\left(\frac{\pi}{2}\right) = 1.000000; \quad f\left(\frac{\pi}{4}\right) = f\left(\frac{3\pi}{4}\right) = 0.707107;$$

$$f\left(\frac{\pi}{8}\right) = f\left(\frac{7\pi}{8}\right) = 0.382683; \quad f\left(\frac{3\pi}{8}\right) = f\left(\frac{5\pi}{8}\right) = 0.923880.$$

$$T_1 = T(f, \{0, \pi\}) = \pi \left(\frac{1}{2} f(0) + \frac{1}{2} f(\pi) \right)$$

$$= \pi \left(\frac{1}{2} \cdot 0 + \frac{1}{2} \cdot 0 \right) = \pi \cdot 0.000000 = 0.000000$$

$$T_2 = T(f, \{0, \frac{\pi}{2}, \pi\}) = \pi \cdot \frac{1}{2} \left(\frac{1}{2} f(0) + f\left(\frac{\pi}{2}\right) + \frac{1}{2} f(\pi) \right)$$

$$= \pi \cdot \frac{1}{2} \left(\frac{1}{2} \cdot 0 + 1 + \frac{1}{2} \cdot 0 \right) = \pi \cdot 0.500000 = 1.570796$$

$$T_4 = T(f, \{0, \frac{\pi}{4}, \frac{\pi}{2}, \frac{3\pi}{4}, \pi\}) = \pi \cdot \frac{1}{4} \left(\frac{1}{2} f(0) + f\left(\frac{\pi}{4}\right) + f\left(\frac{\pi}{2}\right) + f\left(\frac{3\pi}{4}\right) + \frac{1}{2} f(\pi) \right)$$

$$= \pi \cdot \frac{1}{4} \left(\frac{1}{2} \cdot 0 + 0.707107 + 1 + 0.707107 + \frac{1}{2} \cdot 0 \right) = \pi \cdot 0.603553 = 1.896119$$

$$T_8 = T(f, \{0, \frac{\pi}{8}, \frac{\pi}{4}, \frac{3\pi}{8}, \frac{\pi}{2}, \frac{5\pi}{8}, \frac{3\pi}{4}, \frac{7\pi}{8}, \pi\}) = \frac{1}{2} \left(T_4 + \frac{\pi}{4} \left(f\left(\frac{\pi}{8}\right) + f\left(\frac{3\pi}{8}\right) + f\left(\frac{5\pi}{8}\right) + f\left(\frac{7\pi}{8}\right) \right) \right)$$

$$= \frac{1}{2} \left(1.896119 + \pi \cdot \frac{1}{4} (0.382683 + 0.923880 + 0.923880 + 0.382683) \right)$$

$$= \frac{1}{2} (1.896119 + \pi \cdot 0.653281) = \frac{1}{2} (1.896119 + 2.052344) = 1.974232$$

67 / 78

Romberg integration

Example Estimate $\int_0^\pi \sin x \, dx$ using Romberg integration computing $R_{3,3}$.

$$R_{0,0} = 0.000000; \quad R_{1,0} = 1.570796; \quad R_{2,0} = 1.896119; \quad R_{3,0} = 1.974232.$$

$$R_{1,1} = R_{1,0} + \frac{1}{3} (R_{1,0} - R_{0,0}) = \frac{1}{3} (4R_{1,0} - R_{0,0})$$

$$= \frac{1}{3} (4 \times 1.570796 - 0.000000) = 2.094395$$

$$R_{2,1} = \frac{1}{3} (4R_{2,0} - R_{1,0}) = \frac{1}{3} (4 \times 1.896119 - 1.570796) = 2.004560$$

$$R_{3,1} = \frac{1}{3} (4R_{3,0} - R_{2,0}) = \frac{1}{3} (4 \times 1.974232 - 1.896119) = 2.000269$$

$$R_{2,2} = R_{2,1} + \frac{1}{15} (R_{2,1} - R_{1,1}) = \frac{1}{15} (16R_{2,1} - R_{1,1})$$

$$= \frac{1}{15} (16 \times 2.004560 - 2.094395) = 1.998571$$

$$R_{3,2} = \frac{1}{15} (16R_{3,1} - R_{2,1}) = \frac{1}{15} (16 \times 2.000269 - 2.004560) = 1.999983$$

$$R_{3,3} = R_{3,2} + \frac{1}{63} (R_{3,2} - R_{2,2}) = \frac{1}{64} (64R_{3,2} - R_{2,2})$$

$$= \frac{1}{63} (64 \times 1.999983 - 1.998571) = 2.000006$$

68 / 78

Romberg integration

Example Estimate $\int_0^\pi \sin x \, dx$ using Romberg integration computing $R_{3,3}$.

Computed Romberg estimates:

$R_{0,0} = T_1 = 0.000000$			
$R_{1,0} = T_2 = 1.570796$	$R_{1,1} = 2.094395$		
$R_{2,0} = T_4 = 1.896119$	$R_{2,1} = 2.004560$	$R_{2,2} = 1.998571$	
$R_{3,0} = T_8 = 1.974232$	$R_{3,1} = 2.000269$	$R_{3,2} = 1.999983$	$R_{3,3} = 2.000006$

So $\int_0^\pi \sin x \, dx \approx R_{4,4} = 2.000006$.

Exact answer 2.000000, relative error $3 \times 10^{-6} = 0.0003\%$!

69 / 78

Romberg integration

Exercise Estimate $\int_0^1 x^2 dx$ by $R_{2,2}$.

Answer:

$$\begin{aligned}R_{0,0} &= 0.50000 \\R_{1,0} &= 0.37500 \quad R_{1,1} = 0.333333 \\R_{2,0} &= 0.34375 \quad R_{2,1} = 0.333333 \quad R_{2,2} = 0.333333\end{aligned}$$

Exercise Estimate $\int_1^2 1/x dx = \ln 2 \approx 0.693147$ by $R_{2,2}$.

Answer:

$$\begin{aligned}R_{0,0} &= 3/4 = 0.750000 \\R_{2,0} &= 17/24 = 0.708333 \\R_{2,0} &= 1171/1680 = 0.697024 \\R_{1,1} &= R_{1,0} + \frac{1}{3}(R_{1,0} - R_{0,0}) = 25/36 = 0.694444 \\R_{2,1} &= R_{1,0} + \frac{1}{3}(R_{1,0} - R_{0,0}) = 1747/2520 = 0.693254 \\R_{2,2} &= R_{2,1} + \frac{1}{15}(R_{2,1} - R_{1,1}) = 4367/6300 = 0.693175\end{aligned}$$

70 / 78

Adaptive Integration

71 / 78

Adaptive integration

Adaptive Methods In practise, we wish to estimate $\int_a^b f(x) dx$ to some fixed tolerance ϵ .

Suppose we can find an estimate $E(f, [p, q])$ for an integration method over the interval $[p, q]$.

Then the total error for a partition $P = \{x_0, x_1, \dots, x_n\}$ is approximately

$$\sum_{i=1}^n E(f, [x_{i-1}, x_i]).$$

We can *refine* the partition by splitting the subinterval with the largest error to obtain a (hopefully) better approximation.

Over a fixed subinterval $[p, q]$, the (average) error should be at most

$$\frac{q-p}{b-a} \epsilon$$

We can often estimate the error on subinterval $[p, q]$ using two evaluations of a simple integration scheme.

72 / 78

Adaptive trapezoid rule

Adaptive trapezoid rule Composite trapezoid rule; error $\frac{f''(\xi)(b-a)}{12}h^2$.

Assuming $f''(\xi)$ is approximately constant K , can estimate K using two evaluations of the trapezoid rule.

Recall $T_2(f, [a, b]) := T(f, [a, \frac{a+b}{2}, b]) = T(f, [a, \frac{a+b}{2}]) + T(f, [\frac{a+b}{2}, b])$, so

$$T(f, [a, b]) - I \approx \frac{f''(\xi)(b-a)}{12}h^2; \quad T_2(f, [a, b]) - I \approx \frac{f''(\tilde{\xi})(b-a)}{12}(h/2)^2.$$

Assuming $f''(\xi) \approx f''(\tilde{\xi}) \approx K$, we have

$$T_2(f, [a, b]) - I \approx \frac{K(b-a)}{12}(h/2)^2 \approx (T(f, [a, b]) - I)/4.$$

Multiplying through by 4, rearranging, and dividing by 3 gives

$$\frac{1}{3}(T_2(f, [a, b]) - T(f, [a, b])) \approx -(T_2(f, [a, b]) - I).$$

So we obtain error estimate on $[a, b]$ of

$$\begin{aligned} \epsilon &= |T_2(f, [a, b]) - I| \approx \frac{1}{3}|T(f, [a, \frac{a+b}{2}, b]) - T(f, [a, b])| \\ &= \frac{1}{12}(b-a)|f(a) - 2f(\frac{a+b}{2}) + f(b)|. \end{aligned}$$

73 / 78

Adaptive trapezoid rule

Adaptive trapezoid rule Approximate the integral of f over $[p, q]$ by

$$\begin{aligned} \int_p^q f(x) dx &\approx T(f, [p, \frac{p+q}{2}, q]) = T_2(f, [p, q]) \\ &= \frac{q-p}{4}(f(p) + 2f(\frac{p+q}{2}) + f(q)). \end{aligned}$$

Estimate the error of this approximation as

$$\begin{aligned} E_{T_2}(f, [p, q]) &\approx \frac{1}{3}|T(f, [p, \frac{p+q}{2}, q]) - T(f, [p, q])| \\ &= \frac{q-p}{12}(f(p) - 2f(\frac{p+q}{2}) + f(q)) \end{aligned}$$

Subdivide the interval with the largest estimated error until the total estimated error is less than ϵ .

Alternatively, subdivide each interval $[p, q]$ if

$$E_{T_2}(f, [p, q]) > \frac{q-p}{b-a}\epsilon.$$

74 / 78

Adaptive trapezoid rule

Example Evaluate $\int_0^2 1/(3+x^4) dx$ with error $\epsilon = 10^{-2}$.

$$T(f, [0, 2]) \stackrel{4\text{dp}}{=} 0.3860, \quad T(f, [0, 1]) + T(f, [1, 2]) \stackrel{4\text{dp}}{=} 0.2917 + 0.1513 = 0.4430.$$

$E_{T_2}(f, [0, 2]) = 0.0190$ is too high, need to split.

$[p, q]$	$T(f, [p, q])$	$T_2(f, [p, q]) = T(f, [p, \frac{p+q}{2}], q)$	$E_{T_2}(f, [p, q])$
$[0.0, 2.0]$	0.3860	0.4430	0.0190
$[0.0, 1.0]$	0.2917	0.3091	0.0058
$[1.0, 2.0]$	0.1513	0.1377	0.0045
$[0.0, 0.5]$	0.1650	0.1657	0.0002
$[0.5, 1.0]$	0.1441	0.1474	0.0011

So use trapezoid rule with partition $P = [0.0, 0.25, 0.5, 0.75, 1.0, 1.5, 2.0]$.

Total estimated error $0.0011 + 0.0002 + 0.0045 = 0.0058$ (4 dp).

Estimate of integral

$$T(f, P) = T(f, [0.0, 0.25, 0.5]) + T(f, [0.5, 0.75, 1.0]) + T(f, [1.0, 1.5, 2.0])$$

$$\stackrel{4\text{dp}}{=} 0.1657 + 0.1474 + 0.1377 = 0.4508.$$

Exact value 0.4486 (4 dp); error $2.2 \times 10^{-3} < 10^{-2}$.

75 / 78

Adaptive Simpson's rule (Non-examinable)

Adaptive Simpson's rule Composite Simpson's rule; error $\frac{f^{(4)}(\xi)(b-a)}{180}h^4$. Assuming $f^{(4)}(\xi)$ is approximately constant K , can estimate K using two evaluations of Simpson's rule. Let $h = (b-a)/2$.

$$S(f, [a, b]) - I \approx \frac{f^{(4)}(\xi)(b-a)}{180}h^4;$$

$$S(f, [a, c]) + S(f, [c, b]) - I \approx \frac{f^{(4)}(\tilde{\xi})(b-a)}{180}(h/2)^4.$$

Assuming $f^{(4)}(\xi) \approx f^{(4)}(\tilde{\xi}) \approx K$, we have

$$S(f, [a, c]) + S(f, [c, b]) - I \approx \frac{K(b-a)}{180}(h/2)^4 \approx (S(f, [a, b]) - I)/16$$

Multiplying through by 16 and rearranging gives

$$S(f, [a, c]) + S(f, [c, b]) - S(f, [a, b]) \approx -15(S(f, [a, c]) + S(f, [c, b]) - I)$$

So we obtain error estimate on $[a, b]$ of

$$\begin{aligned} \epsilon &= |S(f, [a, c]) + S(f, [c, b]) - I| \approx \frac{1}{15} |S(f, [a, c]) + S(f, [c, b]) - S(f, [a, b])| \\ &= \frac{1}{180} (b-a) (f(a) - 4f(a+h) + 6f(a+2h) - 4f(a+3h) + f(a+4h)) \end{aligned}$$

76 / 78

Adaptive Simpson's rule (Non-examinable)**Adaptive Simpson's rule** Approximate the integral of f over $[p, q]$ by

$$\int_p^q f(x) dx \approx S(f, [p, \frac{p+q}{2}]) + S(f, [\frac{p+q}{2}, q])$$

$$= \frac{q-p}{12} (f(p) + 4f(\frac{3p+q}{4}) + 2f(\frac{p+q}{2}) + 4f(\frac{p+3q}{4}) + f(q)).$$

Estimate the error of this approximation as

$$E_S(f, [p, q]) \approx \frac{1}{15} |S(f, [p, \frac{p+q}{2}]) + S(f, [\frac{p+q}{2}, q]) - S(f, p, q)|$$

$$= \frac{q-p}{180} (f(p) - 4f(\frac{3p+q}{4}) + 6f(\frac{p+q}{2}) - 4f(\frac{p+3q}{4}) + f(q))$$

Subdivide the interval $[p, q]$ if

$$E_S(f, [p, q]) > \frac{q-p}{b-a} \epsilon.$$

Alternatively, subdivide the interval with the largest estimated error until the total estimated error is less than ϵ .

77 / 78

Adaptive Simpson's rule (Non-examinable)**Example** Consider $\int_0^2 1/(1+x^2) dx$ with error $\epsilon = 10^{-4}$.Estimate $S(f, [0, 2]) = 1.066667$, $S(f, [0, 1]) + S(f, [1, 2]) = 1.105128$, $E_S(f, [0, 2]) = 0.0025$. Too high, need to split.

$[p, q]$	$S(f, [p, q])$	$S(f, [p, r]) + S(f, [r, q])$	$E_S(f, [p, q])$	$\frac{q-p}{b-a} \epsilon$
$[0.0, 2.0]$	1.066667	1.105128	2.5×10^{-3}	1.0×10^{-4}
$[0.0, 1.0]$	0.783333	0.785392	1.4×10^{-4}	5.0×10^{-5}
$[1.0, 2.0]$	0.321795	0.321748	3.1×10^{-6}	5.0×10^{-5}
$[0.0, 0.5]$	0.463725	0.46365	4.8×10^{-6}	2.5×10^{-5}
$[0.5, 1.0]$	0.321667	0.321745	5.3×10^{-6}	2.5×10^{-5}

So use Simpson's rule with partition

$$P = [0.0, 0.25, 0.5, 0.75, 1.0, 1.5, 2.0],$$

evaluation points

$$[0.0, 0.125, 0.25, 0.375, 0.5, 0.625, 0.75, 0.875, 1.0, 1.25, 1.5, 1.75, 2.0]$$

Estimate of integral 1.107146; exact value 1.107149; error 2.6×10^{-6} .

78 / 78