

Characterisation of the sparks (full-detector discharges) for detector 3

(June cosmic bench — `g_det3_wknd`)

MX17 cosmic-bench QA

July 5, 2026

1 Setup

A **spark** is an event in which the det3 readout fires more than **50 strips** at once (summed over both planes) — a full-detector discharge, the same cut used to veto sparks in the efficiency breakdown. Run `mx17_det3_p2_det1_overnight_6-27-26 / long_run_p2` (headline det3): 6.9 h, 90 914 detector-firing events, **8 301 sparks** (9.1 % of firing events, mean rate 0.33 Hz). Each spark hit carries a strip position (X=FEU7, Y=FEU8), a time within the readout window, and an amplitude; each event carries a timestamp and, where M3 reconstructs a clean single track, a ray projected onto the det3 plane in the aligned frame (same geometry as the efficiency analysis). We answer four questions.

2 Result in one paragraph

Sparks are **Poisson-random in time** (flat rate, no HV-recovery bursts) but **not random with respect to the muon**: conditioning on a clean single muon, a track that *crosses* the detector sparks at 4.9 % versus only 1.2 % when it clearly *misses* — a **4× enhancement**, so most in-detector sparks are muon-induced on top of a $\sim 1.2\%$ spontaneous baseline. In detector coordinates they are **localised**: strip occupancy peaks sharply at the plane **edges** (plus a few hot strips), while the discharge centroid sits centrally. Temporally the discharge is **neither a single flash nor a propagating “walk”**: strips across a ~ 200 mm region fire over a spread ~ 250 ns window (tail to ~ 1 μ s) with no position–time correlation.

3 Q1 — Are sparks random in time? Yes (Poisson)

The spark rate is flat at 0.33 Hz across the entire 6.9 h run (Fig. 1a): no bursts, no drift, so this is *not* an HV-instability / recovery-oscillation phenomenon. Inter-spark intervals follow a pure exponential over three decades (Fig. 1b) and the sparks-per-10-min counts have $\text{std} = 14 = \sqrt{\text{mean}}$ (Fig. 1c) — textbook Poisson. *Caveat*: Poisson timing is expected for *both* muon-induced and spontaneous sparks (muons themselves arrive Poisson), so time-randomness alone does not settle causation — that is what Q2 is for.

4 Q2 — Muon-induced or random? Muon-dominated

This is the decisive test (Fig. 2a). Taking only events with a clean single M3 muon and asking whether that muon’s projection lands on the detector: A muon that traverses the gas raises the spark probability **4.1×** over one that misses. So $\sim 75\%$ of in-detector sparks are **muon-induced** (the crossing muon’s primary ionisation seeds the discharge) and $\sim 25\%$ ($\approx 1.2\%$ absolute) are a




Figure 1: **Time.** (a) spark rate vs time in run; (b) inter-spark interval pdf with the exponential expectation; (c) counts per 10-min bin vs the Poisson $\pm\sqrt{\text{mean}}$ band. Sparks are random in time with a stable rate.

muon geometry (clean single M3 track)	N	sparks	spark rate
crosses detector (ray in active box)	52 995	2 600	4.91 %
at edge (<30 mm outside box)	9 824	557	5.67 %
clearly misses (>30 mm outside box)	11 858	141	1.19 %
(no clean M3 track — see below)	36 609	5 003	13.67 %

spontaneous baseline present even with no muon in the gas. In M3 coordinates the sparking muons are otherwise unremarkable — roughly uniformly spread over the acceptance (Fig. 2b), i.e. no special beam/telescope location.

The “no clean track” population (a readout confound). 60 % of all sparks fall in events where M3 finds *no* clean track, versus a 33 % baseline — sparks are $2\times$ over-represented there, and those sparks are the *same size* as the tracked ones (median 91 vs 95 strips). The most likely reading is that a large discharge injects common-mode noise into the *shared* DAQ and spoils M3 track reconstruction for that event (or that both are triggered by a messy multi-particle event). Either way it is a selection/cross-talk artefact, not evidence of a large muon-independent spark source, so the clean 4.9 % vs 1.2 % comparison above is restricted to good-track events. *This DAQ cross-talk is itself worth a follow-up.*




Figure 2: **Muon dependence.** (a) spark rate split by where the muon projects — crossing muons spark $4\times$ more than missing ones; (b) M3 crossing positions of sparking muons (roughly uniform); (c) discharge centroid in the detector frame (central).

5 Q3 — Are there particular places that spark? Yes: the edges

Strip occupancy in sparks is strongly non-uniform (Fig. 3a,b). Both planes show large peaks at the detector **edges** (0 and ~ 399 mm; strongest at high- X and low- Y), where field lines concentrate at the active-area boundary — edge strips participate in a large fraction of all sparks. On top of the edge structure the Y plane (FEU8) shows a handful of narrow spikes at fixed positions — individual **hot strips** that fire far above average. Yet the discharge *centroid* sits centrally (Fig. 2c): a typical spark ignites over a ~ 200 mm span (median X span 204 mm, Y span 235 mm of the 399 mm plane) — roughly half the detector, not a point and not the whole plane — with the edges over-weighted. The Y /FEU8 plane discharges harder than X (median 53 vs 40 strips; Fig. 3c), consistent with the FEU8 saturation band seen in the reco.far study. Only 10 % of spark hits actually saturate the ADC, so these are genuine spread discharges, not just electronic rail-outs.




Figure 3: **Where.** (a,b) strip occupancy in sparks for the X and Y planes vs a uniform baseline — edge peaks (both planes) and fixed hot strips (Y); (c) strips firing per spark — Y discharges are larger than X .

6 Q4 — All strips at once, or does the discharge “walk”? Neither

The discharge is *not* a single simultaneous flash: only 1.2 % of sparks have all their hits within 100 ns; the per-spark hit-time spread is $\sigma \approx 253$ ns median (tail beyond 500 ns), somewhat broader than a normal drift cluster (193 ns, Fig. 4a). But it does *not* propagate spatially either: the per-spark position–time correlation is $|r| \approx 0.12$ (only 3 % exceed 0.5), and the spark-centred $\Delta\text{position}-\Delta\text{time}$ map (Fig. 4b) is a clean **cross with no tilt** — a horizontal arm (strips across the whole span firing near-simultaneously) plus a vertical arm (the core keeps firing over $\pm$ hundreds of ns), and no diagonal that a moving front would produce. Example events (Fig. 4c) show strips lighting up across the full 0–400 mm range at scattered times. **Picture:** a spatially-extended discharge that fires a broad region in a near-synchronous burst and then rings/afterpulses for up to ~ 1 μ s, with timing independent of position — not a wave sweeping across the strips.

7 Summary and follow-ups

- **Time:** Poisson-random, stable 0.33 Hz — not HV bursting.
- **Cause:** predominantly **muon-induced** — a crossing muon sparks $4\times$ more than a missing one (4.9 % vs 1.2 %) — on a ~ 1.2 % spontaneous floor. Lowering the resist HV reduces the *rate* (fewer avalanches reach breakdown) but cannot remove the muon-seeded component entirely.




Figure 4: **Time structure.** (a) per-spark hit-time spread vs a normal cluster; (b) spark-centred time vs position (no tilt $\Rightarrow$ no propagating front); (c) two example sparks — strips across ~ 400 mm, spread in time, no position ramp.

- **Place:** **edge-dominated** occupancy plus fixed hot strips (Y/FEU8); discharges span $\sim$ half the plane, centred centrally; Y sparks harder than X.
- **Structure:** a spatially-extended, temporally-spread (~ 250 ns, tail to ~ 1 μ s) discharge with afterpulsing — no propagating “walk”.
- **Follow-ups:** (1) inspect the edge / guard-ring field and the Y/FEU8 hot strips (mask or lower gain); (2) confirm and mitigate the **DAQ common-mode cross-talk** by which a det3 spark spoils M3 tracking (60% of sparks lose the M3 track); (3) an offline spark tag (multiplicity + the ~ 1 μ s duration signature) would clean the reco_far tail (companion note `det3_recofar_analysis`).

Reproduce: `extract_sparks.py` then `make_plots.py`; numbers in `spark_meta.json`. Spark cut (>50 strips) and alignment match `09_efficiency_breakdown.py`.