



Math • Multivariable calculus • Applications of multivariable derivatives
• Quadratic approximations

The Hessian

The Hessian is a matrix that organizes all the second partial derivatives of a function.

Google Classroom Facebook Twitter Email

Background:

- [Second partial derivatives](#)

The Hessian matrix

The "**Hessian matrix**" of a multivariable function $f(x, y, z, \dots)$, which different authors write as $\mathbf{H}(f)$, $\mathbf{H}f$, or \mathbf{H}_f , organizes all second partial derivatives into a matrix:

Are you a student or a teacher?

[Student](#)

[Teacher](#)

$$\mathbf{H}f = \begin{bmatrix} \frac{\partial^2 f}{\partial x^2} & \frac{\partial^2 f}{\partial x \partial y} & \frac{\partial^2 f}{\partial x \partial z} & \cdots \\ \frac{\partial^2 f}{\partial y \partial x} & \frac{\partial^2 f}{\partial y^2} & \frac{\partial^2 f}{\partial y \partial z} & \cdots \\ \frac{\partial^2 f}{\partial z \partial x} & \frac{\partial^2 f}{\partial z \partial y} & \frac{\partial^2 f}{\partial z^2} & \cdots \\ \vdots & \vdots & \vdots & \ddots \end{bmatrix}$$

So, two things to notice here:

- This only makes sense for scalar-valued function.
- This object $\mathbf{H}f$ is no ordinary matrix; it is a matrix with *functions* as entries. In other words, it is meant to be evaluated at some point (x_0, y_0, \dots) .

Are you a student or a teacher?

Student

Teacher

$$\mathbf{H}f(x_0, y_0, \dots) = \begin{bmatrix} \frac{\partial^2 f}{\partial x^2}(x_0, y_0, \dots) & \frac{\partial^2 f}{\partial x \partial y}(x_0, y_0, \dots) & \cdots \\ \frac{\partial^2 f}{\partial y \partial x}(x_0, y_0, \dots) & \frac{\partial^2 f}{\partial y^2}(x_0, y_0, \dots) & \cdots \\ \vdots & \vdots & \ddots \end{bmatrix}$$

As such, you might call this object $\mathbf{H}f$ a "matrix-valued" function. Funky, right?

[The nature of higher order derivatives in a multivariable world]

One more important thing, the word "Hessian" also sometimes refers to the **determinant of this matrix**, instead of to the matrix itself.

Example: Computing a Hessian

Problem: Compute the Hessian of $f(x, y) = x^3$

Solution: Ultimately we need all the second partial derivatives, so we first compute both partial derivatives:

Are you a student or a teacher?

Student

Teacher

$$f_x(x, y) = \frac{\partial}{\partial x}(x^3 - 2xy - y^6) = 3x^2 - 2y$$

$$f_y(x, y) = \frac{\partial}{\partial y}(x^3 - 2xy - y^6) = -2x - 6y^5$$

With these, we compute all four second partial derivatives:

$$f_{xx}(x, y) = \frac{\partial}{\partial x}(3x^2 - 2y) = 6x$$

$$f_{xy}(x, y) = \frac{\partial}{\partial y}(3x^2 - 2y) = -2$$

$$f_{yx}(x, y) = \frac{\partial}{\partial x}(-2x - 6y^5) = -2$$

$$f_{yy}(x, y) = \frac{\partial}{\partial y}(-2x - 6y^5) = -30y^4$$

The Hessian matrix in this case is a 2×2 matrix with the following entries:

Math > Multivariable
calculus > Applications of
multivariable derivatives >
Quadratic approximations
Quadratic approximations



Practice: The Hessian
matrix

Are you a student or a
teacher?

Student

Teacher



Expressing a quadratic form with a matrix



Vector form of multivariable quadratic approximation



The Hessian



Quadratic approximation

Next lesson

Optimizing multivariable f...

$$\mathbf{H}f(x, y) = \begin{bmatrix} f_{xx}(x, y) & f_{yx}(x, y) \\ f_{xy}(x, y) & f_{yy}(x, y) \end{bmatrix} = \begin{bmatrix} 6x & -2 \\ -2 & -30y^4 \end{bmatrix}$$

We were asked to evaluate this at the point $(x, y) = (1, 2)$, so we plug in these values:

$$\mathbf{H}f(1, 2) = \begin{bmatrix} 6(1) & -2 \\ -2 & -30(2)^4 \end{bmatrix} = \begin{bmatrix} 6 & -2 \\ -2 & -480 \end{bmatrix}$$

Now, the problem is ambiguous, since the "Hessian" can refer either to this matrix or to its determinant. What you want depends on context. For example, in optimizing multivariable functions, there is something called the "second partial derivative test" which uses the Hessian determinant. When the Hessian is used to approximate functions, you just use the matrix itself.

If it's the determinant we want, here's what we get:

$$\det \left(\begin{bmatrix} 6 & -2 \\ -2 & -480 \end{bmatrix} \right) = 6(-480) - (-2)(-2) = -2884$$

Uses

Are you a student or a teacher?

Student

Teacher

By capturing all the second-derivative information of a multivariable function, the Hessian matrix often plays a role analogous to the ordinary second derivative in single variable calculus. Most notably, it arises in these two cases:

- [Quadratic approximations of multivariable functions](#), which is a bit like a second order Taylor expansion, but for multivariable functions.
- [The second partial derivative test](#), which helps you find the maximum/minimum of a multivariable function.

Sort by:

Top Voted

[Questions](#)[Tips & Thanks](#)

Want to join the conversation?

[Log in](#)

Are you a student or a teacher?

[Student](#)[Teacher](#)

Jorge R. Martinez Perez-T... 6 years ago

will there be videos and exercises (mostly interested in the exercises) for these topics any time soon?

Reply • Comment

(28 votes)  Upvote  Downvote  Flag [more](#) 



arkanmanva 5 years ago



[more](#) 

Me too. Would love to see exercises in multivariable calculus, differential equations and linear algebra.

Comment

(20 votes)  Upvote  Downvote  Flag [more](#) 

See 1 more reply



Marcel Brown 6 years ago

[more](#) 

Should the determinant in the final step be: $180xy^4 - 4$?

Reply • Comment

(7 votes)  Upvote  Downvote  Flag [more](#) 



Kenz 6 years ago



[more](#) 

I agree partially with Marcel Brown; the determinant is calculated as a 2x2 matrix by $ad-bc$, in this form because $a=180xy^4$ and $b=-4$. However, $ab \cdot \text{coefficient} = 6 \cdot -30 = -180$.

Comment

(12 votes) 

See 1 more reply

Are you a student or a teacher?

[Student](#)

[Teacher](#)



Maria 6 years ago

more ▾

Is the Hessian in any way related to the Jacobian matrix?

Reply • 1 comment

(5 votes)



Upvote



Downvote



Flag

more ▾



Jo Marino 5 years ago

more ▾

More formally: $\mathbf{H}(f(\mathbf{x})) = \mathbf{J}(\nabla f(\mathbf{x}))^T$.

It also relates to the Laplacian as an operator: $\Delta f = \nabla^2 f = \text{trace}(\mathbf{H}(f))$.

Comment

(4 votes)



Upvote



Downvote



Flag

more ▾



gschex1112 6 years ago

more ▾

Why is the last second partial derivative not $-30y^4$?

Reply • Comment

(5 votes)



Upvote



Downvote



Flag

more ▾



Emily H 6 years ago

more ▾

Why is $f_{yx} = d/dy(3x^2 - 2y)$ and not $d/dx(-2x - 6y^5)$? Wouldn't it be the partial derivative with respect to x of the first partial derivative with respect to y ? I ask the same for $f_{xy} = d/dx(-2x - 6y^5)$ not being $d/dy(3x^2 - 2y)$.

Reply • Comment

(3 votes)



Are you a student or a teacher?

Student

Teacher



Tejas 6 years ago

It is both! Whether you derivate with respect to x first then y , or with respect to y first then x , you get the same answer. Notice here that $f_{xy} = f_{yx} = -2$. That is Clairaut's theorem.

Comment

(3 votes)  Upvote  Downvote  Flag [more](#) 



yogesh kumar 6 years ago

[more](#) 

why hessian makes sense only for a scalar valued function?

Reply • 1 comment

(3 votes)  Upvote  Downvote  Flag [more](#) 



Aaron Hargrove 6 years ago

[more](#) 

What are some of the practical applications of the determinant of a Hessian matrix?

Reply • Comment

(2 votes)  Upvote  Downvote  Flag [more](#) 



Tejas 6 years ago

[more](#) 

Evaluating it can tell you whether you are at a maximum, minimum, or a saddle point. It has all the same abilities as a second derivative in a uni-variate function.

Comment

(2 votes) 



Briahnna A 6 years ago

Around the last paragraph, the determinant is
But, I was wonder why, in my multivariate cal

Are you a student or a teacher?

[Student](#)

[Teacher](#)

hessian determinant and f_{xx} to determine local extrema. I was wondering what was the background or the basis for using the determinant of the Hessian Matrix to decide if critical points are local maximum or local minimums? Basically, what is important about the Hessian Determinant? Thank-You!

Reply • Comment

(3 votes)  Upvote  Downvote  Flag [more](#) 



Tejas 6 years ago

[more](#) 

We actually don't use the Hessian to determine whether the critical points are local maxima or local minima. We actually use the Hessian to determine whether they are local extrema or saddle points. As for using f_{xx} , it doesn't have to be f_{xx} . You could just as easily use f_{yy} to determine whether the local extremum is a maximum or minimum.

If it is a local minimum, the gradient is pointing away from this point. If it is a local maximum, the gradient is always pointing toward this point. Of course, at all critical points, the gradient is 0. That should mean that the gradient of nearby points would be tangent to the change in the gradient. In other words, f_{xx} and f_{yy} would be high and f_{xy} and f_{yx} would be low.

On the other hand, if the point is a saddle point, then the gradient vectors will all be pointing around the critical point. Therefore at nearby points, the change in the gradient will be orthogonal to the gradient, not tangent. In other words, f_{xy} and f_{yx} would be high and f_{xx} and f_{yy} would be low, or f_{xx} and f_{yy} would be high and f_{xy} and f_{yx} would be low. The Hessian determinant would be negative.

Comment

(0 votes) 



disha.bandy 4 years ago

Are you a student or a teacher?


[Student](#)

[Teacher](#)

Since Hessian matrices are only used for trigonometric functions, what can be used to optimise a 3 variable trigonometric function?

Reply • Comment

(1 vote)  Upvote  Downvote  Flag [more](#) 



muhammadahme2019 2 years ago

[more](#) 

Where it says The Hessian matrix in this case is a 2×2 matrix with these functions as entries:
the f_{xx} and f_{xy} are flipped. Cuz under f_{xx} it should be f_{xy} . Then under f_{yx} it should be f_{yy} .

Reply • 1 comment

(1 vote)  Upvote  Downvote  Flag [more](#) 

Show more...

[< Vector form of multivariable quadratic approximation](#)

[Quadratic approximation >](#)

Are you a student or a teacher?

[Student](#)

[Teacher](#)