

Regularization

Johanni Brea

Introduction à l'apprentissage automatique

GYMINF 2021

Table of Contents

1. When Linear Models Are Too Flexible

2. Ridge Regression and the Lasso

3. Regularization Examples

When Linear Models Are Too Flexible

In the old days

Typically $n > p$ (much more data than predictors)


For example: predict blood pressure based on age, gender and body mass index (BMI)
(e.g. $n = 200$ patients, $p = 3$).

Nowadays: Big Data

Often $n \approx p$ or $n < p$

For example: predict blood pressure based on
500 000 single nucleotide polymorphisms (SNP)
($n = 200$, $p = 500\,000$).

⇒ **Linear Model perfectly fits the training data.**



Making Linear Models Less Flexible

Option 1: Fix some parameters at zero

$$\hat{y} = f(x) = f(x_1, x_2, \dots, x_p) = \beta_0 + \underbrace{\beta_1}_{\neq 0} x_1 + \underbrace{\beta_2}_{\neq 0} x_2 + \beta_3 x_3 + \dots + \underbrace{\beta_{p-1}}_{\neq 0} x_{p-1} + \beta_p x_p$$

Problem: Many different models to fit; $\binom{p+1}{m}$ combinations of m non-fixed parameters.

Option 2: Favor small parameters

Replace the original loss $L(\theta)$ by $L(\theta) + \text{"penalty for large parameters"}$

Table of Contents

1. When Linear Models Are Too Flexible

2. Ridge Regression and the Lasso

3. Regularization Examples

Ridge Regression (L2 Regularization)

$$L_{L2}(\theta) = L(\theta) + \lambda \|\theta\|_2^2$$

with **regularization constant** λ and (squared) **L2 norm** $\|\theta\|_2^2 = \sum_{i=1}^p \theta_i^2$.

1. The regularization constant λ is a hyper-parameter.
2. Often the intercept θ_0 is not regularized.
3. If $\lambda = 0$: original loss (no penalty)
4. The larger λ , the stronger the impact of the penalty on the result.
5. With increasing λ the model becomes less flexible.
6. With increasing λ all parameters tend to zero; it happens rarely that one is exactly zero.

Lasso (L1 Regularization)

$$L_{L1}(\theta) = L(\theta) + \lambda \|\theta\|_1$$

with **regularization constant** λ and **L1 norm** $\|\theta\|_1 = \sum_{i=1}^p |\theta_i|$.

Points 1-5 from ridge regression are also valid for the Lasso. However:

6. With large λ some parameters are exactly zero (in contrast to ridge regression).

An Alternative Formulation of Regularization

Thanks to a result from constraint optimization (see Karush-Kuhn-Tucker conditions, a generalization of Lagrange multipliers) the above formulations of Ridge Regression and the Lasso are equivalent to a constraint optimization problem:

Ridge Regression

minimize $L(\theta)$ under the constraint that $\|\theta\|_2^2 \leq S$.

The parameters are confined to a p -ball of radius S with center at the origin.

Lasso

minimize $L(\theta)$ under the constraint that $\|\theta\|_1 \leq S$.

The parameters are confined to a hypercube with edge length S , center at the origin and corners on the axes.

S is a (complicated) function of λ and the original loss $L(\theta)$.

With increasing S the model becomes more flexible.

Analytical Solutions for Simple Linear Regression

Notation: $\langle x \rangle = \frac{1}{n} \sum_{i=1}^n x_i$

Ridge Regression

$$L(\theta, \lambda) = \langle (y - \theta_0 - \theta_1 x)^2 \rangle + \lambda \theta_1^2$$

$$\theta_1 = \frac{\langle xy \rangle - \langle x \rangle \langle y \rangle}{\langle x \rangle^2 - \langle x^2 \rangle + \lambda}, \quad \theta_0 = \langle y \rangle - \theta_1 \langle x \rangle$$

Lasso

$$L(\theta, \lambda) = \frac{1}{2} \langle (y - \theta_0 - \theta_1 x)^2 \rangle + \lambda |\theta_1|$$

$$\theta_1 = \frac{\langle xy \rangle - \langle x \rangle \langle y \rangle - \text{sign}(\theta_1) \lambda}{\langle x \rangle^2 - \langle x^2 \rangle} \text{ or } 0 \text{ if } |\langle xy \rangle - \langle x \rangle \langle y \rangle| < \lambda$$

Standardized Inputs for Regularization

Problem

Assume we find in multiple linear regression on the weather data the following parameters

$$\begin{array}{lll} X_1 & \text{LUZ_pressure} & [\text{hPa}] \\ X_2 & \text{LUZ_temperature} & [^\circ\text{C}] \end{array} \left| \begin{array}{ll} \theta_1 = -1 & [\text{km/h/hPa}] \\ \theta_2 = 0.5 & [\text{km/h/}^\circ\text{C}] \end{array} \right.$$

We could have measured the pressure in Pa and get the equivalent result

$$\begin{array}{lll} X_1 & \text{LUZ_pressure} & [\text{Pa}] \\ X_2 & \text{LUZ_temperature} & [^\circ\text{C}] \end{array} \left| \begin{array}{ll} \theta_1 = -1/100 & [\text{km/h/Pa}] \\ \theta_2 = 0.5 & [\text{km/h/}^\circ\text{C}] \end{array} \right.$$

With regularization $\lambda(\theta_1^2 + \theta_2^2)$ we would get different results for measurements in hPa and in Pa, because θ_1 contributes less to the penalty in the latter case.

Solution

Standardize all predictors, such that they have variance 1:

$$\tilde{X}_i = X_i / \sqrt{\text{Var}(X_i)}$$

Quiz

- ▶ The Lasso tends to have larger variance but smaller bias than linear regression.
- ▶ Indicate which is correct: as we increase S from 0 to ∞ in L2 regularized linear regression the training error will be
 1. an inverted U shape.
 2. a U shape.
 3. steadily increasing.
 4. steadily decreasing.
 5. remain constant.
- ▶ Indicate which is correct: as we increase S from 0 to ∞ in L2 regularized linear regression the test error will be
 1. an inverted U shape.
 2. a U shape.
 3. steadily increasing.
 4. steadily decreasing.
 5. remain constant.

Table of Contents


1. When Linear Models Are Too Flexible

2. Ridge Regression and the Lasso


3. Regularization Examples

Polynomial Ridge Regression

$d = 20, \lambda = 0$



$d = 20, \lambda = 10^{-4}$



With a little bit of L2 regularization ($\lambda = 10^{-4}$)
one can prevent overfitting of polynomials with high degrees.

Multiple Logistic Ridge Regression on the Spam Data

$n = 2000$ emails, $p = 801$ features (size of the lexicon)

Without regularization

training misclassification rate: 0.0015


test misclassification rate: 0.1005

With L2 regularization

training misclassification rate: 0.0075

test misclassification rate: 0.0435

The Lasso Path for the Weather Data



As we lower λ , `BER_wind_peak` is the first non-zero factor, `BAS_wind_peak` the second and `LUZ_wind_mean` the third.

Summary

- ▶ Regularization allows to lower the flexibility of a model by restricting the parameters to certain areas of the parameter space.
- ▶ L1 regularization leads to sparse models with some parameters exactly zero
⇒ great for interpretability.